

**Expression of the G<sub>i</sub>-coupled RASSL Ro1 in GFAP-positive cells: a novel model of hydrocephalus**

Elizabeth Joy Sweger

A dissertation submitted to the faculty of the University of North Carolina at Chapel Hill in partial fulfillment of the requirements for the degree of Doctor of Philosophy in the Department of Pharmacology.

Chapel Hill  
2009

Approved by:

Ken McCarthy

Robert Rosenberg

Gary Johnson

Ben Philpot

Patricia Maness

## **ABSTRACT**

ELIZABETH SWEGER: Expression of the Gi-coupled RASSL Ro1 in GFAP-positive cells:  
a novel model of hydrocephalus  
(Under the direction of Ken McCarthy)

Hydrocephalus is a disorder of cerebrospinal fluid dynamics within the central nervous system. Although symptoms of hydrocephalus are often relieved by shunt implantation, complications are common and underlying pathology remains. Understanding the molecular mechanisms of hydrocephalus is critical for development of new therapeutics; however few experimental models allow early events to be studied. Originally designed to study astrocytic G<sub>i</sub> signaling, we developed a transgenic mouse line expressing the G<sub>i</sub>-coupled RASSL (receptor activated solely by synthetic ligand) Ro1 in GFAP-positive cells by crossing the hGFAP-tTA (tet transactivator behind the human glial fibrillary acidic protein promoter) mouse line with the tetO-Ro1/tetO-LacZ mouse line. Surprisingly, we found that all double-transgenic mice developed hydrocephalus by postnatal day 15, while single-transgenic littermate controls appeared normal.

Hydrocephalic Ro1 mice had enlarged lateral and third ventricles, thinned cortex, partial denudation of the ependymal cell layer, abnormal subcommissural organ, and obliteration of the cerebral aqueduct. Severely affected mice had increased phosphoErk and GFAP expression. Giving breeding pairs doxycycline prevented the expression of Ro1 and onset of hydrocephalus in double-transgenic offspring. Double transgenic mice taken off dox at weaning developed enlarged ventricles within 7 weeks, indicating that Ro1 expression also induces hydrocephalus in adults. Double-transgenic mice injected with a Ro1 inverse agonist



when taken off dox did not develop enlarged ventricles or have ependymal detachment, demonstrating that signaling through Ro1 is required for hydrocephalus. We have discovered a new model that allows onset of hydrocephalus to be controlled, providing the unique ability to study the earliest events in juvenile and adult-onset hydrocephalus.

Ependymal denudation is a common feature of hydrocephalus that appears to occur early in the disease. Positive GFAP and Ro1 staining was detected in a subset of ependymal cells from double transgenic mice off dox. Timed studies showed that Ro1 expression began within three days of dox removal and ependymal denudation began within ten days. Affymetrix gene arrays were utilized to screen for changes in ependymal gene expression at five and nine days after dox removal. No differentially expressed genes were detected between double-transgenic mice and controls at either timepoint; RT-PCR also failed to detect significant changes.

## **ACKNOWLEDGEMENTS**

Few difficult undertakings are accomplished without the support and encouragement of others. I am deeply grateful to Ken McCarthy for taking a chance on a student with little research background and for providing advice, discipline and space to grow. I would not have begun this undertaking without the love of learning and the importance of education instilled in me by my parents, Dan and Carol. They demonstrated that anything worth doing should be done well, even if doing so requires sacrifice. My sister Judy, brother Daniel, and fiancé Matt have been an unwavering source of support and my personal cheerleaders. I am also thankful for the friendship and support of my lab mates, past and current. Many thanks go to my committee members for their advice and input, and to Kathy Justice for taking care of the details. Finally, all credit goes to God, who makes all things possible.

## TABLE OF CONTENTS

LIST OF TABLES.....	viii
LIST OF FIGURES.....	ix
LIST OF ABBREVIATIONS.....	xi
Chapter	
I. INTRODUCTION.....	1
Hydrocephalus.....	1
Anatomy of the ventricular system.....	2
Functions of cerebrospinal fluid.....	8
Cerebrospinal fluid production.....	11
Cerebrospinal fluid absorption.....	17
II. HYDROCEPAHLUS: CLASSIFICATION, CAUSES, PATHOLOGY, AND TREATMENT.....	23
Classification.....	23
Causes of hydrocephalus.....	27
Molecular mechanisms of hydrocephalus.....	30
Pathology of hydrocephalus.....	46
Treatment of hydrocephalus.....	60
Experimental hydrocephalus models.....	67
III. A NEW GENETIC MODEL OF HYDROCEPAHLUS.....	74

Materials and methods.....	75
Animals.....	75
Transgenic mouse lines.....	75
Genotyping.....	76
Southern blots.....	76
Doxycycline administration.....	76
Spiradoline administration.....	77
<i>Nor</i> Binaltorphimine administration.....	77
Immunofluorescence.....	77
Ependymal cell dissection.....	78
Xgal histochemistry.....	78
Immunoprecipitation and Western blotting.....	79
Hematoxylin and eosin staining.....	80
Immuncocytochemistry.....	80
Ventricle size.....	81
RNA purification.....	81
Gene microarrays.....	82
Microarray data analysis.....	83
Real time PCR.....	83
Astrocyte cell culture.....	84
cAMP assay.....	84

IV.	CHARACTERIZATION OF THE Ro1 HYDROCEPHALUS MODEL.....	85
	Results.....	89
	Discussion.....	97
V.	EPENDYMAL CELL GENOMIC ARRAYS.....	103
	Results.....	104
	Discussion.....	109
VI.	DISCUSSION.....	112
	Targeting a G <sub>i</sub> -coupled receptor to astrocytes.....	113
	Ro1 expression and cellular localization.....	113
	G <sub>i</sub> signaling and G <sub>i</sub> modulation of behavior.....	114
	G <sub>i</sub> modulation of neuronal excitability.....	115
	Mice expressing Ro1 develop hydrocephalus.....	116
	Doxycycline dosing.....	117
	The Ro1 model.....	117
	Ro1 signaling.....	119
	Possible mechanisms.....	120
	Gene arrays.....	122
	Alternate hypothesis.....	124
	Future directions.....	125
	LITERATURE CITED.....	128

## LIST OF TABLES

### Table

1. Severity of hydrocephalus phenotype in P15 Ro1 mice.....92
2. Gene targets for RT-PCR validation.....110

## LIST OF FIGURES

### Figure

1. The ventricular system.....	3
2. The subarachnoid space.....	4
3. Choroid plexus epithelium and the production of cerebrospinal fluid.....	15
4. Classic view of cerebrospinal fluid circulation.....	19
5. The Ro1 receptor.....	85
6. Tetracycline inducible system.....	86
7. Double transgenic mice off dox develop hydrocephalus.....	87
8. Ro1 expression is restricted to astrocytes and is regulated by dox.....	88
9. Ro1 expression is controlled by doxycycline.....	89
10. LacZ expression is regulated by dox.....	91
11. Ro1 mice maintained off dox develop hydrocephalus.....	91
12. Ro1 mice develop progressive enlargement of the lateral ventricles.....	92
13. Alterations in the ventricular system of hydrocephalic Ro1 mice.....	93
14. Ro1 mice will develop enlarged ventricles if taken off dox at weaning.....	94
15. Blocking Ro1 intrinsic activity prevents ventricle enlargement.....	95
16. Blocking Ro1 signaling prevents ependymal denudation.....	96
17. GFAP expression is upregulated in hydrocephalic Ro1 mice.....	97
18. Phospho-Erk p44/42 levels are elevated in hydrocephalic Ro1 mice.....	98
19. GFAP is expressed in ciliated ependymal cells.....	103
20. Isolated sheets of ependymal cells express GFAP and S100 $\beta$ .....	104

21. Isolated sheets of ependymal cells express GFAP and $\alpha$ -tubulin.....	105
22. Ro1 is expressed in ependymal cells.....	106
23. Time course of Ro1 expression after dox removal.....	106
24. Time course of ependymal denudation after dox removal.....	107



## LIST OF ABBREVIATIONS

ADNF	Activity-dependent neurotrophic factor
BBB	Blood-brain barrier
CNS	Central nervous system
CP	Choroid plexus
CSF	Cerebrospinal fluid
CT	Computed tomography
EC	Ependymal cells
ETV	Endoscopic third ventriculostomy
GFAP	Glial fibrillary acidic protein
GPCR	G protein coupled receptor
HC	Hydrocephalus
ICP	Intracranial pressure
IVH	Intraventricular hemorrhage
KOR	Kappa opioid receptor
LV	Lateral ventricle
MRI	Magnetic resonance imaging
NMDA	N-methyl-D-aspartate
<i>norBNI</i>	<i>NorBinaltorphimine</i>
RASSL	Receptor activated solely by synthetic ligand
SAS	Subarachnoid space
TGF $\beta$	Transforming growth factor $\beta$

VIP                      Vasoactive intestinal peptide

## CHAPTER 1

### INTRODUCTION

**Hydrocephalus.** In the United States approximately 1 in 1000 children are born with the neurological disorder hydrocephalus, sometimes commonly referred to as “water on the brain” (Leech and Payne, 1991; Fletcher et al., 1992a; Johanson and Jones, 2001). Children with hydrocephalus (HC) have an imbalance between the production and absorption of cerebrospinal fluid (CSF), resulting in an accumulation of CSF in the ventricles within the brain. If left untreated, the buildup of fluid causes the cranium to expand in an attempt to accommodate the extra fluid and pressure within the brain increases, resulting in disruption and compression of cerebral structures. More than 50% of children with untreated HC die within 10 years, and over 60% of surviving children have neurological impairments. The development of a one-way valve system to drain, or shunt, excess CSF from the brain was developed in the 1950’s and was the first breakthrough in HC treatment, greatly improving outcomes for HC patients (Laurence and Coates, 1962; Foltz and Shurtleff, 1963; Becker and Nulsen, 1968). While there have been advances in shunt technology, complications associated with shunt dependence and shunt failure frequently require multiple revision surgeries, making shunting a less than ideal treatment. To date there is no cure for HC, nor is there an effective pharmacological treatment for HC. The discovery of new therapies for HC is complicated by the multifactorial etiology of HC, and the molecular mechanisms leading the


onset of HC remain poorly understood. Future advances in HC treatment will be greatly aided by experimental models in which the earliest events of the disease can be studied.

We developed a transgenic mouse line expressing the Gi-coupled RASSL (Receptor Activated Solely by Synthetic Ligand) Ro1 in GFAP-positive cells with the aim of studying the role of GPCR signaling in astrocyte-neuronal communication. Expression of Ro1 was restricted to glia by crossing the transgenic hGFAP-tTA (tet transactivator behind the human glial fibrillary acidic protein promoter) mouse line with the transgenic tetO-Ro1/tetO-LacZ mouse line. This cross produced double transgenic mice that expressed Ro1 only in glia. Surprisingly, we found that all transgenic mice expressing Ro1 reliably developed hydrocephalus. We analyzed these mice in an effort to develop a new model of hydrocephalus that will further our understanding of the pathophysiology of the disease. An overview of the ventricular system, cerebrospinal fluid formation and function, and a review of the literature on HC will aid in understanding the Ro1 model.

**Anatomy of the ventricular system.** The central nervous system (CNS) is bathed and perfused in extracellular fluids that have critical roles in maintaining optimal conditions for brain activity. There are three fluid compartments in the CNS: the ventricles, the subarachnoid space, and the interstitial space. The ventricles and subarachnoid space are filled with cerebrospinal fluid (CSF). The fluid filling the interstitial space is very similar to CSF and is simply called interstitial fluid.

The ventricular system is composed of the lateral ventricles (ventricles one and two), the third ventricle and the fourth ventricle (Figure 1). The ventricles of adult humans hold approximately 30 mLs of CSF (Brodelt and Stoodley, 2007). The choroid plexi, specialized


tissue responsible for most nascent CSF production, are found in each of the ventricles. The lateral ventricles, the largest of the ventricles, have five distinct regions: the anterior (frontal) horn, body, posterior (occipital) horn, temporal (inferior) horn and the collateral trigone or atrium. The lateral ventricles (LV) are separated from each other by the septum and are each



connected to the mid-line third ventricle via the interventricular foramen of Monro. The third ventricle is irregularly shaped and lies between the thalami and above the hypothalamus. Four recesses project from the third ventricle. The optic and infundibular processes are located rostrally below the LV, while the pineal and supra-pineal recesses are located caudally. The third ventricle is connected to the fourth ventricle by the cerebral aqueduct of Sylvius. The aqueduct is the narrowest part of the ventricular system and is thus most susceptible to obstruction in CSF disorders. The fourth ventricle is triangular in shape and is located at the base of the brain ventral to the cerebellum. The median eminence, one

of the circumventricular organs, lies on the floor of the fourth ventricle. There are three foramina in the roof of the fourth ventricle that connect the fourth ventricle to the subarachnoid space. The medial foramen of Magendie is located caudally and opens into the cisterna magna. The two lateral foramina of Luschka connect the fourth ventricle to basal cisterns (Johanson, 2003; McLone, 2004; Di Terlizzi and Platt, 2006).

The subarachnoid space surrounds the brain and spinal cord and contains roughly four times more CSF than the ventricular system. The CNS is covered by three membranes collectively called the meninges (Figure 2). The dura mater is adherent to the skull and contains nerves, blood vessels and venous sinuses. The pia mater tightly covers the external surface of the nervous tissue on top of the superficial glial limiting layer, and is sometimes referred to as the pia-glia. The avascular arachnoid membrane lies in between the dura and pia and bridges the sulci, or grooves, on the surface of the brain. The resulting space



**Figure 2. The subarachnoid space.** Reprinted with permission [www.sbsdefense.com](http://www.sbsdefense.com).

between the pia and arachnoid membrane is the subarachnoid space and is filled with CSF. A loose mesh of fibroblasts and collagen, called the arachnoid trabeculae, connects the arachnoid membrane to the pia and may have a role in directing CSF flow (Yasargil et al., 1984; Segal, 1996; Johnston and Teo, 2000; Brodbelt and Stoodley, 2007). Major blood vessels also lie within the subarachnoid space, and are sheathed by a cuff of pia and arachnoid membrane as they penetrate into the brain. CSF also fills these spaces, called Virchow-Robin spaces (Johanson, 2003).

The interstitial fluid immediately surrounds the neurons and glia and is separated from the CSF in the subarachnoid space by the pia and from ventricular CSF by the ependymal layer lining the ventricles. The interstitial space is the largest of the three fluid compartments, holding roughly 280 mL of fluid in the normal human adult. CSF and interstitial fluid are in almost free communication across the interfaces separating them, and thus impact the composition of one another (Johanson, 2003; Brodbelt and Stoodley, 2007).

CSF transports nutrients and other critical substances from the blood to the central nervous system. Because neurons and glia require a precise extracellular fluid composition for optimal functioning, the composition of CSF needs to be tightly regulated. The blood-brain barrier (BBB) and blood-CSF barrier control the exchange of fluid, ions and proteins between the blood and CSF or interstitial fluid. Most blood vessels in the brain are surrounded by endothelial cells tightly joined by zonula occludens (tight junctions), forming the BBB that severely restrict most solutes from diffusing out of the blood. However, in several specialized areas, including the choroid plexus and median eminence, the blood vessel endothelial layers lack tight junctions, allowing a variety of substances to diffuse from the blood into the interstitial space. In these areas the epithelial layer (choroid plexus) or the

ependymal layer (median eminence) has tight junctions on their apical surface, forming a blood-CSF barrier to protect the brain while allowing these organs to carry out their specialized functions. The arachnoid membrane also has tight junctions to prevent CSF from leaking freely into the dural sinuses (Miller and Leslie, 1994; Di Terlizzi and Platt, 2006; Brodbelt and Stoodley, 2007).

The ependymal cell layer that lines the ventricles separates CSF from interstitial fluid, also acting as a CSF-brain barrier. Although ependymal cells (EC) are joined by zonula adherens and gap junctions, allowing CSF and interstitial fluid to readily communicate across the ependyma, the EC layer has features that allow it to serve as a protective barrier. Ependyma are equipped to remove and/or prevent toxic substances from crossing into the nervous tissue (Del Bigio, 1995; Strazielle and Gherzi-Egea, 2000). Metallothionein in EC takes up copper, zinc and heavy metals from the CSF while transferrin binds up iron (Blaauwgeers et al., 1993; Moos and Mollgard, 1993; Suzuki et al., 1994). Removal of copper and iron may protect the brain from oxidative damage since they are involved in ROS (reactive oxygen species) formation (Halliwell, 1992). Ependymal cells have been shown to sequester proteins, tracers, dyes and even latex beads via pinocytosis (Brightman, 1965; Booz and Wiesen, 1976; Broadwell and Sofroniew, 1993), proving they can take up various potentially harmful substances from CSF. Pinocytotic vesicles then fuse to lysosomes where enzymes are available for degradation, preventing re-entry into the brain. Enzymes for degradation of neuropeptides, including aminopeptidase N, dipeptidyl-peptidase, enkephalinase, endo-oligopeptidase, have been localized to EC (Matsas et al., 1986; Bourne et al., 1989; Birch et al., 1990; Oliveira et al., 1990; Schwartz et al., 1991; Gee et al., 1993). Receptors for some drug classes and enzymes for drug metabolism have also been found in



EC (Benavides et al., 1983; Abramovitz et al., 1988), implicating EC in removal of xenobiotics.

There is some evidence that ependyma can function in immune responses as well. Following injections of interferon- $\gamma$ , ependyma express tissue necrosis factor alpha and major histocompatibility complex II (Tarlow et al., 1993; Lampson et al., 1994). Macrophages are found normally on the apical surface of EC. During infection ependyma have increased expression of vascular adhesion molecule 1 (VCAM-1) and intercellular adhesion molecule 1 (ICAM-1) that may further promote macrophage adhesion (Deckert-Schluter et al., 1994; Strazielle and Gherzi-Egea, 2000).

The ependymal layer is capable of regulating interstitial fluid composition. Microvilli on the apical surface of EC have a glycocalyx coating, and sialic acid, poly-N-lactosamine and D-galactose residues could be involved in the transfer of information about CSF chemical composition (Paulson, 1989; Del Bigio, 1995). Receptors for various growth factors are also present, and EC produce both basic and acidic fibroblast growth factor. Since EC do not proliferate *in situ* in response to growth factors, they may regulate some other function besides growth in EC (Del Bigio, 1995). Peptidases involved in the synthesis or activation of CSF-born neuropeptides, such as peptidyl glycine alpha amidating monooxygenase, gamma glutamyl transferase, tissue kallikrein, furin, and angiotensin converting enzyme (ACE), are found in EC (Del Bigio, 1995). ACE is part of the angiotensinogen (possibly supplied from astrocytes) to angiotensin II conversion pathway, and in rats angiotensin II receptors have been localized to EC (Gehlert et al., 1991). Since angiotensin II is involved in regulating peripheral water balance, it may also play a similar role in the CNS. Vasopressin, found in CSF, is known to increase EC permeability to water

(Rosenberg et al., 1986; Sorensen, 1986; Chen et al., 1993). Furthermore, aquaporin channel 4 has been localized to the ependymal basolateral membrane and suggests that movement of water between interstitial fluid and CSF may be regulated by ependyma (Jung et al., 1994). Thus the BBB and brain-CSF barriers regulate the exchange of substances between the blood and the brain and monitor CSF composition, activities critical for the normal functioning of the central nervous system.

**Functions of cerebrospinal fluid.** CSF also has specific functions critical for maintaining a healthy central nervous system (CNS). First, CSF cushions and supports the brain through a buoyancy effect. CSF is more than 99% water and has a specific gravity nearly equal to that of nervous tissue. As a result, the brain is actually suspended in CSF, reducing the effective weight of the brain by more than 95% and protecting the brain from shearing and tearing forces caused by acceleration and deceleration associated with normal physical activity (Bergsneider, 2001; Johanson, 2003; Brodbelt and Stoodley, 2007). CSF also protects the brain by maintaining a constant intracranial pressure (ICP). As stated by the Monro-Kellie doctrine, blood, brain and CSF are enclosed in a rigid space. Since tissue and fluid cannot compress, a change in the volume of one component must be compensated for by a nearly equal and opposite change in another. As arterial and central venous pressure fluctuates with respiration, exertion and posture changes, CSF volume inversely increases or decreases to maintain ICP within normal ranges. CSF volume is regulated by absorption rates; absorption into venous blood is rapidly enhanced when ICP increases, and is slowed when ICP drops. CSF pressure can change as much as 1-2 mm H<sub>2</sub>O with arterial pulse and 2-10 mm H<sub>2</sub>O with breathing, illustrating the importance of CSF compensation. In the event of a sudden

increase in intracranial blood volume, CSF can also be temporarily stored in the subarachnoid space (Weller et al., 1992; Segal, 1996; Johanson, 2003; Di Terlizzi and Platt, 2006).

In addition to physically protecting the brain, CSF has many roles related to its circulation through the CNS. Micronutrients are transported by the choroid plexus into CSF for distribution to target cells. Vitamin C, folates, deoxyribonucleosides, vitamin B<sub>6</sub>, and trace elements are examples of nutrients that enter brain parenchyma via CSF bulk flow and diffusion across ependymal and pial membranes. CSF also carries hormones and signaling molecules needed for homeostasis, trophic support and neurodevelopment. Growth factors and peptides like transthyretin, insulin-like growth factor, thyroxine, transforming growth factor  $\beta$  (TGF $\beta$ ), growth hormone, bone morphogenic proteins, arginine vasopressin and fibroblast growth hormone are all transported from the blood to the brain in CSF. CSF circulation is an important part of neuroendocrine signaling as centrally released peptides are carried in CSF to their targeted areas. For example, hormone-releasing factors from the hypothalamus (i.e., gonadotropin releasing-hormone, thyrotropin releasing-hormone and growth hormone releasing-hormone) are carried in the CSF to the median eminence (Johanson, 2003; Di Terlizzi and Platt, 2006; Johanson et al., 2008). CSF helps to regulate hypothalamic and respiratory function by carrying growth factors and peptides to the hypothalamus, medulla oblongata and the choroid plexus (Brodbelt and Stoodley, 2007). Tanycytes, specialized ependymal cells located in the third ventricle of adults, have long processes that extend into the neurophil and wrap around blood vessels. They are believed to be a part of the neuroendocrine system since they form a direct link between CSF and the capillary plexus in the hypothalamus (Di Terlizzi and Platt, 2006; Brodbelt and Stoodley,

2007). Additionally, CNS provides a route for parasympaptic neurotransmitter transmission. Neurotransmitters that leak from synapses are carried in the interstitial fluid along perivascular and subependymal pathways (the ependymal basement membrane is continuous with perivascular membranes) to CSF where they may provide a background signaling “tone” (Johanson, 2003).

Finally, CSF transports metabolic wastes, toxins and ions to sites for removal. Proteins from the breakdown of tissues, macromolecules, cellular debris, ions, xenobiotics, and organic anions such as homovanillic acid and 5-OH-indolic acetic acid (metabolites of dopamine and serotonin, respectively) are carried by CSF to the arachnoid villi where they are absorbed into venous blood (Bergsneider, 2001), to the lymphatic system where they are drained into lymph, or to the choroid plexus where they are removed by transporters. The choroid plexus expresses transporters for a variety of substances, including P-glycoprotein and multidrug-resistant protein for taking up organic anions. These transporters also efficiently clear some drug classes and antibiotics. By draining into lymphatics CSF connects the CNS, which lacks its own lymphatic system, to the peripheral lymphatic system and thus provides the potential for foreign substances in the CNS to trigger an immune response. Furthermore, CSF provides a “sink” for diluting excessive ions that flow into CSF from brain interstices. Potassium and  $H^+$  are two important examples; a buildup of  $K^+$  in interstitial fluid causes seizures while too much  $H^+$  leads to ischemia (Johanson, 2003; Di Terlizzi and Platt, 2006; Brodbelt and Stoodley, 2007).

Clearly CSF plays a critical role in normal CNS function beyond physical protection. CSF is ideally situated to regulate the composition of the interstitial fluid that bathes neurons and glia and to facilitate long-range communication between different brain regions. By

supplying needed nutrients and removing excess waste, CSF helps to maintain interstitial fluid concentrations within the narrow ranges required for optimal neuronal function.

**Cerebrospinal fluid production.** Choroid plexus tissue has long been recognized as the site of CSF production. There are four choroid plexi in the brain, one in each of the ventricles. Choroid plexi form frond-like projections from the roof of the third and fourth ventricles and from the walls of each lateral ventricle. The choroid plexi in the lateral ventricles are flat and thin, resembling a leaf, while the choroid plexus of the fourth ventricle is more complex and lobular in appearance. The third ventricle choroid plexus is the smallest, and its appearance is a combination of the other two (Strazielle and Gherzi-Egea, 2000). Studies using isolated ventricles and choroid plexus preparations estimate that 60 to 75 percent of CSF formation is from choroid plexus (Johnston and Teo, 2000; Johanson, 2003; Brodbelt and Stoodley, 2007).

Choroid plexus tissue is composed of a single layer of tightly packed cuboidal epithelial cells surrounding a central core containing a capillary bed and loose connective tissue. The choroid plexus epithelial layer is continuous with the ependymal cell layer that lines the ventricles, but unlike the ependyma, the choroid plexus epithelial layer has tight junctions in between the cells on their apical surface. These tight junctions prevent the majority of hydrophilic ions and molecules from crossing the cell layer into the CSF, although the electrical resistance of choroid plexus epithelium is less than that of the BBB (Johanson, 2003). The choroid plexus epithelium folds into many villi around each capillary, creating processes that project into the ventricles. The basolateral membrane of choroid plexus epithelium also has many interdigitations and infoldings. The villi, along with a brush

border of microvilli on the apical surface, greatly increase the surface area of the choroid plexi. Choroid plexi have a surface area of approximately  $100 \text{ cm}^2$  per gram tissue, highlighting its importance as a secreting and reabsorbing tissue. Not surprisingly, choroid plexi have a high density of mitochondria and an extensive Golgi apparatus to support its secretory function (Strazielle and Gherzi-Egea, 2000; Johanson, 2003). In addition to CSF production, the choroid plexus acts as a filtration system, removing metabolic waste, foreign substances, and excess neurotransmitters from the CSF. Peptidases in the microvilli of the apical surface regulate peptides in CSF (Brodelt and Stoodley, 2007). In this way the choroid plexus has a very important role in helping to maintain the delicate extracellular environment required by the brain to function optimally.

While choroid plexi are believed to produce the majority of CSF, they are not the sole source of CSF formation. Removal of the choroid plexus did not abolish CSF formation in isolated ventricle experiments (Hassin, 1948). In 1969, Milhorat removed the choroid plexus (choroid pexectomy) from rhesus monkeys and compared the development of induced hydrocephalus in the plexectomised monkeys to monkeys with intact choroid plexus, and found only small changes in hydrocephalus severity between the two groups (Milhorat, 1969). Further research showed that bilateral choroid plexectomy reduced CSF formation by only 33-40% (Milhorat et al., 1971; Milhorat, 1974). A human case study found CSF production to be within normal range in a 5 year old child who had undergone choroid plexectomy as an infant (Milhorat et al., 1976).

Sources for extraventricular CSF production have not been definitively established; however proposed sources include capillaries and metabolic processes in the parenchyma, ependymal cells, and spinal subarachnoid space (Johanson et al., 2008). The endothelium

surrounding capillaries in the brain appear to be capable of secreting fluid and osmolites. Like choroid plexus epithelium, capillary endothelium is polarized with different ion transporters and channels expressed on the basal and apical surfaces.  $\text{Na}^+\text{K}^+\text{ATPase}$ , a pump essential in CSF formation in choroid plexus, is expressed on the apical surface and thus could drive the net secretion of sodium and chloride, the main components of CSF (Segal, 1996; Abbott, 2004). Metabolically, water is produced in cells as a byproduct when glucose is oxidized to  $\text{CO}_2$  and could thus contribute to interstitial fluid formation, although not enough water is generated this way to account for the total volume of interstitial fluid (Rapoport, 1976; Abbott, 2004). Ependymal cells have several characteristics suggesting they have secretory capabilities. In rat ependyma, mRNA for the secretory proteins chromogranin B, secretogranin, and sulfated glycoprotein-2 have been found (Senut et al., 1992; Danik et al., 1993; Gee et al., 1993). Granulophysin, a membrane protein associated with exocytosis, is present in human ependymal cells (Hatskelzon et al., 1993) and surface blebs and vesicles have been observed in the ependymal cells lining the cerebral aqueduct in rabbits and pigeons (Meller and Dennis, 1993; Mestres and Rascher, 1994). During development, secretory droplets can be found on the apical surface of third ventricle ependyma (Booz, 1975). Although there are some reports of CSF from spinal sources (Sato et al., 1973), multiple studies have failed to find significant spinal subarachnoid space CSF production in cats, dogs and monkeys (Lux and Fenstermacher, 1975; del Pozo-Reyes et al., 1978; Hamer and Sahar, 1978), making the subarachnoid space an unlikely source for CSF formation.

CSF is an aqueous solution composed mainly of sodium, chloride and bicarbonate, similar to blood plasma from which it is derived. Nascent CSF contains the following ion


concentrations (in mEq/kg H<sub>2</sub>O): 158 Na<sup>+</sup>, 138 Cl<sup>-</sup>, 25 HCO<sub>3</sub><sup>-</sup>, 3.3 K<sup>+</sup>, 1.7 Ca<sup>2+</sup>, and 1.5 Mg<sup>2+</sup>. CSF is not, however, simply a filtrate of plasma. CSF differs from plasma with slightly higher levels of Cl<sup>-</sup> and Mg<sup>2+</sup> and lower levels of Ca<sup>2+</sup> and K<sup>+</sup> (Johanson, 2003; Di Terlizzi and Platt, 2006). Ca<sup>2+</sup> and K<sup>+</sup> levels in particular must be kept stable in CSF as small changes in their concentrations impact CNS excitability. Accordingly, changes in plasma concentrations of these ions have little or no effect on CSF concentrations (Crone and Christensen, 1981). Protein, urea, and glucose are present in CSF but at levels 30-40% less than plasma, and amino acids are present in only trace amounts (Johanson, 2003). Normally CSF is clear and colorless and has a total nuclear cell count of only 0-2 cells/ $\mu$ L (Meinkoth et al., 1998); any cloudiness or color is a sign of pathology.

CSF is formed in a two step process, ultrafiltration of plasma from blood and active net secretion of Na<sup>+</sup>, Cl<sup>-</sup> and bicarbonate ions (HCO<sub>3</sub><sup>-</sup>) into the ventricles. Capillaries in choroid plexus are large and have thin, fenestrated endothelial walls, allowing plasma, macromolecules and ions to filter from the blood into choroid plexus interstitial fluid. Under normal conditions, blood flow to the choroid plexus capillaries is quite high, flowing ten times faster than cerebral blood flow (Johanson, 2003), and filtration of plasma is proportional to the hydrostatic pressure gradient between blood and choroid plexus interstitial fluid (Strazielle and Ghersi-Egea, 2000; Johanson et al., 2008).

The net flow of ions from plasma across choroid plexus basolateral and apical membranes and into the ventricular space is driven by the coordinated activity of ion transporters and channels (Figure 3). On the apical membrane, sodium is transported into CSF by ATP-dependent Na<sup>+</sup>K<sup>-</sup>ATPase and Na<sup>+</sup>K<sup>+</sup>2Cl<sup>-</sup> cotransporters. The activity of these pumps and cotransporters keeps the internal concentration of sodium around 20-30 mM,



establishing a favorable sodium gradient for secondary active transport of sodium across the basolateral membrane by  $\text{Na}^+\text{-H}^+$  exchange and  $\text{Na}^+\text{-HCO}_3^-$  cotransporters (Bergsneider, 2001; Praetorius et al., 2004; Bouzinova et al., 2005). Along with the bicarbonate ions brought into the choroid plexus (against its concentration gradient) by  $\text{Na}^+\text{-HCO}_3^-$  cotransporters, bicarbonate is accumulated in the cell as carbonic anhydrase hydrates  $\text{CO}_2$  into  $\text{HCO}_3^-$  and  $\text{H}^+$ . The buildup of  $\text{HCO}_3^-$  allows chloride ions to be brought into the choroid plexus epithelium by  $\text{Cl}^-\text{-HCO}_3^-$  exchange in the basolateral membrane. Chloride may also enter by a  $\text{K}^+ \text{- Cl}^-$  cotransporter as the KCC3 isoform has been localized to the



**Figure 3. Choroid plexus epithelium and the production of cerebrospinal fluid.** CSF production, the net flow of bicarbonate ( $\text{HCO}_3^-$ ), chloride ( $\text{Cl}^-$ ) and sodium ions ( $\text{Na}^+$ ), occurs through the coordination of channels, transporters and cotransporters in the choroid plexus epithelium.  $\text{HCO}_3^-$  accumulates in the cytoplasm via  $\text{Na}^+\text{-HCO}_3^-$  cotransporters and  $\text{Cl}^-\text{-HCO}_3^-$  exchange pumps in the basolateral membrane and by carbonic anhydrase (c.a.) activity.  $\text{HCO}_3^-$  and  $\text{Cl}^-$  enter the CSF through inward rectifying anion channels in the apical membrane. Sodium is actively pumped into CSF by apical membrane  $\text{Na}^+\text{K}^+\text{ATPase}$  and  $\text{Na}^+\text{K}^+2\text{Cl}^-$  cotransporters, creating a favorable concentration gradient that allows sodium to enter the cell via  $\text{Na}^+\text{-H}^+$  exchange and  $\text{Na}^+\text{-HCO}_3^-$  cotransporters in the basolateral membrane. Abbreviations:  $\text{Na}$ =sodium ion,  $\text{K}$ =potassium ion,  $\text{HCO}_3^-$ =bicarbonate ion, c.a.=carbonic anhydrase,  $\text{Cl}$ =chloride ion. Reproduced in agreement with the BioMed Central open access policy. Johanson *et al. Cerebrospinal Fluid Research* 2008 5:10 doi:10.1186/1743-8454-5-10

basolateral membrane (Pearson et al., 2001). Inward rectifying anion channels in the apical membrane allow chloride and  $\text{HCO}_3^-$  to enter CSF by moving down their respective electrochemical gradients. The resulting hyperosmolar solution causes water to move into CSF via aquaporin 1 channels on the choroid plexus apical membrane (Bergsneider, 2001; Brown et al., 2004; Johanson et al., 2008). There is, of course, reverse movement of ions due to reabsorptive fluxes to the blood, and other ions such as calcium and magnesium are moved between blood and CSF; however these fluxes, although important for CNS function, are not instrumental in CSF production (Johanson, 2003; Brown et al., 2004).

CSF is made continuously and is estimated to turn over three times in twenty-four hours in humans, which corresponds roughly to 600 mL of new CSF/day in healthy adults (Johanson, 2003). Choroid plexus (human) produces CSF at 0.3-0.6 mL/min/g, although there is some question as to whether CSF is produced constantly at this rate. In a quantitative MR study monitoring CSF flow, CSF formation followed circadian rhythms, with a maximum rate of production at 2:00 AM and a minimum rate of production at 6:00 PM (Nilsson et al., 1992). There are other factors that influence CSF production as well. The choroid plexus receives neural input via sympathetic input; sympathetic stimulation decreases CSF production while removal of sympathetic input greatly increases production. Although the choroid plexus receives cholinergic and peptidergic innervations, their influences on choroid plexus function are not well understood (Johnston and Teo, 2000).

The role of ICP and blood flow on CSF production also remains unclear. As mentioned previously, the rate of CSF production is dependent on the hydrostatic pressure gradient between choroid plexus interstitial fluid and blood plasma. Thus it seems likely that changes in blood flow and ICP would affect the rate of CSF production. Blood flow to the


choroid plexus is regulated by endocrine and neurohumoral inputs (Shapira et al., 1992; Strazielle and Gherzi-Egea, 2000; Johanson et al., 2008) that seem ideally suited for controlling CSF formation. Yet experiments manipulating interstitial fluid and blood flow have had conflicting results on CSF formation. For example, administration of acetazolamide, an inhibitor of carbonic anhydrase, caused a 50% decrease in CSF formation while simultaneously increasing choroidal blood flow (Faraci et al., 1990). A similar result is seen with vasoactive intestinal polypeptide (VIP). When given intraventricularly, blood flow increased 20% yet CSF formation dropped by 30%. Choroidal blood flow also increased when VIP was administered i.v., but CSF formation remained unchanged (Nilsson et al., 1991). Conversely in acute hydrocephalus, increased ventricular CSF pressure causes an increase in choroid plexus interstitial fluid pressure, reducing plasma filtration and subsequent CSF formation. Blood flow to the choroid plexus is also reduced in HC (Johanson et al., 2008). Overall, most evidence suggests that within the rather narrow limits of normal physiology, CSF production is relatively independent of both ICP and blood flow (Bergsneider, 2001). It is conceivable that the observed decreases in CSF formation are compensating for increased ICP caused by increased blood flow in CNS, however this area remains controversial.

**Cerebrospinal fluid absorption.** The classic pathway of CSF movement in the CNS (Figure 4) is for CSF to circulate by bulk flow from its site of origin at the choroid plexus, through the ventricles and basal cisterns, down the posterior side of the spinal cord, up the anterior side, and through the cranial subarachnoid space to the midline convexity where it is absorbed into the dural sinuses by arachnoid villi (Welch and Friedman, 1960; Arkhinova,

1966; Di Chiro, 1966; Du Boulay, 1966). CSF movement is believed to be driven by new CSF production, postural effects, pulse waves generated by the choroid plexus, arterial or respiratory pulsations, and cilia movement (Gonzalez Santander et al., 1984; Roth et al., 1985; Johnston and Teo, 2000; Bergsneider, 2001). The arachnoid villi (also known as pacchionian granulations or arachnoid granulations) are invaginations of the arachnoid membrane into the dural sinuses. Their role in CSF absorption was first described by Welch and Friedman (1960), who believed they contained one-way valves that open when CSF pressure exceeds venous pressure. Radiocisternograms, where large molecule radioisotope-labeled tracers are injected into CSF, showed an accumulation of tracers over the cerebral convexity where the arachnoid villi are located (Di Chiro, 1966; Bergsneider, 2001). Together these studies were interpreted as evidence for the arachnoid villi being the main site of CSF absorption (Milhorat, 1987; Bergsneider, 2001; Di Terlizzi and Platt, 2006).

It is now evident that CSF movement through the CNS is more complex than originally described. Flow-sensitive MR imaging shows bulk flow of CSF through the ventricular system but no net flow around the spinal cord or towards the cerebral convexity (Schossberger and Touya, 1976; Greitz, 1993; Bergsneider, 2001; Abbott, 2004). Instead, CSF movement in the subarachnoid space and around the spinal cord is pulsatile in nature, which is caused by changes in brain blood volume during the cardiac cycle. Blood flows into the brain during systole and out during diastole, causing the brain to move up and down like a piston as subarachnoid CSF is dispelled to accommodate increased blood volume. During systole, CSF in the subarachnoid space and cisterns moves into the compliant spinal canal, while CSF in the brain moves first retrograde through the aqueduct and foramen of Magendie and then moves anteretrograde at mid-systole. During diastole CSF again moves retrograde

through the aqueduct and foramen (Greitz et al., 1997a; Bergsneider, 2001). This pulsatile CSF flow could account for the concentration of tracers around at the arachnoid villi. Furthermore, in radionuclide cisternography studies there is a second site of tracer accumulation located at the lumbar-sacral sac that is unaccounted for by the bulk flow model (Greitz and Hannerz, 1996). As new CSF flows from the foramina of Luschka and Magendie it mixes with and dilutes the tracers. The accumulations of tracer in radiocisternograms are thus due to a lack of dilution since these sites are the furthest away from the flow of new



**Figure 4. Classic view of cerebrospinal fluid circulation.** Cerebrospinal fluid is produced by the choroid plexus and moves by bulk flow through the lateral, third, and fourth ventricles into the subarachnoid space. After flowing through the spinal canal and around the brain, CSF is finally absorbed into the dural sinus via the arachnoid granulations (or arachnoid villi).

Image obtained from [www.corkrey.com](http://www.corkrey.com)

CSF.

The evidence against CSF circulating via bulk flow raises the question of CSF absorption. If most CSF does not move to the arachnoid villi, then where is it absorbed?

Even before the pulsatile movement of subarachnoid CSF was demonstrated, the arachnoid villi as the primary site for CSF drainage were a topic of debate. Human infants do not develop arachnoid villi until the fontanelles fuse, meaning there must be other routes for CSF drainage; and in spite of numerous references to valve-like channels in arachnoid villi, no such structures have ever been found (Greitz et al., 1997a, b). Furthermore, other studies found various tracers appeared in blood or nervous tissue faster than could be accounted for by bulk flow. Albumin, inulin and sucrose injected intrathecally into rats reached half-maximal concentrations in the blood and CSF in 100, 80 and 30 minutes, respectively. By bulk flow it should take 6-8 hours to reach the arachnoid villi (Reed and Woodbury, 1963; Greitz et al., 1997a, b). In 1972, iodinated albumin injected into lumbar subarachnoid space was absorbed directly into the spinal canal (Soderborg and Lying-Tunell, 1972). CT cisternography showed that contrast media penetrates into brain parenchyma (Drayer and Rosenbaum, 1977), and albumin and x-ray contrast media were found in blood only minutes after injection into the subarachnoid space (Greitz et al., 1997a). When radioactive water was injected into the carotid artery, nearly all was found to penetrate the brain and almost 50% returned to blood within two minutes, indicating that the movement of water and the movement of solutes are not necessarily the same (Oldendorf, 1970; Greitz et al., 1997a; Bergsneider, 2001). This was confirmed by comparing 3H-water and 3H-inulin injected into the lateral ventricles of cats. The 3H-water was rapidly absorbed into adjacent cerebral capillaries while 3H-inulin moved bidirectionally along the ventricular system and was slowly absorbed into blood (Bulat et al., 2008; Vladic et al., 2009).

The current concept of CSF flow and absorption focuses on the nasal lymphatic system as a primary site for CSF drainage. Lymph and CSF have been thought to interact for

over a century, but work within the last decade has shown that the majority of CSF in multiple mammalian species (including human and non-human primates) flows out of the subarachnoid space along the perineural spaces of the olfactory nerves, through the cribriform plate, and into the submucosa, where lymphatic ducts form a collar around the olfactory nerves. CSF is ultimately returned to venous blood through the cervical lymphatics (Abbott et al., 1985; Bradbury, 1985; Johnston et al., 2004; Zakharov et al., 2004; Johnston et al., 2005; Koh et al., 2005; Johanson et al., 2008). Spinal nerve rootlets also connect with lymphatic vessels, and CSF is absorbed similarly, although on a smaller scale, in the spinal cord (Johanson et al., 2008).

A secondary site of CSF absorption is believed to be through blood vessels in the parenchyma. Water moves freely between blood and interstitial fluid, and larger molecules (albumin, iodide, horseradish peroxidase) are evidently able to enter the blood, probably through carriers or by pinocytosis (Greitz et al., 1997a). CSF in the subarachnoid space enters the parenchyma through Virchow-Robin spaces (Figure 2) and across the ependymal layer, preferentially flowing in the subependymal space, white matter tracts, and around blood vessels. Once in the parenchyma, CSF mixes with interstitial fluid and can be absorbed through capillary endothelium (Brightman, 1965; Abbott, 2004).

Although arachnoid villi are no longer believed to be the primary site of CSF absorption, further investigation into their function has suggested they do have a role as an auxiliary drainage route when CSF pressures are elevated (Johnston et al., 2004; Johanson et al., 2008). Significant concentrations of soluble and particulate tracers are found in the dural sinuses when CSF-sinus blood pressure gradients exceed 20 cm H<sub>2</sub>O, but not at physiological pressure gradients of 5 cm (Mann et al., 1979; McComb, 1983, 1992; Johnston et al., 2004).

The original thought that CSF moves through the arachnoid villi via valve-like structures is questionable, however. Large vacuoles seen in the basal surface of arachnoid villi epithelium and the presence of pinocytotic vesicles on the endothelium of sinuses indicate that fluid is transported into blood instead of flowing through an open channel (Johnston and Teo, 2000).

The ventricular system and CSF carry out multiple functions within the CNS, providing physical protection, communication routes, and a means for removing harmful substances from the brain. The physiology of CSF production, means of movement through the CNS, and routes of absorption are clearly not fully understood but are obviously critical for normal functioning of the tightly regulated environment of the CNS. A disturbance in any of the components (i.e. CSF production, physical pathways, or absorption routes) has serious consequences, including hydrocephalus. Hydrocephalus can result from perturbations of any and all parts of CSF physiology, and is described further in the next chapter.



## **CHAPTER 2**

### **HYDROCEPHALUS: CLASSIFICATION, CAUSES, PATHOLOGY, AND TREATMENT**

Hydrocephalus (HC) is a multifactoral, progressive brain disorder characterized by accumulation of cerebrospinal fluid (CSF) in the brain and subsequent expansion of the ventricles. HC is the most common neurosurgical problem in children with an incidence of 1-2 cases per 1000 live births (Peacock and Currer, 1984; Leech and Payne, 1991; Casey et al., 1997; Johanson et al., 2001; Fletcher et al., 2002). The advent of better diagnostic tools has allowed adult onset of HC to be recognized as an increasing health issue. Approximately 69,000 to 80,000 new cases of HC (both adult and childhood) are diagnosed each year in the United States (Bondurant and Jimenez, 1995), although the actual number of cases may be underreported since HC in adults can mimic other neurological disorders (Casmiro et al., 1989a).

#### **Classification**

Communicating vs. non-communicating. HC cases can be divided into two broad categories, communicating vs. non-communicating. Either type may be congenital, where the patient is born with HC, or acquired, where HC develops after birth. Both congenital and acquired HC can be acute, where the ventricles are enlarging and intracranial pressure (ICP) is high and/or unstable, or chronic, where ventricles are enlarged but ICP is normal or slightly elevated but

stable. Frequently the same initial insult can cause multiple types of HC, depending on when the insult occurs in conjunction with undetermined genetic and environmental factors. For example, intracranial hemorrhage leading to HC can occur during gestation, thus causing congenital HC, or at any age after birth, causing acquired HC. The cause of hemorrhage can vary from genetic disorders to traumatic injury, and the resulting HC can be communicating or non-communicating, acute or chronic. Although various systems for classifying HC can be found, the communicating vs. non-communicating system seems to be the most common.

HC in which the ventricular system is free from physical obstruction is called communicating HC. Communicating HC thus must result from either increased production or decreased absorption of CSF, although cases where CSF flow is blocked outside the ventricular system (i.e., in the basal cisterns or parasagittal arachnoid villi) are frequently included in this category. Despite the extremely diverse etiology of HC, abnormalities of CSF circulation leading to HC are usually due to compromised absorption into venous blood; HC due to increased production of CSF is extremely rare (Mori et al., 1995; McAllister and Chovan, 1998; Johanson et al., 2001). Bleeding into the ventricles and infection of the meninges result in compromised CSF absorption and are the leading causes of communicating HC (Beni-Adani et al., 2006). Non-communicating HC occurs when physical obstruction of the ventricular system prevents normal CSF flow. Congenital abnormalities, tumors, genetic defects, stenosis of the aqueduct, meningitis and hemorrhage are common sources of obstruction (Vertinsky and Barnes, 2007).

Chronic hydrocephalus. In 1965 Hakim and Adams described cases of HC where intracranial pressure (ICP) was normal and coined the term “normal pressure

hydrocephalus”, or NPH to describe this condition. Today it is recognized that there are several different types of HC without elevated ICP and the term chronic hydrocephalus is used to describe these cases. Chronic HC is defined as all cases with both ventriculomegaly (Evans index greater than 0.3) and normal or chronic low-grade increase of ICP (Edwards et al., 2004). This definition includes compensated (arrested) HC, *ex vacuo* HC, adult onset stenosis of the aqueduct (Harrison et al., 1974; Oi et al., 2000) and normal pressure hydrocephalus (NPH), both secondary NPH where HC occurs after a clear insult (i.e., meningitis, SAH, trauma) and idiopathic NPH where no clear precipitating factor is present (Hebb and Cusimano, 2001; Zemack and Romner, 2002). Chronic HC is seen most frequently in older children or adults. Like acute HC, chronic HC has multiple precipitating causes and has a similar pathology, although chronic HC progresses more slowly as the adult brain appears better able to compensate than the immature or developing brain (Edwards et al., 2004).

The actual number of chronic HC cases and its rate of development is unknown. In the US, adult cases of chronic HC are estimated at approximately 40,000/year, comprising more than half of new hydrocephalus cases (Bondurant and Jimenez, 1995). Other studies estimate 2.6 new cases per 100,000 people each year world-wide (Hoglund et al., 2001). Although not all cases of chronic HC have cognitive impairments, chronic HC can present symptoms that are diagnosed as dementia, and an estimated 5-10% of patients with dementia are thought to actually have chronic HC (Fisher, 1982; Hakim et al., 2001; Kuba et al., 2002; Vale and Miranda, 2002).

Compensated congenital hydrocephalus, also known as arrested HC, occurs when the ventricles spontaneously cease enlarging and stabilize (Whittle et al., 1985). Increased CSF

absorption by circumventricular and intraaxial blood vessels is thought to account for the halt of ventricular expansion, although cases of spontaneous ventriculostomy are not uncommon. Compensation occurs in 10-15% of congenital HC cases, and usually occurs when the child is between 1 and 2 years of age (Schick and Matson, 1961; Hochwald et al., 1969; Nonaka et al., 2008). Compensation does not mean patients are healthy, however. An estimated 2/3 of untreated children with compensated HC have below normal intelligence, and progressive loss of cognitive function may occur throughout adulthood (Laurence and Coates, 1962; Yashon, 1963; Oi et al., 2000; Edwards et al., 2004; Nonaka et al., 2008). They also have an increased risk of epileptic seizures and sudden death (Rickert et al., 2001; Black and Graham, 2002). Although the brain has compensated for the initial cause of HC, there is little tolerance for further insult to CSF pathways, so even modest straining, exertion or injury can cause instantaneous death from ischemia in cardio-respiratory centers in the brainstem (Botez et al., 1977; Dobkin, 1978; Dickerman et al., 2003; Edwards et al., 2004). The deterioration of cognition in adulthood further suggests that HC is not fully compensated and is actually progressing at a very slow rate. Children with compensated HC are generally not shunted, but newer research suggests shunting would be beneficial as neurological damage does occur once a threshold of ventricular enlargement is reached. The slower the expansion, the larger the ventricles can be before significant axonal damage and cognitive and motor impairments occur, which would explain the delay in onset of symptoms (Del Bigio et al., 2003).

Ex vacuo hydrocephalus. *Ex vacuo* hydrocephalus occurs when brain parenchyma atrophies, resulting in enlarged ventricles (Bradley, 2001b). *Ex vacuo* HC historically was not considered to be related to CSF hydrodynamics, but this conclusion is controversial

(Silverberg et al., 2003; Silverberg, 2004). Proper CSF circulation is being recognized as important for progenitor cell proliferation and migration, particularly in the germinal matrix of the subventricular zone, since CSF carries growth factors and other signaling molecules necessary for normal development. Neuronal death, loss of white matter, and defects in neuronal precursor proliferation and migration can all result in loss of parenchyma (Johanson et al., 2001). While disrupted CSF hydrodynamics is not the only cause of improper development, it is conceivable that *ex vacuo* HC could be related to loss of proper CSF flow.

### **Causes of hydrocephalus**

Physical malformations. Developmental malformations that physically change the ventricular system frequently have HC as part of their pathology. The most common developmental cause of HC is Chiari II (or Arnold-Chiari) malformation, where the cerebellum and brainstem extend into the foramen magnum and block normal CSF flow. Approximately one third of infantile HC cases are caused by Chiari II malformations (Vertinsky and Barnes, 2007). A second physical cause of HC is Dandy Walker malformation, a developmental abnormality in which the cerebellum is underdeveloped, resulting in a misshapen fourth ventricle that drains CSF into a cyst. Approximately 1 in 5000 infants is born with Dandy-Walker malformation (Parisi and Dobyns, 2003), and approximately 20-80% of cases develop HC after the neonatal period due to obstruction of the foramina of Magendie and Luschka (Vertinsky and Barnes, 2007). Other less common developmental abnormalities also cause HC, but will not be covered here.

Stenosis of the aqueduct. The aqueduct of Sylvius is the narrowest part of the ventricular system and closure, or stenosis, of the aqueduct is a frequent finding in both acquired and congenital HC. Primary stenosis is caused by developmental abnormalities such as forking, septation and gliosis. Onset of HC from primary stenosis generally occurs during the neonatal period or in infancy. Secondary stenosis can occur after various types of insult, including hemorrhage, infection and tumors (Vertinsky and Barnes, 2007). Stenosis of the aqueduct also occurs in HC cases without a clear precipitating factor. Approximately 45% of congenital HC cases have no known cause, and fetal HC with aqueduct stenosis constitutes a major portion of these cases (Castaneyra-Perdomo et al., 2006). Whether stenosis is the cause of HC or a consequence of HC in these cases is unknown, and contradictory evidence from experimental models has made this a topic of debate.

Infection. Infection blocks drainage of CSF from the ventricular system by causing inflammation of the meninges, limiting the flow of CSF through the subarachnoid space to absorption sites. Meningitis is, along with hemorrhage, one of the most common causes of congenital HC, but also causes HC in infants, children and adults. Congenital HC from infection occurs when sub-clinical viral infections in pregnant women infect the fetus; post-natal meningitis can be acquired via sinus or ear infections. Sources of infection include gram-negative bacteria, group B streptococcus, pneumococcus, *Listeria*, neisseria, tuberculosis, cytomegalovirus, toxoplasmosis, mumps, rubella and varicella (Margolis and Kilham, 1969; Herndon et al., 1974; McComb, 1997; Vertinsky and Barnes, 2007).

Hemorrhage. Intraventricular hemorrhage (IVH) is a complication of premature birth (Cherian et al., 2004a; Crews et al., 2004) and occurs in 12-20% of preterm births where the infant weighs less than 1500g. Intraventricular hemorrhage in preterm births (under 32 weeks gestation) results from the combination of a very rich blood supply to the subventricular zone (required to support the high metabolic requirements of this region during development), fragile blood vessels due to lack of glial support, occasional hypoxia (due to respiratory complications), and unregulated swings in arterial and venous pressure. Together these events cause bleeding into the ventricles. Red blood cells and their breakdown products can block the ventricular system or arachnoid villi, leading to HC (Cherian et al., 2004a). Post-hemorrhagic ventricular dilation (PHVD) affects 15-35% of infants with IVH. Motor dysfunction is seen in approximately two thirds of PHVD infants while one third have cognitive dysfunction (Davis et al., 1987; Sheth, 1998; Kazan et al., 2005).

Severe traumatic brain injury can also result in hemorrhage and subsequent HC. HC may occur from IVH, subarachnoid hemorrhage (SAH), blood clots that block CSF drainage, or from venous compression caused by swelling. Glial scars that form after injury can also hinder proper CSF flow. While onset of HC can occur within weeks of the insult, in some cases HC develops very slowly and takes months, even years, to become evident.

Genetic disorders. Known genetic abnormalities leading to HC account for a very small percentage of HC cases. X-linked recessive mutations in the neural cell adhesion molecule (CAM) gene for L1 neural cell adhesion molecule are known to cause HC (Van Camp et al., 1993; Weller and Gartner, 2001). There are case reports of inherited HC that presents as an

autosomal recessive disorder in some families, but no genes responsible have been identified (Johanson et al., 2001). However, the large number of experimental HC models resulting from genetic alterations suggests that other cases of human HC with unidentifiable causes may also have a genetic component.

### **Molecular mechanisms of hydrocephalus**

HC has an extremely diverse etiology and is found as part of a variety of neurological disorders. Yet the progression and outcomes of HC are similar, implying that some common underlying molecular mechanisms are present. These underlying molecular mechanisms have yet to be fully elucidated and HC remains poorly understood, due at least in part to the difficulty in differentiating between changes that are a consequence of HC from those that are causing HC. Research from experimental HC models has provided some valuable insights into the pathology of HC which are summarized below.

Disregulation of cytokines. In addition to secreting CSF, the choroid plexus secretes a variety of growth factors necessary for proper CNS function, including FGF-1, FGF-2, TGF- $\beta$ 1, NGF, HGF-SF and IGF-II. Signaling molecules present in CSF are important for regulating cell growth and development. The subventricular zone lies very close to the lateral ventricles and contains progenitor cells that continue to proliferate into adulthood. The proximity of the subventricular zone to CSF suggests that the growth factors present in CSF would impact proliferation in this region. Areas of the brain more distant from the ventricles are also affected by CSF contents. For example, Cajal-Retzius cells, which regulate the laminar organization of the cortex, send fibers from their location in the marginal



zone to the ventricles during development, implying that during development the CSF contains factors important for normal cortical development (Johanson et al., 2001). Accordingly, changes in cortical layering and progenitor proliferation have been observed in HC.

Restriction of CSF flow in HC likely causes signaling molecules to become concentrated in the ventricles and alterations in cytokine levels could account for observed changes in progenitor cells. In support of this theory, altered cytokine levels are found in CSF from various experimental models of HC, and CSF obtained from affected H-Tx rats inhibits the proliferation of cultured neuronal precursor cells from normal controls (Vetsika et al., 1999; Owen-Lynch et al., 2003). If the concentration of growth factors in the CSF is changed in HC, one would expect a compensatory feedback on their production. Research investigating the choroid plexus *in vitro* showed a decrease of EGF, FGF-2, and TGF- $\beta$ 1 mRNA and a 10-fold increase in IGF-II mRNA in the choroid epithelium from HC animals (Johanson et al., 2001).

While the observed changes in CSF may very well be a consequence of HC, altered levels of TGF- $\beta$ 1 and FGF-2 levels have been shown to cause HC (Galbreath et al., 1995; Wyss-Coray et al., 1995; Pearce et al., 1996). Ventricular infusions of recombinant basic fibroblast growth factor (FGF-2) in rats and transforming growth factor  $\beta$ 1 (TGF- $\beta$ 1) in mice both result in chronic HC characterized by ventriculomegaly, reduced CSF formation and increased fibrosis and collagen deposits in arachnoid villi with a patent aqueduct and normal ICP. In rats given FGF-2, enlargement of the lateral ventricles begins after 2-5 days of infusions and ependymal cells over proliferate, forming rosettes that protrude into the third ventricle (Tada et al., 1994; Johanson et al., 1999b). Two separate transgenic models of HC

were developed by over-expressing TGF- $\beta$ 1 (Kitazawa and Tada, 1994; Tada et al., 1994; Galbreath et al., 1995; Wyss-Coray et al., 1995). In both models TGF-  $\beta$ 1 is expressed under control of the GFAP promoter and results in communicating HC. The time of onset and severity of HC in both models correspond to the level of TGF-  $\beta$ 1 expression (Crews et al., 2004); in one model, low expressing mice that are clinically normal develop hydrocephalus following injury (Wyss-Coray et al., 1995). TGF-  $\beta$ 1 is also elevated in a rat model of post-hemorrhage ventricular dilation (Cherian et al., 2004b). In human HC cases, increased risk of developing chronic communicating HC is associated with elevated levels of CSF TGF-  $\beta$  1 in patients with subarachnoid hemorrhage (SAH) (Flood et al., 2001) and increased TGF-  $\beta$  1 expression is found in preterm births with post-hemorrhagic hydrocephalus (Kitazawa and Tada, 1994; Whitelaw et al., 1999).

The TGF-  $\beta$  family of growth factors regulates development, cell growth, differentiation, homeostasis, tissue repair, and deposition of extracellular matrix proteins (ECM) (Roberts and Sporn, 1996; Benveniste, 1998). All three isoforms of TGF- $\beta$  and their receptors are expressed in neurons, microglia, and astrocytes (Flanders et al., 1991; Wang et al., 1995; Bottner et al., 1996; Ata et al., 1997; Slotkin et al., 1997; Lippa et al., 1998). Overexpression of TGF- $\beta$ 1 in perivascular astrocytes results in increased deposition of ECM proteins like collagen, laminin and fibronectin. This may interfere with the ability of CSF to be absorbed into the vasculature (Cohen et al., 1999). TGF- $\beta$ 1 tg mice also have decreased expression and reduced activity of matrix metalloproteinase 9 (MMP9) along with an altered expression pattern of its inhibitor, tissue inhibitor of MMPs 1 (TIMP1). MMP9 breaks down ECM proteins, and decreased MMP9 activity could contribute to the accumulation of fibronectin and laminin found in the ECM surrounding the microvasculature, the choroid

plexus and the arachnoid granulations (Del Bigio, 1993; Wyss-Coray et al., 1995; Zechel et al., 2002). Together these findings suggest that CSF absorption and flow of interstitial fluid could be impacted by the thickened ECM and thus lead to the development of spontaneous hydrocephalus. Observed abnormal migration of cortical neurons during development may also contribute to the pathology (Crews et al., 2004).

Ependymal cell denudation. Loss of ependymal cells lining the lateral ventricles and the aqueduct is commonly seen in human cases of HC as well as in animal models (Page and Leure-duPree, 1983; Jimenez et al., 2001; Dominguez-Pinos et al., 2005). Although ependymal denudation is considered by some to result from stretching of the ventricles as CSF accumulates (Sarnat, 1995; Kiefer et al., 1998), there is a significant amount of research indicating that the loss or damage of ependymal cells can in and of itself cause hydrocephalus. Loss or damage of ependyma may lead to HC because regulation of bulk flow of fluids between the brain parenchyma and CSF is lost (Rodriguez, 1976; Sarnat, 1995; Jimenez et al., 2001; Batiz et al., 2006). A study of eight human fetuses with communicating HC showed progressive ependymal cell loss as early as 16 weeks gestation when HC was only moderate, making loss of ependyma due to stretching unlikely (Johnson and Johnson, 1968; Margolis and Kilham, 1969; Nielsen and Baringer, 1972; Nielsen and Gauger, 1974; Bruni et al., 1988b; Jones and Bucknall, 1988; Yamada et al., 1991; Dominguez-Pinos et al., 2005). In one large scale study of a spontaneous HC model, Rodriguez et al (2006) found that ependymal denudation and abnormal neurogenesis (stemming from molecular and cellular level abnormalities) preceded the onset of HC in a set temporal and spatial pattern,

and strongly states that the changes seen in the brain (ie, loss of ependyma, closure of the aqueduct) are not the result of mechanical processes but are causal in the pathology of HC.

Although multiple experimental HC models have ependymal denudation, denudation in the *hyh* mutant mouse model of HC has been the most extensively studied. *Hyh* mice are a spontaneous HC model recently discovered to have a point mutation in the gene encoding the alpha-SNAP protein. Although how a decrease in alpha-SNAP causes HC is not understood, a programmed denudation of ependymal cells begins on E12 before the mice develop enlarged ventricles and continues until P14. The pattern of denudation follows the pattern of ependymal maturation, thus in HC mice the areas where ependyma mature sooner (ie lateral ventricles) also have denudation sooner (Jimenez et al., 2001; Wagner et al., 2003; Batiz et al., 2006).

There are two distinct phases in the development of HC in the *hyh* mouse. During embryonic development the lateral ventricles, the fourth ventricle, and the floor of the aqueduct undergo denudation and the rostral end of the aqueduct becomes stenosed. This is followed by the onset of moderate HC. After birth, the caudal and dorsal walls of the aqueduct undergo denudation, followed by the obliteration of the caudal region of the aqueduct. This obliteration occurs within the first postnatal week, preferentially during the first 3 days (56%). This is also the time when in normal mice the ependymal cells of the aqueduct lose vimentin expression and undergo ciliogenesis (Wagner et al., 2003). Following obliteration of the aqueduct, the mice develop severe HC.

Denudation follows the developmental maturation of ependyma (rostral to caudal, ventral to dorsal), and developmental defects in brain structures correlates with areas of denudation. Not all ependymal cells detach, however. Patches of specialized ependyma that

resist denudation are found on the circumventricular organs, the dorsal and ventral walls of the rostral end of the aqueduct, in a well defined region of the floor of the aqueduct, and on the wall of the collicular recess. This suggests that a defect in the neuroepithelium/ependymal cell lineage may be responsible for both abnormal development of brain structures and hydrocephalus (Paez et al., 2007).

Scanning electron microscopy of the ventricle walls shortly after birth revealed numerous macrophages in areas that had recently undergone denudation. Detached ependyma are frequently associated with clusters of macrophages. Detaching ependymal cells appear normal, ie, not dying, and have a smooth, spherical appearance (Jimenez et al., 2001; Wagner et al., 2003). Macrophages also are found on areas that will soon undergo denudation and contact exclusively multi-ciliated ependyma. Macrophages are notably lacking on areas that had undergone denudation early in development and on the specialized ependyma that never detach.

Wagner et al (2003) postulated that ependyma in *hyh* mice may have a defect in glycosylation of *N*-linked sialoglycoproteins, resulting in galactose as the terminal sugar residue. This defect would only occur at the stage of differentiation when that enzyme would be active, which would explain the link between differentiation and detachment. Macrophages have receptors for galactose, which could explain the interaction of macrophages with ependymal cells prior to denudation. Ependymal denudation and ependymal rosettes in the neuropil are seen in chick embryos after N-cadherin is blocked by a single injection of antibody into the CSF (Ganzler-Odenthal and Redies, 1998), demonstrating that interfering with surface proteins can lead to ependymal denudation and abnormal migration. Ependymal denudation and sub-ependymal rosette formation are also

seen in infants with post hemorrhage ventricular dilation and in a rat model of post hemorrhage ventricular dilation, although changes in N-cadherin or other surface proteins has yet to be demonstrated (Fukumizu et al., 1995, 1996; Takano et al., 1996; Cherian et al., 2004a).

Subcommissural organ abnormalities. Improper functioning of the SCO is another abnormality associated with HC. Although the exact link is not established, an abnormal SCO has been described in multiple congenital and experimental HC models and in humans (Overholser et al., 1954; Takeuchi et al., 1987; Bruni et al., 1988b; Irigoien et al., 1990; Yamada et al., 1992; Castaneyra-Perdomo et al., 1994; Danielian and McMahon, 1996; Louvi and Wassef, 2000; Estivill-Torrus et al., 2001; Jimenez et al., 2001; Perez-Figares et al., 2001; Blackshear et al., 2003; Baas et al., 2006; Sweger et al., 2007). The SCO is postulated to regulate electrolytes and monitor pressure (Dominguez et al., 1993) and is known to secrete the negatively charged glycoproteins that form the Reissner fiber. The Reissner fiber (RF) extends the length of the aqueduct and may function to prevent the stenosis or obliteration of the aqueduct (Overholser et al., 1954) by repelling the negatively charged glycocalyx of the ependyma. In support of this theory, the sialic acid residues of the RF face the aqueductal lumen (Fernandez-Llebrez et al., 1987; Wagner et al., 2003) and the main RF glycoprotein SCO-spondin contains multiple anti-adhesive microdomains (Gobron et al., 1996; Monnerie et al., 1998; Meiniel, 2001). Loss of the RF is sufficient to cause aqueductal stenosis and HC (Cifuentes et al., 1994; Perez-Figares et al., 1998; Vio et al., 2000; Wagner et al., 2003). Since the SCO is a specialized type of ependyma, an ependymal cell lineage defect could also be responsible for SCO abnormalities.

The SCO also secretes soluble glycoproteins and is, along with the choroid plexus, the first secretory organ to develop in the brain (Somera and Jones, 2004). SCO glycoproteins would be able to diffuse throughout the ventricles during development, prior to the directional CSF flow that develops after communication between the ventricular system and the subarachnoid space is established (Jones and Sellars, 1982; Somera and Jones, 2004). In support of this theory, the choroid plexus has receptors for SCO-spondin, an important SCO glycoprotein that is also implicated in neuronal maturation (Gobron et al., 1996; Gobron et al., 2000; Meiniel, 2001). Soluble SCO sialoglycoproteins may be important for maintaining intact ependymal cell layers, since loss of ependymal surface sialic acid residues is sufficient to cause detachment, leading to obliteration of the aqueduct and HC (Grondona et al., 1996).

Altered levels of SCO secretions and glycoproteins have been found in CSF from animal models of HC and in human fetuses with HC (Irigoin et al., 1990; Rodriguez et al., 1990; Bonadio et al., 1992; Castaneyra-Perdomo et al., 1994; Castaneyra-Perdomo et al., 2006; Martinez-Pena y Valenzuela et al., 2006; Gonzalez-Marrero et al., 2007). Castaneyra-Perdomo et al (2006) found an additional four RF glycoprotein (AFRU) bands in multiple types of human HC that were not present in normal CSF, while Martinez-Pena y Valenzuela et al (2006) found 5 additional bands in the CSF of spontaneously hypertensive rats with ventricular dilation. SCO AFRU immunoreactivity is also reduced in these rats. In WKY rats, several protein bands (AFRU) are absent from the CSF of both spontaneous and induced HC, and the SCO immunoreactivity for AFRU is also reduced (Gonzalez-Marrero et al., 2007).

Changes in SCO glycoprotein immunoreactivity precede the onset of HC in several experimental models. LEW/Jms rats have a poorly developed SCO and posterior commissure and an abnormally shaped dorsal midbrain. These abnormalities precede closure of the aqueduct and onset of HC (Yamada et al., 1992). In H-Tx rats, reduction in RF glycoprotein immunoreactivity (AFRU) in the SCO is seen at E16, while the closure of the aqueduct and the loss or reduction of SCO ependymal cells does not begin until E17 and the onset of HC does not occur until P0. Loss of glycoprotein staining was always found in HC rats, while closure of the aqueduct and ventricular dilation were not always present (some HC rats had dilated ventricles but open aqueducts, while some had closed or narrowed aqueducts but normal ventricles. However, most had both enlarged ventricles and narrowed or closed aqueduct). Embryos with low AFRU reactivity also showed delayed maturation of the brain prior to E17 (Somera and Jones, 2004). These findings suggest that some pathological process involving the SCO is in place prior to the aqueduct narrowing or closing, and this process is exacerbated or accelerated when the aqueduct closes.

Loss of cilia function. Choroid plexus epithelial cells and the ependymal cells lining the ventricles and circumventricular organs are a mixture of single and multi-ciliated cells. Functions of cilia can vary, depending on their type. Motile cilia are involved in the transport of fluids (Satir and Sleight, 1990) and intracellular signal transduction. A large protein complex, the IFT particle, moves along the cilia axoneme via kinesin II complex (anterograde) and dynein motor proteins (retrograde). This movement is called intraflagellar transport (IFT) and is important in intracellular signal transduction (Kozminski et al., 1993; Rivkin et al., 1997; Pan et al., 2003; Wang and Snell, 2003). Immotile cilia are less



understood, but are believed to regulate ion transport and act as mechanosensors, similar to immotile cilia in the kidney where they detect flow of fluid through the collecting ducts (Lee et al., 1999; Choi et al., 2001; Praetorius and Spring, 2001; Banizs et al., 2005). Immotile cilia also have a possible role in chemoreception. Neuronal primary cilia have olfactory receptors, somatostatin-3 (SST3) receptors and serotonin receptors (Handel et al., 1999; Brailov et al., 2000). All of these functions could have roles in regulating the formation and composition of CSF.

Hydrocephalus frequently is a component of the complex pathologies associated with stunted or absent cilia in animal models (Bruni et al., 1988a; Torikata et al., 1991; Chen et al., 1998; Taulman et al., 2001; Kobayashi et al., 2002; Sapiro et al., 2002; Zhang et al., 2005) and is also seen in human cases of primary ciliary dyskinesia (PCD) (Jabourian et al., 1986; De Santi et al., 1990; Picco et al., 1993; Zammarchi et al., 1993; al-Shroof et al., 2001; Wessels et al., 2003). HC develops in mice lacking regulatory factor X (RFX3), a transcription factor that regulates ciliogenesis. The SCO is also abnormal in these mice, with reduced SCO-spondin expression and undifferentiated ependymal cells (Baas et al., 2006). Mutations in Polaris (IFT88, *Tg737<sup>orpk</sup>*), wimple (IFT172), Spage6, Hfh-4 (a transcription factor), polymerase  $\lambda$ , and kinesin II (Kif3A and Kif3B) complexes all result in cilia dysfunction and HC in mutant mice (Marszalek et al., 1999; Murcia et al., 2000; Nonaka et al., 2008). It remains unclear whether cilia abnormalities result from HC or are a contributing cause (Taulman et al., 2001), although in mice lacking the ependymal-specific axonemal dynein heavy chain *Mdnah5*, loss of ependymal flow (the movement of CSF through the aqueduct) due to dysfunctional motile cilia precedes stenosis of the aqueduct and the postnatal development of HC (Ibanez-Tallon et al., 2004).

In *Tg737<sup>orpk</sup>* mutants, ventricle ependyma have significantly fewer cilia and the cilia that are present are shortened and abnormal (Taulman et al., 2001). However, motile cilia dysfunction is not believed to be the cause of HC in *Tg737<sup>orpk</sup>*, as the pathology develops on P1, prior to motile cilia development (which occurs around P3). Rather, the immotile cilia on choroid plexus epithelial cells likely have a sensory role, similar to primary cilia in renal tubules. Accordingly, CSF from *Tg737<sup>orpk</sup>* mice has a high chloride concentration, indicating a dysfunction in ion transport. An inwardly rectifying chloride channel on the apical surface of choroid plexus epithelial cells is responsible for transporting chloride into CSF. This channel is regulated by cAMP, and intracellular cAMP levels are elevated in choroid plexus epithelium (Banizs et al., 2005). Isolated mutant choroid plexus has lower internal pH and increased  $\text{Na}^+$ -dependent  $\text{HCO}_3^-$  transport activity, resulting in increased CSF production. Inhibiting PKA activity reverses the phenotype to normal; conversely, normal choroid plexus can be converted to mutant phenotype by adding dibutyryl-cAMP (Banizs et al., 2007).

Clearly, the role of cilia dysfunction in HC pathology is far from determined. Interestingly, proteins essential for cilia formation and function are also required for normal cell cycle progression in non-ciliated cells (Mahjoub et al., 2002; Kim et al., 2004; Bahe et al., 2005; Yang et al., 2006). It is tempting to speculate that changes in cell populations observed in HC models could be linked to some of these proteins, but additional research is needed to understand the connection between cilia function and the pathology of HC.

Oxidative stress. Oxidative stress is increasingly recognized as a contributing factor of various neurological disorders and could be responsible for part of the pathology in HC. NO, a free radical, is toxic at elevated concentrations and nNOS (neuronal nitric oxide synthase)

is strongly upregulated in striatum and periventricular regions in affected H-Tx rats (P21) (Socci et al., 1999). It remains unclear if increases in nNOS expression are causative or compensatory. The NOS/NO/cGMP signaling pathway has been shown to inhibit the Na-K-ATPase activity in choroid plexus, implying that CSF production could be changed by increased NO production. Little research has been done to date on the role of reactive oxygen species in HC, so oxidative stress presents an area for further investigation (Johanson et al., 2001; Johanson and Jones, 2001).

Aging. Changes associated with normal aging in the brain may contribute to the development of adult-onset HC. Turnover of CSF declines with age, allowing potentially toxic agents, such as amyloid beta, to accumulate within the brain. This accumulation is enhanced in patients with hydrocephalus and could contribute to cognitive decline. Reduced CSF production associated with aging is caused at least in part by thickening of the choroid plexus basement membrane, reduced perfusion of the choroid plexus and increased vasopressin levels (Serot et al., 1997a; Serot et al., 1997b; Johanson et al., 1999a; Czosnyka et al., 2001; Preston, 2001; Serot et al., 2003). Absorption of CSF also declines with age as the leptomeninges (leptomeningeal fibrosis) and arachnoid villi become thickened, and deposits of amyloid beta in these tissues may further impair absorption of CSF (Bellur et al., 1980; Bech et al., 1997; Hamano et al., 1997; Albeck et al., 1998). Regression of the subcommissural organ is found in human adults, although the functional significance of this finding is not known (Rodriguez et al., 2001). There is also a reduction in the production and circulation of anti-oxidants and free radical scavengers in the CSF, including glutathione peroxidase and vitamins C and E (Cloeze et al., 1989; Tayarani et al., 1989; Czosnyka et al.,

2001; Preston, 2001; Silverberg et al., 2002; Serot et al., 2003; Edwards et al., 2004). All of these changes are typical of the aging brain, and thus do not cause HC in and of themselves. However, there could be underlying factors that make a subset of individuals susceptible to HC, and preventing or reversing age-related changes could be therapeutic in these cases.

Hypertension. Reabsorption of CSF through parenchymal routes (i.e., absorption into venous blood) is now believed to be a significant component of CSF circulation. Hypertension reduces the ability of blood vessels to absorb CSF, causing CSF to accumulate within the brain (Haidri and Modi, 1977; Shukla et al., 1980; Casmiro et al., 1989b; Newton et al., 1989; Bradley et al., 1991; Krauss et al., 1996; Boon et al., 1999; Tullberg et al., 2001). Spontaneously hypertensive rats have progressive ventricular dilation accompanied by reduced subcommissural organ AFRU (glycoprotein) immunoreactivity and altered CSF protein composition (Ritter et al., 1988; Martinez-Pena y Valenzuela et al., 2006). Again, although not every patient with hypertension develops HC, hypertension increases the risk of developing HC. Elevated intracranial pressure can also result from hypertension and cerebrovascular disease. Veins are thought to account for 70-80% of the capacitance in the brain, adjusting blood flow to compensate for increases in intracranial pressure. Hypertension and closure of veins due to cerebrovascular disease thus reduces cerebral compliance. This is of particular concern for patients with compensated HC since cerebrovascular compromise increases the risk of decompensation (Bergsneider et al., 1998; Bateman, 2000; Bradley, 2001a; Czosnyka et al., 2001; Hakim et al., 2001; Yau et al., 2002; Cirovic et al., 2003; Edwards et al., 2004).

Genetic mutations. Mutations in the gene for L1 (L1CAM) are the only genetic mutations known to cause HC in humans. L1 is a neural cell adhesion molecule involved in axon guidance, neuronal migration, signal transduction pathways, and is implicated in learning and memory (Brummendorf and Rathjen, 1995; Scholey et al., 1995; Hortsch, 1996; Fransen et al., 1997; De Angelis et al., 2002). There are at least 70 different mutations of L1 in humans, all of which result in corpus callosum hypoplasia, mental retardation, adducted thumbs, spastic paraplegia and hydrocephalus (CRASH syndrome) (Bickers and Adams, 1949; Fransen et al., 1995; Wong et al., 1995; Kenwrick et al., 1996; Fransen et al., 1997). Other human cases of inherited HC have been documented, but the responsible genes remain unidentified. There are numerous genetic mutations in experimental animals that have HC as part of the resulting phenotype; these may lead to the identification of other genetic HC susceptibilities in humans. There are too many models to cover in detail here, however several well-known and relevant mutations are summarized below.

The suppressor of cytokine signaling (SOCS) family of proteins is involved in the termination of cytokine and growth factor signaling (Starr et al., 1997; Krebs and Hilton, 2000). Approximately 50% of SOCS7 null mice develop HC and die by 15 weeks of age. The SCO is disorganized and is the only apparent abnormality, although the aqueduct was not specifically mentioned as being open and normally shaped (Krebs et al., 2004). Although its specific function is not known, SOCS7 is expressed primarily in the brain and binds to insulin receptor substrate 2 (IRS-2), IRS-4, p85 subunit of PI3K (phosphatidylinositol-3 kinase), vinexin, Nck, Ash, phospholipase C $\gamma$  and epidermal growth factor receptors (Matuoka et al., 1997; Martens et al., 2004). SOCS7 may regulate other tyrosine kinase pathways, including Erk1/2, Akt, c-Kit (Mooney et al., 2001; Bayle et al., 2004). The

appearance of HC in SOCS7 null mice thus supports the evidence for dysregulation of growth factors in the mechanisms of HC.

Pituitary adenylate cyclase-activating polypeptide (PACAP) receptor type 1 (PAC1) is a G protein coupled receptor for PACAP that is expressed throughout the CNS. Both PAC1 null mice and transgenic mice overexpressing PAC1 develop HC (Hannibal et al., 2001; Otto et al., 2004; Lang et al., 2006). In transgenic mice overexpressing PAC1 the severity of HC correlates with the number of transgene copies. Neuronal proliferation in the cortex and subcommissural organ is decreased at E15.5. Other findings include increased apoptosis, reduced corpus callosum, and thinning of the cerebral cortex, particularly layer V.

PAC1 activates both PKA and PKC, and mice lacking the PKC substrates MARCKS (myristoylated, alanine-rich C-kinase substrate), MARCKS-related protein, p190 RhoGAP or the PKA substrate CREB (cAMP response element-binding protein) also have enlarged ventricles, reduced corpus callosum and other neural tube defects, highlighting the importance of these signaling pathways (Chen et al., 1996; Rudolph et al., 1998; Brouns et al., 2000; Huang et al., 2002). Significantly, two of the susceptibility loci for HC the H-Tx rat include the PAC1 and PACAP genes (Jones et al., 2001a; Jones et al., 2001b; Lang et al., 2006). Ependymal cells express PAC1 and tg PAC1 mice have fewer cilia that are shorter and disorganized compared to normal controls. Phospho-CREB is elevated in tg cilia, suggesting disruption of PKA/PKC signaling could be responsible (Lang et al., 2006).

Hydrocephaly with hop gait (*hyh* mouse) is a spontaneous autosomal recessive mutant born with severely dilated ventricles and a small cerebral cortex. Affected mice have a programmed denudation of the ependymal cell layer in the lateral ventricles and aqueduct that corresponds with specific time points in ependymal differentiation (Jimenez et al., 2001;

Wagner et al., 2003). *Hyh* mice have a missense mutation M105I in Napa, the gene for soluble N-ethylmaleimide sensitive factor attachment protein alpha ( $\alpha$ -SNAP). The mutation results in a 40% reduction of  $\alpha$ -SNAP protein levels in homozygous mice. Alpha-SNAP is important for intercompartmental transport of proteins, apical protein localization and cell fate determination; *hyh* mice have abnormal localization of proteins like beta-catenin and E-cadherins (Chae et al., 2004; Hong et al., 2004). This mutation apparently affects ependymal cell lineage, although ependymal detachment may be the result of improper localization of adhesion molecules.

*Hydrocephalus3 (hy3)* mice also have a spontaneous autosomal-recessive mutation that causes lethal HC. The *hy3* mouse mutant is caused by rearrangement in the *Hydin* gene in which exons are rearranged or deleted, causing a premature stop codon. This mutation causes defective CSF reabsorption (Lawson and Raimondi, 1973; Davy and Robinson, 2003). More recently two transgenic mouse lines, *hy3* and OVE459, were generated in which the inserted transgene disrupted two different alleles of *Hydin*. *Hy3* and OVE459 mice have identical pathophysiology to each other and to the spontaneous *hy3* mutant (Lawson and Raimondi, 1973; Johanson et al., 2001; Robinson et al., 2002; Davy and Robinson, 2003). *Hydin* is expressed specifically in ependymal cells of the ventricular system and the *Hydin* gene product may have a role in cilia function (Davy and Robinson, 2003).

Two other genetic mutations of interest are Musashi 1 and RFX4\_v3 knockout mice. Musashi1 protein (*msi1*) is a RNA-binding protein found in ependymal cells and neuronal precursors. Homozygous *msi1*<sup>-/-</sup> mice have abnormal proliferation of ependymal cells surrounding the aqueduct of Sylvius, resulting in stenosis of the aqueduct and obstructive HC. Neuronal precursors appear relatively normal, suggesting that HC again results from an

ependymal cell defect (Sakakibara et al., 2002). Regulatory factor X4 variant 3 (RFX4\_v3) is a transcription factor whose expression is restricted to the brain and regulates the expression of multiple genes critical for development, including Wnt pathway genes and BMP pathway genes (Zhang et al., 2007). Mice lacking a single allele fail to develop the subcommissural organ and have congenital HC. Removing both alleles causes severe midline malformations and death (Blackshear et al., 2003).

### **Pathology of hydrocephalus**

While the exact cause(s) of HC remains unclear, the pathological changes associated with HC are fairly consistent and include physical, motor, cognitive, and behavioral abnormalities. The severity or extent of damage depends upon the age of onset, how quickly ventricles enlarge, and the magnitude of ventriculomegaly.

Loss of periventricular white matter. During the progression of HC, the periventricular white matter begins to atrophy and the corpus callosum and fornix become thinned (Kiefer et al., 1998; Del Bigio et al., 2003). The build-up of CSF in the ventricles causes stretching of the surrounding white matter, damaging the axons, myelin and blood vessels that it contains (Del Bigio, 2001), and cognitive deficits result when the deterioration of the fimbria/fornix connections (connecting the hippocampus to the forebrain) reach a threshold level of atrophy (Gadsdon et al., 1979; Egawa et al., 2002; Del Bigio et al., 2003; Del Bigio, 2004). Although stretching accounts for at least part of the white matter thinning, there is also a reduction in myelin content from the loss of axons and from impaired myelin production that begins prior to axonal damage (Del Bigio et al., 1997b; Del Bigio, 2000). Loss of oligodendrocytes in the



subependymal zone by apoptosis and necrosis could account for the decreased myelination (Del Bigio and Zhang, 1998). In chronic hydrocephalus increased CNPase activity (an oligodendrocyte marker enriched in immature myelin) in periventricular white matter and increased expression of CGaIT (indicative of myelin production) in the dorsal cerebrum myelin suggests that the turnover rate of myelin is increased (Del Bigio et al., 2003). Induction of HC in immature rats delays myelin production, but myelin can be restored if shunting is done before loss of axons occurs (Del Bigio et al., 1997c).

Loss of CSF flow. CSF contains many signaling molecules, growth factors, cytokines, and neurotransmitters that are present at specific times during development and under certain physiological conditions (Kasaian and Neet, 1989; Johnson et al., 1992b; Kitazawa and Tada, 1994; Nogi et al., 1997; Suzaki et al., 1997; Korhonen et al., 1998; Arnold et al., 1999; Ikeda et al., 1999; Riikonen et al., 1999; Grouzmann et al., 2000; Whalen et al., 2000). CSF normally circulates rapidly throughout the ventricular system and reaches most areas of the brain (Proescholdt et al., 2000), making CSF an important communication route. In HC, normal CSF circulation is lost, causing CSF to accumulate in the ventricles. Loss of CSF flow results in abnormal progenitor cell proliferation and accumulation of toxic waste products (Johnson et al., 1992a; Marin-Padilla, 1998; Nicholson, 1999; Mashayekhi et al., 2002). The neurological deficits that remain in cases of early onset HC, even after shunting, may be the result of improper CSF flow (McAllister and Chovan, 1998; Mashayekhi et al., 2002).

Grey matter atrophy. Thinning of the cortex is nearly always seen in both human and experimental cases of HC. While elevated intracranial pressure can cause cortical thinning and defects in progenitor cells are generally thought to be a consequence of HC, abnormal proliferation or differentiation of cortical neurons may also contribute to the pathology of HC. The germinal cells of the subventricular zone may not proliferate or migrate normally if hydrocephalus begins prenatally, impacting normal neuronal layering (Nojima et al., 1998; Khajeh et al., 2002; Owen-Lynch et al., 2003). One group reported that overexpressing cre recombinase in nestin-positive cells (neuronal precursors, radial glia and ependymal cells) resulted in genotoxicity, and the subsequent development of HC was due to impaired proliferation and death of neuronal precursors (Mignone et al., 2004; Forni et al., 2006).

To better understand the loss of grey matter Miyan et al 1998 studied the H-Tx rat at P0, 5, 10, 15, 20 and 25, using nucleic acid staining to compare the germinal matrix (subventricular zone), corpus callosum and frontal cortex of HC rats with normal littermates. Although present at birth, the corpus callosum and germinal matrix are reduced in HC rats at P5 and are nearly gone by P25, while both are still present in normal rats. The frontal cortex, while retaining laminar architecture, appears thinner at birth with a loss of space surrounding nuclei (the authors attribute this to loss of glial processes or to compression) and continues to shrink with time. Apoptosis does not seem to contribute to the loss as the number of apoptotic cells is actually decreased in HC rats in all three regions and at all timepoints. Early shunting of HC rats did not restore normal development, suggesting that elevated intracranial pressure was not responsible for the loss of gray matter (Miyan et al., 1998).

Altered trophic factor levels have been reported in early onset hydrocephalus and may be responsible for abnormal cortical development (Fukumitsu et al., 2000; Miyan et al.,

2001; Mashayekhi et al., 2002). Obstruction of the aqueduct occurs at P18 in the H-Tx rat, coinciding with the peak of neurogenesis and mass migration to the developing cortex; loss of subventricular zone neurogenesis and decreased migration is observed subsequent to aqueductal stenosis. The loss of normal CSF movement may prevent signals necessary for normal development from reaching the subventricular zone, resulting in a thinned cortex due to the loss of neuronal migration to the cortex and/or the loss of astrocytes and oligodendrocytes. (The cell populations primarily affected depends on age of onset. Neuronal progenitors are mostly generated prenatally while oligodendrocyte precursors are generated postnatally in the rat (Mashayekhi et al., 2002; Owen-Lynch et al., 2003; Khan et al., 2006). Progenitor cells apparently migrate prematurely in HC rats, resulting in fewer cells overall being produced in the germinal epithelium. The decrease in proliferation is likely due to factors (either the presence of an abnormal inhibitory factor, the lack of a normal growth factor, or the accumulation of normal growth factors to a concentration at which they become inhibitory) found in the CSF following obstruction of flow since progenitor cells from HC rats are able to proliferate *in vitro* when cultured with normal CSF. Progenitor cells from normal rats survive but do not proliferate *in vitro* in CSF from HC rats, suggesting that the HC CSF is not toxic. Intracranial pressure is not elevated in H-Tx rats and nor is apoptosis occurring, thus the loss of neurons due to decreased proliferation is likely the cause of cortical thinning (Jones, 1985; Miyan et al., 1998; Mashayekhi et al., 2002). Alternatively, Del Bigio et al (2003) found the cortex was thinner in early stages of chronic HC, but the total cerebral volume remained the same, suggesting that cerebral thinning is due mainly to stretching and not tissue loss (Del Bigio et al., 2003). They did not look to see if cell populations or water content were different, however.

Neuronal damage. Axonal damage in periventricular white matter is one of the earliest observations in HC. Periventricular axons are destroyed by mechanical injury from stretching, loss or decrease of cerebral blood flow, and the accumulation of waste products and toxins that accumulate in stagnant CSF (Del Bigio, 1993). Axonal degeneration correlates with ventricle enlargement and becomes more severe over time (Ding et al., 2001a, b). As the lateral ventricles enlarge with accumulating CSF, periventricular axons become stretched, activating L-type calcium channels in axons (Del Bigio, 2000; Johanson et al., 2001). Proteolytic calpains damage cytoskeletal proteins of axons in a calcium dependent manner and are partially responsible for the progressive loss of periventricular axons (Del Bigio, 2000).

Axon degeneration (as seen by silver impregnation in kaolin induced HC) occurs in layers IV-VI of the sensorimotor cortex, neostriatum, hippocampus, subiculum, fimbria, and corpus callosum. Stretching eventually causes disconnection of neurons in the striatum and cerebral cortex, and damage to these structures is associated with deficits in motor learning, memory, and nonverbal cognitive skills (Woody, 1986; Asanuma and Keller, 1991; van der Knaap et al., 1991; Fletcher et al., 1996b; Deiber et al., 1997; Shadmehr and Holcomb, 1997; van der Knaap et al., 1997; Del Bigio and Zhang, 1998). The corticospinal tract of the cervical spinal cord also is damaged in rats with chronic HC and is further evidence of damage to the motor cortex axons (Del Bigio, 2001; Del Bigio et al., 2003).

Disorganization and degeneration of thalamic neurons, subplate neurons (responsible for cerebral cortex and thalamus connections), and pyramidal neurons of the cerebral cortex is common in HC (McAllister et al., 1985; Robertson et al., 2000; Ulfing et al., 2001; Csillik

et al., 2002; Mori et al., 2002; Khan et al., 2006). Neurons, particularly in cortical layers V and VI, may have diminished capacity for activation in progressive HC. Reduced numbers of c-fos-immunoreactive cells and reduced c-fos protein expression is seen as early as P12 in H-Tx rats. Neuronal death does not appear to account for the reduction in c-fos since large numbers of normal-looking neurons are found in the areas of diminished c-fos expression (McAllister et al., 1999). Altered neuronal precursor proliferation and migration, ischemia and deafferentation may account for these changes (Oi et al., 1989).

While axonal damage is common, acute changes and neuronal death are rare in HC (Del Bigio and Zhang, 1998; Ding et al., 2001b, a; Del Bigio, 2004). However, there may be neuronal damage that is subtle but important to pathology (Del Bigio, 1993). Some investigators report finding “dark” neurons, swelling, and vacuolization in a small number of neurons in hippocampus, thalamus, caudate nucleus and in cortical and acetylcholinergic neurons (Gopinath et al., 1979; Wright et al., 1990; Hale et al., 1992; Ulfig, 2002). Other findings include neurofibrillary tangles on cortical and hippocampal neurons, altered expression of synaptic vesicle proteins, and loss of dendritic spines and branches (Ball, 1976; Ball and Vis, 1978; Fan and Pezeshkpour, 1987; Bret et al., 1990; Kriebel et al., 1993; Harris et al., 1996b; Del Bigio and Zhang, 1998; Del Bigio et al., 2003), which may explain the finding of impaired LTP generation in the hippocampus of HC rats (Katayama, 1992).

Ependymal denudation. The ependymal cell layer is lost in portions of the ventricular system in HC. Ependymal denudation is most frequently observed over white matter, particularly the dorsolateral angle and roof of the LV (Lawson and Raimondi, 1973; Page, 1975). In humans, there are case reports of normal, stretched, and completely denuded ependymal

layers throughout the ventricular system (Weller and Shulman, 1972; Bannister and Mundy, 1979; Del Bigio et al., 1985). Ependyma have a limited ability to proliferate, so as the ependymal cells are lost a glial scar usually forms in its place (Del Bigio, 1993). Remaining ependymal cells become flattened, lose their cuboidal shape, appear stretched and lose microvilli, (Price et al., 1976; Weller et al., 1978; James et al., 1980; Rascher et al., 1987; Kiefer et al., 1998). Ependymal denudation may contribute to the loss of directional CSF flow, resulting in stagnation and accumulation of CSF in the ventricles. The rate of ventricular enlargement may determine extent of ependymal loss (Collins, 1979), however as detailed previously there is substantial evidence for ependymal denudation prior to ventricular dilation.

Cerebral compression. Expansion of the ventricular system causes extracellular fluid and blood to be expelled from the brain in order to maintain a normal intracranial pressure (Hakim et al., 1976; Del Bigio, 1993). As extracellular space is compressed, diffusion of water and small molecules (including neurotransmitters) through the cerebral cortex is impinged (Nicholson and Sykova, 1998; Massicotte et al., 2000; Shoesmith et al., 2000; Sykova et al., 2001). Thus compression alters the extracellular environment in grey matter with unknown consequences. The reduced movement of fluids through the brain also causes stagnation in the large extracellular spaces of the periventricular white matter, which are then exposed to abnormally high concentrations of neurotransmitters and waste products found in the CSF of hydrocephalic patients (Lux et al., 1970; Del Bigio, 1989). The morphology of the blood vessels and vasculature is changed by compression, which could further reduce

blood flow and lead to ischemia (Epstein, 1952; Wozniak et al., 1975; Oka et al., 1985; Okuyama et al., 1987; Nakada et al., 1992; Ding et al., 2001a).

Altered cerebral vascularization and metabolism. Reduced cerebral blood flow (CBF) of up to 50% has been demonstrated in hydrocephalic humans and experimental models using PET, SPECT, MRI, CT and Doppler blood flow studies (Goh and Minns, 1995; Shih and Tasdemiroglu, 1995; Nakano et al., 1996). Capillaries are reduced in various experimental models and human cases (Gadson and Emery, 1976; Oka et al., 1985; Higashi et al., 1986; Okuyama et al., 1987; Del Bigio and Bruni, 1988; Sato et al., 1988; Jones et al., 1991; Nakada et al., 1992). In a study designed to compare capillary density in acute vs. chronic HC, Luciano et al (2001) found an initial decrease in cortical capillary density similar to previous studies. However, this acute decrease in capillaries was followed by recovery and increased density over a 12 week period. The functional significance of capillary recovery remains uncertain because cerebral blood flow is still reduced in chronic HC (Higashi et al., 1986; Luciano et al., 2001). HC may cause ischemic preconditioning that is neuroprotective in subsequent ischemic insults (Furuta et al., 1993; Barone et al., 1998; Ding et al., 2001a). This ischemic tolerance may in part explain why more neuronal death is not seen in HC.

Given the reduction in cerebral blood flow, it is not surprising that multiple groups have found changes in oxidative metabolism. During acute HC, increased lactate concentrations and lactate/pyruvate ratios in the frontal lobe, cortex and hippocampus are indicative of increased anaerobic metabolism, and hypoxic metabolites are found in the CSF from HC children (Del Bigio, 1989; Chumas et al., 1994; Braun et al., 1997; Hidaka et al., 1997; Braun et al., 1998; Braun et al., 1999; Socci et al., 1999; Braun et al., 2000; Massicotte

et al., 2000; Kondziella et al., 2002; Kawamata et al., 2003). Glucose metabolism also becomes impaired during HC. The CA3 region of the hippocampus is the first region to show such disturbances, which may contribute to problems with memory in HC patients (Kawamata et al., 2003). Magnetic resonance spectroscopy (MRS) studies with [1-<sup>13</sup>C]glucose and [1,2-<sup>13</sup>C]acetate injections in kaolin-induced HC rats show that multiple [1,2-<sup>13</sup>C]acetate metabolites are significantly changed by week 4, while only one metabolite from [1-<sup>13</sup>C]glucose is changed by week 6. Since acetate is only taken up by astrocytes and glucose is mostly taken up by neurons, it appears that astrocyte metabolism is primarily affected in HC (Kondziella et al., 2003). Additionally, transport of glutamine from astrocytes to neurons in the cerebrum is decreased, which could eventually lead to glutamatergic neurons losing function (Shank et al., 1985).

Changes in neurotransmitter concentrations and intracellular osmolytes are reported by many different groups, however there are considerable discrepancies between studies. Some studies found reduced glutamine, glutamate, aspartate, creatinine, N-acetyl-aspartate, taurine, alanine, dopamine, norpinephrine, GABA, ATP and cAMP levels in H-Tx rats (Miwa et al., 1982; Higashi et al., 1986; Miyake et al., 1992; Minamikawa et al., 1994; Harris et al., 1997; Jones et al., 1997; Tashiro et al., 1997; Kondziella et al., 2002). In others, glutamine, glutamate, aspartate, GABA, norepinephrine, dopamine, noradrenaline, homovanillic acid and serotonin concentrations are increased (Miwa et al., 1982; Del Bigio et al., 1998; Kondziella et al., 2002). Differences in the ages of the animals and time after HC induction may account for some of the contradictory findings between studies.



Choroid plexus and circumventricular organs. Few morphological changes in the choroid plexus are documented in HC. Atrophied epithelial cells, microvilli distortion, vacuolization, intracellular inclusions and enlarged intercellular spaces are generally attributed to reduced production of CSF in an attempt to compensate for increased intracranial pressure (De, 1950; Hochwald et al., 1969; Dohrmann, 1971; Lawson and Raimondi, 1973; Go et al., 1976; Miyagami et al., 1976; Di Rocco et al., 1977; Collins and Woollam, 1979; Liszczak et al., 1984; Preston et al., 2003). More recently elevated cAMP in choroid epithelial cells has been documented in the *Tg737<sup>orpK</sup>* mutant mouse (Banizs et al., 2005). As discussed previously, the subcommissural organ may be smaller, disorganized, or have reduced levels of glycoproteins. Reports of abnormalities in other circumventricular organs include reduced size and increased expression of angiotensin II receptors in the subfornical organ (Acikgoz et al., 1999; Somera and Jones, 2004).

Extracellular matrix. Increased depositions of fibronectin, laminin and vitronectin are seen in the rat model of PHVD and in human cases (Cherian et al., 2004b), particularly around blood vessels. The increased perivascular and parenchymal ECM may impede absorption of CSF, resulting in a build-up of CSF in the ventricles over time.

Changes in glia. A common reaction to CNS damage, astrogliosis is found in the perivascular and periventricular white matter in HC (Del Bigio, 1993; Takei and Sato, 1995), although whether gliosis is beneficial or harmful remains unclear. Elevated levels of GFAP and S-100b in the CSF of HC patients indicate irreversible loss of astrocytes and glia (since these are structural proteins not secreted by glia) (Albrechtsen et al., 1985; Tullberg et al.,

1998; Beems et al., 2003). However, increased subependymal gliosis and proliferation is higher in cats given an effective shunting procedure (compared to no shunt and ineffective shunt), even though the clinical outcome of effectively shunted animals was better (Takei and Sato, 1995). Astrocytes also proliferate and migrate to areas of ependymal denudation, forming a superficial layer. These cells express vimentin, normally only expressed by ependyma, which may indicate they have undergone differentiation (Wagner et al., 2003; Dominguez-Pinos et al., 2005).

Microglial activation in HC has yet to be fully understood. Activated microglia are the brain's macrophages, and increased numbers of activated microglia and clusters of phagocytic microglia were present in the periventricular white matter of P12 mild and severe HC rats and in the cortical grey matter of P21 HC rats (Mangano et al., 1998). Cells positive for the microglial marker mac-H were seen in older HC human fetuses (40 weeks) in the neuropil under areas without denudation and lying on the surface of denuded areas. However, younger fetuses did not have mac-H positive cells (Dominguez-Pinos et al., 2005) and no increases in microglia were found in rats with induced chronic HC (Del Bigio et al., 2003).

Cognitive deficits. The most common complication of HC in human patients is cognitive dysfunction (Young et al., 1973; Raimondi and Soare, 1974; Fletcher et al., 1992b; Fletcher et al., 1992a; Fletcher et al., 1996b; Fletcher et al., 1996a). 68% of shunted HC children have low IQ (Hoppe-Hirsch et al., 1998) and 72% of surviving premature birth infants with HC have “mental deficiencies” (Fernell et al., 1990; Fernell et al., 1991b). Cognitive impairment correlates to ventricle size. Damage to axons in the periventricular white matter

is believed to be the main cause of cognitive deficits (Del Bigio, 1993; Del Bigio et al., 2003; Khan et al., 2006), although abnormal development of some cortical neuronal populations may be responsible for some of the neurological impairments seen in HC children.

Impaired memory retention is another consequence of HC. Patients with chronic HC show signs of memory loss in adulthood (30's and 40's), perhaps caused by fluctuations in intracranial pressure and ventricle size over time (Hagberg and Sjorgen, 1966; Jones et al., 1995b; Scott et al., 1998; Dennis, 2000; Oi et al., 2000; Egawa et al., 2002; Khan et al., 2006). The hippocampus and septohippocampal cholinergic system (SHC) are believed to be critical for spatial learning and memory. Although only subtle changes have been observed in the hippocampus of HC brains, the axons of the SHC are located in structures that are damaged in HC, including the fimbria-fornix, hippocampal projections, and medial septum (Wenk, 1984; Toumane et al., 1988; Kriebel and McAllister, 2000). The alveus, fimbria and fornix are atrophied in humans and there are signs of axonal damage in these structures in HC rats (Del Bigio et al., 2003). Shim and colleagues (2003) found significant loss of cholinergic cells in the SHC pathway in kaolin-injected rats that correlated with severe deficits in learning and memory (assessed by morris water maze & passive avoidance tests to evaluate spatial learning and explicit memory, respectively). Both the loss of cholinergic neurons in the SHC pathway and behavioral deficits correlate with ventricle size, and deficits are worse in chronic HC than in acute HC (Shim et al., 2003). Shunting, if done early enough, can prevent memory deficits, arguing that shunting should be done even in cases where HC seems to have become arrested (Jones et al., 1995a).

Behavioral deficits. Autism and autism spectrum disorders (ASD) are more prevalent in children with HC than in the general population, affecting approximately 20% of HC children as compared to 0.3-0.6% of children without ventriculomegaly (Fernell et al., 1991a, b; Chakrabarti and Fombonne, 2001; Kielinen et al., 2004; Lindquist et al., 2005; Persson et al., 2005; Lindquist et al., 2006). Other behavioral problems are common in children with HC. Unlike the general population, problems are reported equally in boys and girls (normally boys are reported to have more problems) (Connell and McConnel, 1981; Fernell et al., 1991a, b; Fletcher et al., 1995; Williams and Lyttle, 1998). The percentage of HC children with behavioral problems is as high as 60%, whereas only 3-5% of all school age children have reported issues (Rutter et al., 1976), and problems in HC children are reported to be extreme (Lindquist et al., 2006). Behavioral problems in HC children strongly correlate with the degree of cognitive impairment. 60-90% of children with IQ <70 have behavioral problems while 50-20% of children with IQ >70 have problems and only 12% of children with IQ>85 have problems. Preterm birth does not correlate with behavioral problems in HC children, although children with infantile HC are more likely to be hyperactive (Lindquist et al., 2006; Persson et al., 2006).

Motor dysfunction. Most children with HC have some form of motor disability, and gait disturbances are a common feature in adult onset or chronic hydrocephalus (McAllister and Chovan, 1998; Nowak and Topka, 2006; Bugalho and Guimaraes, 2007; Fraser and Fraser, 2007; Factora and Luciano, 2008; McGirt et al., 2008; Mori, 2008; Williams et al., 2008). 78% of premature surviving infants with HC have cerebral palsy and 60% of shunted HC children have motor deficits (Fernell et al., 1990; Hoppe-Hirsch et al., 1998). Like other

impairments caused by HC, motor deficits may correlate with ventricle size. Del Bigio et al (2003) report in rats with induced chronic HC, only animals with the greatest ventricle enlargement displayed motor deficits on the rotating cylinder. However, this was true only during the final (9 month) testing. HC rats also performed significantly more poorly during the 8 month time point but performance did not correlate to ventricle size. HC rats also tended to be less active and had significantly less rearing, but again ventricle size did not correlate with activity (Del Bigio et al., 2003). In the kaolin model, HC rats had readily observable differences in gait and posture but qualitative differences in performance on the rotating cylinder were minimal (Khan et al., 2006). Motor deficits may result from atrophy of the corpus callosum as severe thinning of the corpus callosum has been observed in animal models and humans. Stretching and/or disconnection of axons in the striatum or the cerebral cortex could also result in motor deficits (Gadsdon et al., 1979; Giroud and Dumas, 1995; Del Bigio et al., 2003).

Other impairments. Hormonal imbalances, seizures, sensitivity to sound and light, urinary incontinence, and visual deficits are seen with a higher than normal frequency in HC patients. Impaired depth perception and upward gaze/ocular movement is found in 24-30% of HC children (Zeiner et al., 1985; Hoppe-Hirsch et al., 1998) and 56% of surviving premature infants with HC have epilepsy (Fernell et al., 1990). Reduced body weight and short stature seen in some HC cases are likely due to hormonal imbalances and are possibly related to pituitary gland dysfunction or morphological abnormalities seen in hypothalamic nuclei (Del Bigio, 1993; Lopponen et al., 1996; Yoshino et al., 1999; Abdolvahabi et al., 2000). Rats with chronic HC gain less weight than controls even though they retain similar activity levels

(ie, they are physically able to eat) (Del Bigio et al., 2003). It is possible that stretching of the third ventricle affects the hypothalamus. If so, regulation of appetite or the pituitary gland could result in decreased weight gain.

### **Treatment of hydrocephalus**

Diagnosis. The first diagnosis of hydrocephalus is credited to Hippocrates, who recognized that the enlarged head of some infants resulted from fluid accumulation in the brain and attempted to puncture the fontanelle (Davidoff and Chamlin, 1959). Today, macrocephaly (increasing head circumference) coupled with elevated intracranial pressure symptoms (nausea, bulging fontanelle, split sutures, poor feeding, irritability, headache, lethargy, vision impairment, declining developmental milestones) are indications of HC in infants (Vertinsky and Barnes, 2007). Ultrasound and CT (computed tomography) can detect enlarged ventricles and periventricular edema in fetuses and infants and are used in the initial diagnosis of HC to determine the need and urgency for surgical intervention. Ventricular enlargement is determined by taking the ratio of the diameter of the frontal horn to the maximum brain width (the Evans index). An Evans index greater than 0.3 is considered to be ventriculomegaly (Barkovich and Edwards, 1992; Vertinsky and Barnes, 2007). CT is also useful for follow-up evaluations on shunts and for determining shunt placement (Babcock et al 1988).

MRI (magnetic resonance imaging) has better resolution and contrast than CT and is also capable of detecting tumors that may be blocking the ventricular system. MRI uses no ionizing radiation, gives slice and 3D imaging, and shows circulation and CSF flow (Vertinsky and Barnes, 2007). MRI is used for obtaining anatomical detail for more

complicated surgeries (ie, to tell if CSF is moving through the aqueduct and to assist in catheter placement when cysts or isolated 4<sup>th</sup> ventricles are present) (Gammal et al., 1987; Quencer, 1992; Vertinsky and Barnes, 2007). MRI is also used to obtain detailed 3<sup>rd</sup> ventricle anatomy for endoscopic third ventriculostomy (ETV) surgeries and for determining ETV patency after surgery (Takahashi, 2006). SPECT (single photon emission computed tomography) can be used to evaluate cerebral blood flow and thus help determine if immediate surgical/medical intervention is required in children and adults with ventriculomegaly, but is not used as commonly as CT or MRI (Nayak et al., 2005).

In older children and adults, macrocephaly does not occur due to the hardening of the skull. Primary symptoms of HC in adults are gait disturbance, urinary incontinence, dementia, headaches, and vomiting (Hakim et al., 2001). Demonstration of enlarged ventricles by MRI or CT is used to confirm the diagnosis of HC. Gait disturbance presentation can vary, but typically presents as a slow, shuffling step characterized by a shortened stride, reduced foot-floor clearance and instability (Soelberg Sorensen et al., 1986). Frequent falls are common due to a loss of postural stability (Haan et al., 1987; Blomsterwall et al., 1995). Gait and stability frequently improve after shunt placement (Edwards et al., 2004). Urinary incontinence is due to an increased urgency coupled with difficulty reaching the bathroom because of gait disturbances (Ahlberg et al., 1988; Hakim et al., 2001). Cognitive impairments associated with HC are symptoms of frontal lobe impairment and include inattention, apathy, slowed responses, impaired executive functions, disinhibition, forgetfulness, and excessive sleepiness (Rosen and Swigar, 1976; Masters and O'Grady, 1992; Lindqvist et al., 1993; Chemelli et al., 1999; Iddon et al., 1999; Hakim et al., 2001; Dauvilliers et al., 2003; Lindquist et al., 2005; Lindquist et al., 2008). Mild to moderate

cognitive deficits usually present later, after gait disturbances are manifest, and can fluctuate daily. Severe deficits or dementia that occurs prior to gait disturbances typically are not helped by shunting (Graff-Radford and Godersky, 1986; Iddon et al., 1999).

Surgical treatments. Left untreated, mortality rates of HC are 50-96% (Laurence and Coates, 1962; Yashon, 1963; Eckstein and Macnab, 1966). Surgical treatment of HC involves either redirecting CSF flow by shunt placement or creating a new pathway for CSF flow with endoscopic procedures. Currently shunt placement is the most widely used treatment for HC and is the most common neurosurgery in children (Hoffman and Smith, 1986; Pople et al., 1990; Beni-Adani et al., 2006). Shunts are used in cases of communicating HC and obstructive HC that includes an absorption problem. The first successful valved shunt surgery was described in 1952 by Nulsen and Spitz who diverted CSF into jugular vein. Currently CSF is diverted into the abdomen, lungs, or heart and the use of adjustable valves allows the drainage rate to be tailored to the individual as the rate needed varies widely among patients (Nulsen and Spitz, 1951; Takei and Sato, 1995; Zemack and Romner, 2002; Edwards et al., 2004).

Survival and outcomes of HC have greatly improved with use of shunts, although early placement, preventing severe damage, is best (Scarff, 1963; Takei and Sato, 1995; Del Bigio et al., 1997c). Brain weights and measurements of shunted and normal brains are comparable (Gadsdon et al., 1979). Ventricles become smaller, bulk flow of CSF is restored and edema of periventricular tissues is lost (Del Bigio et al., 1997a; Jones and Andersohn, 1998). Blood flow and metabolism improve, although capillaries generally do not regenerate (Wozniak et al., 1975; Del Bigio and Bruni, 1988; da Silva et al., 1994; da Silva et al., 1995;



Harris et al., 1996a; Nayak et al., 2005). Thinning of the corpus callosum and fornix is stopped, but few, if any, new oligodendrocytes are generated (Del Bigio et al., 2003; Fukushima et al., 2003). There are improvements in learning ability and behavior (Del Bigio et al., 1997a; Hawkins et al., 1997). In cases of chronic HC, 60-80% of patients given a shunt have clinical improvement as evidenced by a decrease in CSF outflow resistance and mean intracranial pressure (Vanneste, 2000; Petrella et al., 2008).

While shunt placement has greatly improved the life expectancy of HC patients, it also has the highest complication rate of all pediatric neurosurgeries (Renier et al., 1984; Walters et al., 1984; Oi and Matsumoto, 1985; Ammirati and Raimondi, 1987; Pople et al., 1990; Sainte-Rose et al., 1991; Pople, 1992; Bajpai et al., 1997; Baskin et al., 1998b). In the US, \$100 million/year is spent on shunt procedures, and nearly half are revision surgeries (Bondurant and Jimenez, 1995). 70% of shunts fail within 10 years, and most children have multiple replacements, with a revision average of 1.6 to 3.6 revisions per patient (Tung et al., 1991; McCallum and Turbeville, 1994). Shunts can fail due to mechanical malfunction, infection, obstruction, migration, disconnection, or inadequate length for a growing child. Ventriculovascular shunts may develop clots in the catheter or blood vessel. Along with mechanical failure, infection is the most serious complication. Infection occurs in 2-39% of shunts (Yogev and Davis, 1980; Nelson et al., 1984; Blount et al., 1993). A majority (60%) of infections are caused by *Staphylococcus epidermidis* and *S.aureus* and 90% of infections occur within 6 months of placement, indicating that infection occurred at time of placement (Schoenbaum et al., 1975; Ersahin et al., 1994; Ronan et al., 1995; Turgut et al., 2005). Shunt obstruction is most common in cases of HC caused by tumors; other sources of obstruction include choroid plexus tissue, reactive glia, ependyma and cellular debris

(Lazareff et al., 1998). Overdrainage, slit ventricle syndrome, and craniosynostosis are also possible complications of shunts (Kang and Lee, 1999; Virella et al., 2002; Martinez-Lage et al., 2005; Daszkiewicz and Barszcz, 2007; Clark et al., 2008; Rekate, 2008).

In addition to surgical risks, shunt placement does not address all aspects of HC pathology. The ependymal layer does not regenerate (Del Bigio, 1993; Kiefer et al., 1998), and damaged axons generally cannot be restored (Del Bigio, 2001). There is some recovery in grey matter, but little repair of axons in white matter (Aoyama et al., 2006).

Periventricular gliosis remains (Takei and Sato, 1995) and may contribute to the observation of stiffened ventricular walls, resulting in incompliance to changes in pressure (Kiefer et al., 1998). Additionally, the developmental consequences of removing or altering the composition of CSF are unknown.

Endoscopic procedures are used to treat obstructive, non-communicating HC by creating new pathways for CSF flow. Because they do not address problems involving reabsorption of CSF, endoscopic surgeries are generally not effective for treatment of communicating HC. Third ventriculostomy (ETV) is the most common endoscopic treatment and provides a bypass around the site of blockage by making an opening in the floor of the third ventricle. Success depends on the cause of HC. ETV are 70-85% successful in congenital HC cases with aqueductal stenosis or tumor, but are only successful in half of cases where HC is due to intraventricular hemorrhage or infection (Grant and McLone, 1997; Buxton et al., 1998; Buxton et al., 1999; Cinalli et al., 1999; Hopf et al., 1999; Fukuhara et al., 2002).

A successful ETV allows the patient to avoid shunt dependence and the accompanying complications as there is little to no risk of overdrainage, no foreign matter

left in the brain, no tubing to become blocked or fail, and low risk of infection. However, ETV have a high failure rate in infants (30-90% require shunts later) due to ongoing problems with CSF absorption. Success is better in older children, although the exact reasons are unclear (Baskin et al., 1998a; Cinalli et al., 1998; Cinalli et al., 1999; Siomin et al., 2001; Beems and Grotenhuis, 2002; Edwards and Pople, 2003; Rekate, 2004; Baykan et al., 2005; Dusick et al., 2008). Better classification of HC should help in determining which cases (ie, purely obstructive) would benefit from endoscopic treatment (Beni-Adani et al., 2006). Additionally, ventricular volume post ETV does decrease but remains higher than normal population, and there are risks of uncontrolled bleeding during operation and failure leading to death (Cinalli et al., 1999; Hader et al., 2002; Feng et al., 2004).

A new experimental treatment for preterm infants after intraventricular hemorrhage, called DRIFT (drainage, irrigation and fibrinolytic therapy), shows promise for preventing post-IVH HC. The theory behind DRIFT is to wash out as much blood, cytokines and free-radical causing agents (like excessive iron) from the ventricles as possible before ventricular dilation and HC can occur. Infants are also given plasminogen activator prior to irrigation with aCSF. A small pilot study was highly successful, but large-scale and long-term studies have yet to be completed (Cherian et al., 2004a).

Pharmacological treatments. There are no current pharmacologic therapies for HC, although there have been multiple attempts. Drugs designed to inhibit CSF production include acetazolamide, furosemide, isosorbide, and ouabain. Acetazolamide (Diamox) is a carbonic anhydrase inhibitor administered in the 1950's and reduces CSF production by as much as 60% *in vitro* and in experimental models. However there was no clinical improvement in HC

symptoms (Elvidge et al., 1957). Furosemide is a diuretic and acts by inhibiting renal sodium reabsorption. Again no improvement of HC symptoms was seen in a randomized drug trial (Shinnar et al., 1985; Witte et al., 1986; Kennedy et al., 2001; Whitelaw et al., 2001; Horinek et al., 2003). Isosorbide is a hyperosmolar agent that was promoted in the 1970's, but has not gone through randomized testing and lacks evidence for clinical efficacy (Lorber, 1975; Lorber et al., 1983; Del Bigio, 2004). Ouabain is a sodium-potassium ATPase pump inhibitor, but has widespread toxicity and thus is not suitable as a pharmacological treatment (Srebro, 1969; Gueli et al., 1976).

Agents designed to restore blocked CSF pathways also show little promise. Intrathecal injections of hyaluronidase did not show improvement (Schoeman et al., 1991). Urokinase, streptokinase, and tissue plasminogen activator were tested in hopes that they could break up blood clots following intraventricular hemorrhage, but had minimal success (Hansen et al., 1997a; Hansen et al., 1997b; Whitelaw, 2001; Whitelaw et al., 2001; Yapicioglu et al., 2003). Reducing inflammation of the meninges seems like a logical way to improve CSF absorption, but treatment with anti-inflammatory agents was also met with minimal success (Wilkinson et al., 1974; Erler and Klaber, 2001).

The most promising treatments for HC involve neuroprotective agents. Nimodipine is an L-type calcium channel antagonist that improves cerebral blood flow in models of ischemic injury (Scriabine and van den Kerckhoff, 1988; Scriabine et al., 1989; Schmidt et al., 1990; Welty and Horner, 1990; Feigin et al., 1998) and may help in HC by preventing the activation of calpains in axons that occurs during stretch injury, thus preventing proteolytic axonal damage (Maxwell et al., 1991; Povlishock and Jenkins, 1995; Stys, 1998; Buki et al., 1999; Del Bigio, 2000). In the kaolin rat model of HC, nimodipine treatment for 2 weeks

(started 2 weeks after induction of HC) prevented motor and cognitive deficits and preserved white matter (Del Bigio and Massicotte, 2001), although the myelin content and ventricle size did not differ from untreated controls. L-type calcium channels are also needed for synaptogenesis, so more work needs to be done to determine safe dosage levels (Vigers and Pfenninger, 1991; Kirsch and Betz, 1998; Del Bigio and Massicotte, 2001).

Magnesium sulfate, a calcium antagonist that blocks NMDA channel receptors and voltage and receptor-operated calcium channels, is another agent with neuroprotective potential. In experimental models of spinal cord injury, magnesium sulfate reduces lipid peroxidation, hypoxic-ischemic damage and the size of traumatic injury lesions (Simpson et al., 1994; Marinov et al., 1996; Suzer et al., 1999; Sameshima and Ikenoue, 2001; Vink et al., 2001). In the kaolin rat model (2 weeks of treatment given 2 weeks after induction), rats had reduced GFAP levels in the cerebrum but showed no improvement on the rotarod test, and higher doses produced significant sedation (Khan et al., 2003). Magnesium sulfate has been used in managing pregnant women with eclampsia, but recent studies show an association between its use and poor neurologic outcomes for the infants (Nelson and Grether, 1995; Mittendorf et al., 2002). Thus magnesium sulfate is not a viable HC treatment.

Mexiletine and riluzole are sodium channel blockers that showed some potential in various neurological injury models but also provided no protective effects in kaolin rat model of HC (Del Bigio et al., 2002).

### **Experimental hydrocephalus models**

Induced models. Experimental models of HC have been created using a variety of substances injected into the ventricular system to block CSF flow. The first model of HC

involved inserting a cotton plug into the aqueduct of dogs (Dandy, 1919). Since then, many other agents have been used to induce HC, including blood (Bachs and Walker, 1953), India ink, cyanoacrylate glue, balloons, silicone oil, neuraminidase, infectious agents, and kaolin. Typically these are inserted or injected into the aqueduct, lateral, or fourth ventricle. Silicone oil is injected into the cisterna magna and produces a sterile mechanical obstruction to CSF out of the fourth ventricle (Wisniewski et al., 1969; Johnson et al., 1999). Neuraminidase is injected into the lateral ventricles and causes HC by cleaving surface sialic acid residues from the surface of ependymal cells, causing detachment and subsequent obliteration of the aqueduct (Grondona et al., 1996). Multiple agents have been used to cause meningitis and subsequent HC, including mycobacterium tuberculosis (De, 1950), influenza (Margolis and Kilham, 1977), mumps (Johnson and Johson, 1968), reovirus (Margolis and Kilham, 1969; Nielsen and Baringer, 1972), MHV3 mouse hepatitis virus 3 (Tardieu et al., 1982), and RS human respiratory syncytial virus (Lagace-Simard et al., 1982). In addition to inflammation, some of these agents also cause ependymal damage, denudation and aqueductal stenosis. A complication of most of these induced methods is the presence of an inflammatory response or physical tissue distortion that is not present in most cases of HC, thus the relevance of findings from models with inflammation is not clear.

Kaolin injection into the cisterna magna is an inexpensive and simple model of chronic noncommunicating hydrocephalus (Del Bigio and Zhang, 1998; Del Bigio et al., 2003; Khan et al., 2006). Kaolin causes sterile inflammation of the meninges, blocking the flow of CSF from the ventricular system to the intracranial and spinal subarachnoid space (the foramina of Luschka and Magendi are obstructed) (Kondziella et al., 2002). The development of HC takes place in several phases. The acute phase occurs after kaolin

injection and lasts for approximately four weeks and is characterized by maximal intracranial pressure and resistance to CSF outflow. After the fourth week, resistance and intracranial pressure drop, but ventricles continue to expand. This intermediate phase lasts 6-8 weeks, after which the model resembles chronic or normal pressure HC, although some believe it is most similar to postmeningitis HC in human infants (Oi et al., 1996; Kondziella et al., 2002).

Hydrocephalic rats have atrophy of periventricular white matter, reactive astrocytes, damaged neuronal axons and glial cell death. Deficits in motor and cognitive function are associated with damage to the corpus callosum, corticospinal tract and hippocampal fimbria/fornix projections. Severity of these deficits is related to the age of onset and the rate at which hydrocephalus progresses (Del Bigio and Zhang, 1998; Del Bigio et al., 2003). Severely HC rats have a learning disability in water maze. Early shunting can prevent this, while late shunting shows gradual but incomplete restoration of learning ability (Overholser et al., 1954; Del Bigio et al., 1997c).

HC can also be induced by altering nutrition. When female rats are fed a diet lacking in vitamins B12 or folic acid, they give birth to pups with HC. Embryos have stenosis of the aqueduct that begins between E16 and E18, with ventricle enlargement and HC beginning at E18 (Overholser et al., 1954; Newberne and O'Dell, 1959). Embryos from folic acid-deficient mothers also have loss of ependyma in the cerebral aqueduct.

Spontaneous models. Several spontaneously occurring congenital HC models are also commonly studied. One highly studied model is the H-Tx rat. Approximately 33% of H-Tx rats develop severe HC in late gestation following subcommissural organ abnormalities and obliteration of the cerebral aqueduct at E18 (Jones and Bucknall, 1988; Somera and Jones,

2004). Genetic studies show HC results from complex genetic factors with incomplete penetrance. Several loci for putative HC gene mutations have been identified, including one located near the gene *Igfb4* (insulin-like growth factor binding protein) (Jones et al., 2000). Approximately 15% of affected H-Tx rats have symptoms of HC after birth but live a normal life span. These are called hH-Tx rats, and are a model of compensated or spontaneously arrested HC. hH-Tx rats have a slow loss in learning ability as they age, similar to human cases of chronic HC (Kiefer et al., 1998; Nonaka et al., 2008).

*Hyh* mice are another spontaneous model that has been extensively researched. *Hyh* mice have a point mutation in the Napa gene encoding alpha-SNAP, causing a 40% reduction in protein levels (Chae et al., 2004; Hong et al., 2004). A four year study shows that 22.4% of *hyh* mice develop HC; 70% of these progress rapidly and die within 2 months while the rest survive up to 2 years (with no motor deficits and little to no enlargement of the head). The survivors develop spontaneous ventriculostomies that allow CSF to drain from the ventricles directly into the subarachnoid space (Batiz et al., 2006).

Other spontaneous models have been used in HC research but less is known about their pathology. In the SUMS/NP mouse, lateral ventricle enlargement begins on E15, the cerebral aqueduct is stenosed or obliterated prenatally, and ependymal development in the aqueduct is abnormal with dysfunctional cilia and discontinuous ependymal detachment. The cause for HC in this model has yet to be identified (Jones et al., 1987; Bruni et al., 1988a). The LEW/Jms rat also has aqueductal stenosis from unknown causes (Yamada et al., 1992). *Ch* mice have a point mutation in *Mf1* (Foxc1), a forkhead/winged helix gene. The resulting mutant protein lacks its DNA-binding domain (Kume et al., 1998) and a defective chondroitin sulfate proteoglycan is produced (Green, 1970; Gruneberg and Wickramaratne,



1974; Richardson, 1985). Hydrocephalus begins on E12, possibly due to abnormal subarachnoid space or loss of ependymal cells since the cerebral aqueduct remains patent (Green, 1970; Gruneberg and Wickramaratne, 1974). In WIC-Hyd rats an unknown genetic mutation causes cilia dysfunction and HC (Torikata et al., 1991; Nakamura and Sato, 1993). Other spontaneous HC models include *hop sterile* (hop) mice and *obstructive hydrocephalus* (oh) mice.

Transgenic models. Numerous lines of transgenic animals develop HC for a variety of known and unknown reasons. HC develops in two separate lines of mice lacking the L1 neural adhesion molecule, and severity ranges from mild ventriculomegaly to severe HC (Dahme et al., 1997; Fransen et al., 1998). The aqueduct is patent at early stages of HC but becomes stenosed as HC progresses (Rolf et al., 2001). Hypoplasia of the corticospinal tract is a consistent finding and mirrors the human CRASH syndrome (Fransen et al., 1995; Cohen et al., 1998; Fransen et al., 1998). A mutation in *hydin*, a cilia protein expressed exclusively in ventricular ependymal cells, causes HC in *Hy3* mice (Davy and Robinson, 2003). Ependymal denudation precedes stenosis of the aqueduct, which occurs late in the pathogenesis (Raimondi et al., 1976; Jimenez et al., 2001).

Not all transgenic models have aqueductal stenosis. Two lines of mice over-expressing TGF- $\beta$ 1 in GFAP positive cells develop progressive communicating HC and have increased depositions of perivascular extracellular matrix proteins (Galbreath et al., 1995; Wyss-Coray et al., 1995; Cohen et al., 1999), reduced LTP in hippocampal slices, and impaired spatial learning in Morris water maze (Gerlai et al., 1995). The conditional knockout of *Cnp1* in oligodendrocytes results in neuronal degeneration in white matter, HC

and early death (Lappe-Siefke et al., 2003). Interestingly, myelin and myelination are normal, which shows that oligodendrocytes have axonal support functions beyond myelination. The function of CNP (2',3'-cyclic nucleotide 3'-phosphodiesterase) in oligodendrocytes is unclear, but it can interact with cytoskeletal proteins (Laezza et al., 1997; Bifulco et al., 2002). Both full knockouts and mice with reduced expression of nonmuscle myosin heavy chain II-B (NMHC-B, involved in cell polarity, cell shape, cell motility, cytokinesis) results in HC. Severity correlates with the level of gene expression. Ependymal cells denude and the aqueduct is malformed, likely due to weak cell-cell interactions of neuroepithelial cells and possibly resulting in improper differentiation. There are defects in neural cell migration as well (Uren et al., 2000). The conditional knockout of *Hif-1 $\alpha$*  in nestin expressing (neural) cells results in HC and impaired memory (Tomita et al., 2003). HIF-1 $\alpha$  is a hypoxia-inducible factor involved in cell survival during ischemia, and also has role in vascular development. Null mutants of *Spag6*, *Hfh4*, *Pol  $\lambda$* , *Polaris*, *Mdnh5* all have cilia dysfunction and HC (Chen et al., 1998; Taulman et al., 2001; Ibanez-Tallon et al., 2002; Kobayashi et al., 2002; Sapiro et al., 2002).

Hydrocephalus has long been recognized as a serious neurological disorder. Modern advances in imaging techniques and biomedical technology has made a tremendous impact in the treatment of patients with the disorder. Surgical treatments have provided a greatly prolonged life expectancy for persons with HC, but their quality of life could be improved if pharmacological treatments for reversing or preventing HC were available. As summarized above, HC has a complicated etiology and past attempts for drug therapies were only marginally successful at best. Experimental HC models have been useful in identifying common features of HC, including abnormalities in ependymal cells, cilia, the

subcommissural organ, and neuronal progenitor cells. Yet the question of cause and effect remains for many of these mechanisms, and the current experimental models have not been able to effectively address these issues. Many models cannot predict which animals will develop HC, making the distinction between early causative events and pathological responses very difficult. Other models are complicated by the presence of pathologies distinct from HC, again making the identification of the molecular mechanisms of HC tricky. Further advances in HC treatment will be aided by a model that allows for control of HC onset and accurate prediction of affected animals.

### **CHAPTER 3**

#### **A NEW GENETIC MODEL OF HYDROCEPHALUS**

We developed a transgenic mouse line that expresses the Gi-coupled RASSL (Receptor Activated Solely by Synthetic Ligand) Ro1 in GFAP-positive cells with the aim of studying the role of GPCR signaling in astrocyte-neuronal communication. Surprisingly, we found that all transgenic mice expressing Ro1 developed hydrocephalus. We analyzed these mice in an effort to develop a new model of hydrocephalus that will further our understanding of the pathophysiology of the disease.

Expression of Ro1 was restricted to glia by crossing the transgenic hGFAP-tTA (tet transactivator behind the human glial fibrillary acidic protein promoter) mouse line with the transgenic tetO-Ro1/tetO-LacZ mouse line. This cross produced double transgenic mice that expressed Ro1 only in glia. All double transgenic mice developed hydrocephalus by p15, while single transgenic littermate controls appeared normal. Hydrocephalic double tg mice displayed enlarged ventricles, partial denudation of the ependymal cell layer, altered subcommissural organ morphology, and obliteration of the cerebral aqueduct. Severely hydrocephalic mice also had increased levels of phosphoErk and GFAP expression.

Administration of doxycycline to breeding pairs suppressed Ro1 expression and the onset of hydrocephalus in double transgenic offspring. Double transgenic mice maintained on dox never developed hydrocephalus, however if they were taken off doxycycline at weaning, double transgenic mice developed enlarged ventricles within seven weeks,

indicating that Ro1 expression also induces hydrocephalus in adults. Signaling through Ro1 is required for hydrocephalus to occur since both ventricular enlargement and ependymal denudation were prevented by administering a KOR inverse agonist to double transgenic mice taken off doxycycline. This study discovered a new model of hydrocephalus in which the rate of pathogenesis can be controlled, enabling the study of the pathogenesis of both juvenile and adult onset hydrocephalus.

## **Materials and methods**

Animals. All experiments were performed in accordance with guidelines set forth by the Institutional Animal Care and Use Committee at the University of North Carolina at Chapel Hill and federal guidelines. Animals were maintained in climate-controlled housing with a 12 h light/dark cycle and were given food and water *ad libitum*. The day of birth was defined as postnatal day 0 (P0).

Transgenic mouse lines. The B6(C3)-Tg(GFAP-tTA)96Kdmc mouse line was made by isolating the tTA gene from pUHD 15-1 (kindly provided by Hermann Bujard, Universität Heidelberg, Heidelberg, Germany), and cloning it downstream of the 2 kb human GFAP promoter in place of lacZ in the pGfa2lac1 vector (kindly provided by Michael Brenner, University of Alabama at Birmingham). The GFAP-tTA-mP1 cassette was placed between four copies of genomic insulators from chicken  $\beta$ -globin gene (Chung et al., 1993).

Transgenic mice were obtained by standard methods and backcrossed to C57BL/6J for at least five generations. FVB/N-TgN(tetO-Ro1-LacZ)CONK mice were obtained from The Jackson Laboratory (Bar Harbor, ME) and crossed to GFAP-tTA mice to get double-

transgenic mice. KOR knock-out mice (Hough et al., 2000) were obtained courtesy of Dr. John E. Pintar (University of Medicine and Dentistry New Jersey, Piscataway, NJ).

Genotyping. Polymerase chain reaction (PCR) was used to screen mice for transgene presence. For B6(C3)-Tg(GFAP-tTA) transline, the following primers were used: 5' CCCTTGGAATTGACGAGTACGGTG 3' (forward) and 5' TGGTGTATGAGCGGCGGCGACGGCAG 3' (reverse). For FVB/N-TgN(tetO-Ro1-LacZ)CONK transline, primers against the LacZ gene were utilized: 5' ATCCTCTGCATGGTCAGGTC 3' (forward) and 5' CGTGGCCTGATTCATTCC 3' (reverse).

Southern blots. To identify KOR KO mice, genomic DNA was digested with BamHI overnight at 37°C. DNA fragments were separated on a 0.8% agarose gel (95 volts for 3.5 hours) and transferred to Zeta-Probe GT membrane (Bio-Rad Laboratories, Hercules, CA) using an alkaline transfer method. DNA was immobilized by UV crosslinking. The <sup>32</sup>P-labelled hybridization probe was made using the Megaprime<sup>TM</sup> DNA labeling kit (Amersham Life Science, Buckinghamshire, England) and 100 ng of the 0.8 kb XbaI/EcoRI fragment of kappa genomic clone from plasmid GC#3 (kindly provided by John Pintar, University of Medicine and Dentistry New Jersey, Piscataway, NJ). The membrane was hybridized in QuikHyb buffer (Stratagene Cloning Systems, La Jolla, CA) and exposed to Kodak Biomax Film overnight at -70° C.

Doxycycline administration. Selected animals were given 25 or 50 µg/ml doxycycline hyclate (dox; Sigma, St. Louis, MO) in the drinking water. Amber bottles wrapped in aluminum foil were used to protect the dox from light and the water was changed once a week. Breeding

pairs on dox were given dox water at the time of mating and maintained on dox continuously. Breeding pairs off dox were never exposed to dox. For timed studies, mice were taken off dox at P30.

Spiradoline administration. Spiradoline mesylate (0.6  $\mu\text{g}/\mu\text{L}$  in ddH<sub>2</sub>O, U-62066, Sigma, St. Louis, MO) was given to mice (3 mg/kg, i.p.) 15-30 minutes before behavioral testing or euthanizing.

NorBinaltorphimine administration. nor-Binaltorphimine dihydrochloride (Tocris, St. Louis, MO) was suspended in PBS to make a 1 mM solution and aliquots were stored at -20°C. Mice were injected with norBIN (20 mg/kg, s.c.) every three days, starting with the day doxycycline was removed from the drinking water. For behavioral tests, mice were injected with norBIN 18-24 hours prior to testing.

Immunofluorescence. Adult (P45–P60) mice were deeply anesthetized and transcardially perfused with cold 4% paraformaldehyde/0.1 M NaPO<sub>4</sub>, pH 7.4. Brains were removed and postfixed overnight at 4°C in the same buffer. To detect the Ro1 receptor, sections were probed for FLAG, the peptide (DYKDDDDK) epitope tag fused to the N terminus of Ro1. For FLAG staining, 50  $\mu\text{m}$  sections were cut on a vibratome, washed in alternating PBS and PBS/0.5% Triton X-100 (Tx-100) washes, and blocked in PBS/0.5% Tx-100/10% NGS/0.2% gelatin/3% milk. Sections were incubated with primary antibodies (rabbit anti-FLAG; Sigma) diluted in PBS/0.5% Tx-100/3% NGS/0.2% gelatin overnight at 4°C with gentle agitation. Sections were extensively washed in PBS and PBS/0.2% Tx-100. Secondary antibodies (goat anti-rabbit IgG conjugated to Alexa-488; Invitrogen, Carlsbad, CA) were diluted in PBS/0.5% Tx-100/5% NGS/0.2% gelatin and applied for 2–4 h at room temperature. For

GFAP and NeuN staining, fixed brains were cryoprotected in PBS/30% sucrose overnight at 4°C. Sections (14 µm) were cut on a cryostat, blocked for 2–4 h in PBS/20% normal goat serum/2% BSA/0.2% Tx-100 at room temperature, and incubated overnight with primary antibodies (mouse anti-GFAP, 1:500, Sigma; rabbit anti-GFAP, 1:500, Roche, Mannheim, Germany; mouse anti-NeuN, 1:1000, Chemicon, Temecula, CA; rabbit anti-ERK1/2 and anti-phosphoERK1/2, Cell Signaling, Beverly, MA; mouse anti-acetylated tubulin, Sigma, St. Louis, MO; mouse anti-S100β, Sigma, St. Louis, MO; anti-cleaved caspase 3, Cell Signaling, Beverly, MA) diluted in the same buffer. After washing, sections were incubated with secondary antibodies (goat anti-mouse conjugated to Alexa-488 and goat anti-rabbit conjugated to Alexa-594; Invitrogen) diluted 1:400 in blocking buffer for 2–4 h at room temperature.

Ependymal cell dissection. Adult mice were anesthetized and euthanized by decapitation. Coronal sections (400 µm) containing the lateral ventricles (usually four sections) were cut on a vibratome (Leica VT 1000S; Leica, Nussloch, Germany). Using a dissecting microscope, fine point microdissection tweezers (World Precision Instruments, Sarasota, FL) and 30 gauge needles, the ependymal cell layer(s) were teased away from the underlying tissue.

Xgal histochemistry. Adult mice (P30–P45;  $n = 6$ ) were deeply anesthetized and transcardially perfused with cold 4% paraformaldehyde/0.1 M NaPO<sub>4</sub>, pH 7.4. One hundred micrometer sections were cut on a vibratome. Sections were washed three times (10 min each) at room temperature in Xgal rinse buffer (0.1 M KPO<sub>4</sub>, pH 7.4, 2 mM MgCl<sub>2</sub>, 0.01% sodium deoxycholate, 0.2% Igepal CA-630) and stained overnight at 37°C in Xgal stain



buffer [rinse buffer plus 1 mg/ml 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside, 5 mM  $K_3Fe(CN)_6$ , and 5 mM  $K_4Fe(CN)_6$ ].

Immunoprecipitation/Western blotting. Fresh brain tissue (P21–P50;  $n = 15$ ) was homogenized in ice-cold lysis buffer (50 mM Tris-HCl, pH 7.4, 100 mM NaCl, 5 mM EDTA, 50 mM NaF, 10 mM NaPP, 200  $\mu$ M  $Na_3VO_4$ , 1% Tx-100, 100  $\mu$ g/ml PMSF, 1  $\mu$ g/ml leupeptin, 2  $\mu$ g/ml aprotinin, 1  $\mu$ g/ml soybean trypsin inhibitor, 1  $\mu$ g/ml pepstatin A, 10  $\mu$ g/ml benzamidine) and sonicated on ice. Lysates were incubated on a rotator for 1 h at 4°C. Supernatants were precleared and incubated for 2 h at 4°C with M2 FLAG antibody (Sigma) and protein A/Sepharose beads (Sigma) that had been blocked with BSA. Beads were boiled in Laemmli buffer, run on an SDS-PAGE gel, and blotted onto a nitrocellulose membrane. The blots were blocked in 5% milk–TBS/0.1% Tween 20 (TBST) for 2 h at room temperature and probed overnight at 4°C with a 1:250 dilution of a rabbit anti-FLAG polyclonal antibody (Sigma) in 5% milk/TBST. Membranes were washed in TBST, probed with HRP-conjugated goat anti-rabbit IgG (1:10,000), and processed for ECL. For GFAP, fresh brains (P21–P35;  $n = 6$ ) were dissected in ice-cold PBS, and tissues were homogenized in a volume of homogenization buffer (50 mM HEPES, pH 7.3, 150 mM NaCl<sub>2</sub>, 1.5 mM MgCl<sub>2</sub>, 1 mM EGTA, 10% glycerol, 1% Tx-100 with Complete protease inhibitor (Roche Diagnostics, Mannheim, Germany) equal to 10x tissue weight. The samples were sonicated and centrifuged. Equal amounts of protein were run on a Tris-glycine gel, blotted onto a nitrocellulose membrane, and probed with rabbit anti-GFAP antibody (DakoCytomation; 1:5000) for 48 h at 4°C.

Hematoxylin and eosin staining. Brains from perfused animals (P30–P66;  $n = 14$ ) were fixed in 10% formalin overnight, rinsed in distilled water, and stored in 70% ethanol. The fixed brains were taken to the University of North Carolina histology core for paraffin embedding. Sections (6  $\mu\text{m}$ ) were cut on a sliding microtome. Sections were incubated in xylenes to remove paraffin, rehydrated in serial ethanol washes, and incubated for 6 minutes in hematoxylin (Sigma, St. Louis, MO). After rinses in 0.25 acid alcohol and lithium carbonate, sections were counterstained in eosin Y solution (Sigma, St. Louis MO), dehydrated in serial ethanol washes, cleared in xylenes, and mounted in Cytoseal<sup>TM</sup> XYL (Richard-Allan Scientific, Kalamazoo, MI).

Immunocytochemistry. Brains from adult mice (P30–P45;  $n = 11$ ) were fixed in 4% paraformaldehyde, cryoprotected in PBS/30% sucrose, and cut in 35  $\mu\text{m}$  sections on a cryostat. Sections were rehydrated in PBS and incubated in 100% methanol at  $-20^{\circ}\text{C}$  for 10 min. Sections were washed three times (10 min each) in TBST and once in TBS before incubating for 2 h in blocking buffer (5.5% NGS/5.5% BSA/TBST) at room temperature. Avidin/biotin was blocked using an avidin/biotin blocking kit (Vector Laboratories, Burlingame, CA) according to the manufacturer's protocol. Primary antibodies [rabbit anti-phospho-p44/42 mitogen-activated protein (MAP) kinase (Thr202/Tyr204); Cell Signaling Technology, Beverly, MA; 1:250; or mouse anti-GFAP; Sigma; 1:500] were diluted in blocking buffer, and sections were incubated overnight at  $4^{\circ}\text{C}$ . Biotinylated goat anti-rabbit secondary antibody (Vectastain Elite ABC kit; Vector Laboratories) was diluted 1:500 in TBS/3% BSA, and sections were incubated for 2 h at room temperature. Sections were then incubated for exactly 30 min in 0.6%  $\text{H}_2\text{O}_2$  at room temperature. ABC reagent was prepared according to kit instructions, and sections were incubated for 2 h at room temperature. DAB

reagent (Vector Laboratories) was added, and the reaction was allowed to proceed until sufficient staining developed. Coverslips were mounted using Vectashield with DAPI (4',6'-diamidino-2-phenylindole) (Vector Laboratories).

Ventricle size. Brains fixed in 4% paraformaldehyde were embedded in agarose and cut coronally into 500  $\mu\text{m}$  (P15;  $n = 22$ ) or 375  $\mu\text{m}$  (P0;  $n = 24$ ) sections on a vibratome (Leica VT 1000S; Leica, Nussloch, Germany). Sequential sections were viewed under a light microscope (Zeiss, Oberkochen, Germany), and images were collected using MetaMorph imaging software. The section at the level of striatum and anterior commissure was selected from each animal, and the size (maximum width) of the lateral ventricles was measured using the MetaMorph software and divided by the maximum brain width. The resulting ratio of ventricle to brain diameter provides a measurement of hydrocephalus severity (Jones et al., 2001a). Hydrocephalic phenotypes were classified as mild (0.20–0.40), moderate (0.41–0.60), or severe ( $>0.60$ ).

RNA purification. Ependymal cell dissections (as described above) were done in RNAlater (Ambion, Foster City, CA) using RNase-free instruments. Total RNA was purified using the RNeasy Lipid Mini Kit (Qiagen, Germantown, MD) according to the manufacturer's protocol. RNA was eluted in a final volume of 60  $\mu\text{L}$  RNase-free water containing SUPERase-In™ (1 U/ $\mu\text{L}$ , Ambion, Foster City, CA) and concentrated using a Savant SpeedVac Concentrator (Thermo Scientific, Waltham, MA). RNA concentration was determined using NanoDrop™ 1000 spectrophotometer (Thermo Scientific, Waltham, MA) and samples were sent to the University of North Carolina Genomic and Bioinformatics core

facility for RNA quality analysis. Samples with a quality score less than 7.0 were discarded. Samples were stored at -20° C if not used immediately.

Gene microarrays. Affymetrix Mouse GeneChip Gene 1.0ST microarrays were used to detect changes in gene expression between Ro1 double transgenic mice and single transgenic littermate controls at 5 (n=3) and 9 (n=4) days after dox removal. 0.5 to 1.0 ug of total RNA from each animal was sent to the University of North Carolina Functional Genomics core facility. Each sample was amplified to obtain 7 µg of total RNA that was used to synthesize cDNA. A custom cDNA kit from Life Technologies was used with a T7-(dT)<sub>24</sub> primer for this reaction. Biotinylated cRNA was then generated from the cDNA reaction using the BioArray High Yield RNA Transcript Kit. The cRNA was then fragmented in fragmentation buffer (5X fragmentation buffer: 200mM Tris-acetate, pH8.1, 500mM KOAc, 150mM MgOAc) at 94°C for 35 minutes before the chip hybridization. 15 µg of fragmented cRNA was then added to a hybridization cocktail (0.05 µg/µl fragmented cRNA, 50 pM control oligonucleotide B2, *BioB*, *BioC*, *BioD*, and *cre* hybridization controls, 0.1 mg/ml herring sperm DNA, 0.5 mg/ml acetylated BSA, 100mM MES, 1M [Na<sup>+</sup>], 20mM EDTA, 0.01% Tween 20). 10 µg of cRNA was used for hybridization. Arrays were hybridized for 16 hours at 45°C in the GeneChip Hybridization Oven 640. The arrays were washed and stained with R-phycoerythrin streptavidin in the GeneChip Fluidics Station 400. After this, the arrays were scanned with the Hewlett Packard GeneArray Scanner. Affymetrix GeneChip Microarray Suite 5.0 software was used for washing, scanning, and basic analysis. Sample quality was assessed by examination of 3' to 5' intensity ratios of certain genes.

Microarray data analysis. CEL and CHP files were sent to Expression Analysis, Inc. (Durham, NC) for two-group comparison with permutation analysis for differential expression (PADE). As no genes were found to be differentially expressed with a false discovery rate (FDR) less than 1, genes with a fold change  $\geq .25$  and a p-value  $\leq 0.5$  (as determined by student's T-test assuming equal variance) were selected for pathway analysis. The DAVID bioinformatics database (<http://david.abcc.ncifcrf.gov/home.jsp>, National Institute of Allergy and Infectious Diseases, NIH) was used for gene-annotation enrichment analysis, KEGG pathway and BioCarta pathway mapping. Gene lists and expression data were sent to GeneGo Bioinformatics Software, Inc. (St. Joseph, MI) for detailed pathway analysis using the MetaCore platform.

Real time PCR. RNA (2.5-3.0  $\mu\text{g}$  total RNA) was converted to cDNA using random primers (250 ng/ $\mu\text{L}$ ) and SuperScript™ II reverse transcriptase (Invitrogen, Carlsbad, CA) in 20  $\mu\text{L}$  total volume following the manufacturer's protocol. PCR was carried out using 1  $\mu\text{L}$  cDNA for genes of interest or 1  $\mu\text{L}$  of a 1:10 cDNA dilution for the GAPD reference gene and 20X TaqMan® gene expression assay primer/probe sets (Applied Biosystems, Foster City, CA) in a 20  $\mu\text{L}$  final volume. Reactions were run in triplicate on a 7500 Fast Real-Time PCR System (Applied Biosystems, Foster City, CA) using the default PCR thermal cycling conditions. The reporter dye signal (FAM) was normalized using the passive reference dye, ROX, to eliminate volume-handling error. Fold change was determined using the comparative  $C_T$  method ( $\Delta\Delta C_T$ ).

Astrocyte cell culture. Litters were collected 0-2 days postnatal and lab protocol was followed to obtain sections of cortex from the pups. DNA preparations from the tails were screened for the presence of the tTA and lacZ genes by PCR. The cortex cell suspensions from individual pups were plated in 12 well plates, one pup per well. After seven days RASSL/tTA positive cultures were combined and cultures of wild-type or tTA only cells were combined. The cells were then plated either in 24 well plates for cAMP assays or on polylysine-coated 15 mm circle coverslips.


cAMP assay. 24-well plate cultures were incubated in 0.8 uCi  $^3\text{H}$ -Adenine per well for two hours. 0.2 mM IBMX was added to block the breakdown of cAMP. Cultures were then incubated in plain media, isoproterenol (1 uM), clonidine (10 uM), spiradoline (1 uM), isoproterenol and clonidine, or isoproterenol and spiradoline for fifteen minutes. Cells were lysed in 5% cold TCA. ATP and cAMP were eluted using Dowex and Alumina columns. Radioactivity of both elutions was measured in a scintillation counter and the data normalized for cell population.

## CHAPTER 4

### CHARACTERIZATION OF THE RO1 HYDROCEPHALUS MODEL


Glial cells, the majority of which are astrocytes, carry out numerous functions critical for the central nervous system to develop and operate normally. Certain of these functions are passive and supportive in nature. Astrocytes envelope neurons and contact blood vessels, providing metabolic and nutritive support. Astrocytes prevent neuronal excitotoxicity by buffering extracellular potassium and removing excess neurotransmitters from synapses (Gabriel et al., 1998; Walz, 2000; D'Ambrosio et al., 2002).

During development glia are critical for normal migration of neuronal precursors. Recent studies suggest astrocytes can actively modulate or initiate neuronal signaling (Walz, 1989; Keyser and Pellmar, 1994; Vernadakis, 1996; Carmignoto et al., 1997; Pasti et al., 1997; Araque et al., 1999; Araque et al., 2000; Araque et al., 2001; Pasti et al., 2001; Araque et al., 2002; Fiacco and McCarthy,



2004). If astrocytes actively regulate neurophysiology, alterations in astrocyte-neuronal signaling could contribute to the etiology of many CNS disorders. However, isolating astrocytic signaling systems to test this hypothesis is difficult since astrocytes and neurons express a similar complement of neuroligand receptors (Porter and McCarthy, 1996). Consequently, neuronal responses to astrocyte signaling under normal or pathological conditions have been difficult to measure directly.

We generated a mouse line expressing the  $G_i$ -coupled Ro1 RASSL (Receptor




Activated Solely by Synthetic Ligand) in glial fibrillary acidic protein (GFAP)-positive cells to directly assess the role of astrocytic G-protein coupled receptors (GPCRs) in neurophysiology. Ro1 is a kappa opioid receptor (KOR) modified by replacing its second extracellular loop with the delta opioid receptor second extracellular loop (Figure 5) and adding the FLAG® epitope tag to the N terminus. Conklin and colleagues demonstrated that Ro1 has a 200-fold decrease in binding affinity for dynorphin, the endogenous ligand for



KOR. Ro1 also has significantly reduced signaling in response to multiple dynorphin and enkephalin peptides while retaining its ability to bind spiradoline and *norbinaltorphimine*, a small molecule agonist and inverse agonist, respectively (Coward et al., 1998; Redfern et al., 1999). This is possible since the endogenous peptides bind to the KOR second extracellular loop while small molecule drugs bind in transmembrane regions of the receptor (Schwartz, 1994). Expressing Ro1 in glia on a KOR knockout background allows us to activate Gi-coupled receptor signaling in glia while measuring changes in neuronal excitability.

Restricting Ro1 expression to GFAP-positive cells required crossing two lines of transgenic mice, the Ro1 line and the tet-transactivator (tTA) line. The Ro1 line carries the Ro1 gene under the control of the




tetO promoter; the tTA line carries the tTA gene under the control of a 2.0 kB fragment of the human glial fibrillary acidic protein (hGFAP) promoter. In order for the Ro1 receptor to be expressed, tTA must bind the tetO promoter and initiate transcription. Since tTA expression is restricted to GFAP-positive cells, double transgenic progeny will only express Ro1 in glia. The timing and level of Ro1 expression in double transgenic mice can be manipulated with doxycycline. Doxycycline (dox) prevents Ro1 expression by binding to

tTA, thus preventing tTA from binding to the tetO promoter (Figure 6). Dox is able to cross the blood brain barrier, so Ro1 expression is turned off by maintaining transgenic mice on dox water and turned on by removing dox from the drinking water.

Crossing Ro1 and tTA transgenic lines produced double transgenic (tg) mice in expected numbers. However, the double tg mice unexpectedly developed severe hydrocephalus when maintained off dox (Figure 7). This is the first indication that activating a Gi-coupled GPCR in GFAP-positive cells can lead to hydrocephalus. Gi-linked signaling is involved in various physiological processes including proliferation, chemotaxis, hormone secretion, modulation of neurotransmission and contraction of cardiomyocytes (Pace et al., 1991; Spiegel, 1992b, a; Wickman and Clapham, 1995; Jan and Jan, 1997). The classical Gi signaling pathway is the inhibition of adenylate cyclase resulting in a decrease of cAMP


production, and it is now recognized that Gi signals via the ERK/MAPK pathway as well. Although how altering Gi-linked signaling cascades in glia results in HC is unclear, the prevalence of GPCRs and their importance in the CNS makes it unsurprising that changes in GPCR cascades causes pathology.



## Results


**Expression of Ro1.** Mice expressing Ro1 in GFAP-positive cells were generated by crossing the hGFAP-tTA mouse line to the tetO-Ro1/tetO-LacZ mouse line. This genotype was moved onto a KOR knock-out background (Hough et al., 2000) to prevent spiradoline from activating endogenous kappa opioid receptors. KOR<sup>-/-</sup>/hGFAP-tTA<sup>+/+</sup> or KOR<sup>-/-</sup>/hGFAP-tTA<sup>+/+</sup> mice were crossed to KOR<sup>-/-</sup>/tetO-Ro1/tetO-LacZ<sup>+/+</sup> mice to produce double tg (hGFAP-tTA<sup>+/+</sup>/tetORo1/tetO-LacZ<sup>+/+</sup>) mice, single tg (hGFAP-tTA<sup>+/+</sup> or tetO-Ro1/tetO-LacZ<sup>+/+</sup>) mice, and mice with neither transgene, all on the KOR background. Double tg mice were produced in expected numbers and are indistinguishable from littermates at birth. To determine that Ro1 expression is restricted to glia, brain sections from double tg mice were probed with anti-FLAG (the epitope tag fused to Ro1) antibodies plus either anti-GFAP (Glial Fibrillary Acidic Protein, an astrocytic marker) or anti-NeuN (neuronal nuclei, a neuron-specific nuclear protein) antibodies. Positive FLAG staining is apparent in most brain regions, including cortex, hippocampus, cerebellum and thalamus. FLAG staining is seen in GFAP-positive cells but not in NeuNpositive cells (Figure 8A and 8B), confirming that Ro1 expression is restricted to glial cells.

Since Ro1 expression is driven by the tetO system, dox can be used to manipulate when Ro1 is expressed. To verify that dox represses Ro1



expression, brain tissue extracts from double tg animals (P60) that had been maintained off dox, double tg animals that had been maintained on 25  $\mu\text{g/mL}$  dox and C57Bl/6J wild-type control animals were immunoprecipitated with a monoclonal Flag antibody. A band at approximately 44 kDa, consistent with the predicted size of the receptor, along with bands representing multimers of the receptor, were detected in extracts from double tg animals maintained off dox. No FLAG-specific bands were detected in wild type mice or double tg mice on 25  $\mu\text{g/mL}$  dox, indicating that 25  $\mu\text{g/mL}$  dox represses Ro1 expression (Figure 9). Since Ro1 tg mice also carry a tetO-LacZ transgene, we used Xgal histochemistry to confirm that dox regulates expression (Figure 10). No Xgal staining was detected in any brain region of sections from double tg mice on 25  $\mu\text{g/mL}$  dox. Strong Xgal staining was observed in the hippocampus, thalamus and brainstem of double tg mice maintained off dox while weak Xgal staining was seen in the cortex.

Characterization of hydrocephalus. Although double tg mice *off dox* appear grossly normal at birth, by postnatal day 15 (P15) 100% of double tg mice off dox begin to exhibit macrocephaly characterized by a swollen, dome-shaped cranium (Figure 11B). By 12 weeks of age all double transgenic mice off dox develop severe hydrocephalus and approximately 50% die (Figure 11C). Survival studies were terminated at 12 weeks due to the severe morbidity of the remaining double tg animals. Dissection of the brain revealed greatly enlarged ventricles filled with fluid, indicative of hydrocephalus and impaired CSF homeostasis. Blood on the surface of the brain was frequently observed in hydrocephalic mice and on occasion excess CSF was seen between the skull and brain surface. Control single tg littermates (hGFAPtTA<sup>+/+</sup> only or tetO-Ro1/tetO-lacZ<sup>+/+</sup> only) on or off dox and



double tg mice on 25 or 50  $\mu\text{g}/\text{mL}$  dox appear normal at P15 and do not develop macrocephaly at any age, indicating that the insertion of either or both transgenes does not cause hydrocephalus.

To further characterize the progression of hydrocephalus in double tg mice, the ventricle size of litters *off dox* were examined at P0 and P15 (Figure 12). At P0 double tg mice

( $n=10$ ) and control littermates ( $n=14$ ) showed no enlargement of the lateral ventricles. At P15, however, all double tg mice ( $n=12$ ) exhibited enlarged ventricles with a ventricle to brain ratio greater than 0.30 while control littermates ( $n=10$ ) had an average ratio of 0.10.

We observed

variation in the


severity of the


hydrocephalus

phenotype (Table 1)

as defined in

Materials and






**Methods.** The severity of hydrocephalus is not linked to homozygosity as all animals measured were heterozygous for both transgenes. The severity of hydrocephalus continued to progress with age since all double tg animals off dox developed severe hydrocephalus by P30 (Figure 12G and author's observations).

Brains of four control mice (single transgenic, P30 to P125) and five

**Table 1. Severity of hydrocephalic phenotype in P15 Ro1 mice**

<i>Phenotype</i>	<i>Average ratio</i>	<i>n</i>
<i>Mild</i>	0.31	4
<i>Moderate</i>	0.50	6
<i>Severe</i>	0.67	2

hydrocephalic double tg (P30 to P68) mice off dox were serially sectioned and stained with hematoxylin and eosin for a more detailed histological analysis. The most striking feature of the hydrocephalic brains was the greatly enlarged lateral ventricles (Figure 13G). The third ventricle, optic recess, pineal recess and intraventricular foramen were also enlarged. The



**Figure 13. Alterations in the ventricular system of hydrocephalic Ro1 mice.** Hematoxylin and eosin staining of coronal sections from single-transgenic control (P45, left column) and Ro1 (P33, right column) mice. **A,B**, In Ro1 mice (**B**), the third ventricle at the level of the periventricular hypothalamic nucleus is reduced in size and the ependymal cell layer appears thinner compared with controls (**A**). **D,F,H,J**, The subcommissural organ is disorganized in Ro1 mice (**D**), and there is a partial denudation of ependymal cells lining the ventricular walls (**F**, arrowhead; **J**). The remaining ependymal cells have fewer cilia (**F**, arrow), and the aqueduct is obliterated with no apparent ependymal cell layer (**H**). **C,E,G,I**, In control mice, the subcommissural organ (**C**) and aqueduct (**G**) show normal morphology. The ependymal layer is intact with multiple cilia per cell (**E**, arrow; **I**). **E** and **F** are enlargements of the boxes in **C** and **D**, respectively. Mice were maintained off dox. Aq=aqueduct, SCO=subcommissural organ, III=third ventricle, PC=posterior commissure. Scale bars: **A,B,G,H,I,J**, 50  $\mu$ m; **C,D**, 100  $\mu$ m; **E,F**, 30  $\mu$ m.


cerebral cortex was reduced in thickness, with the most severely affected animals exhibiting an almost completely atrophied cortex. All cortical layers appeared to be present (data not shown). The septum and retrosplenial cortex-hippocampus transition were disrupted; other brain regions, including the hippocampus, appeared compressed and caudally displaced. Red blood cells indicative of recent hemorrhaging were found throughout the brain (data not shown).

The white tract tissue underlying the lateral ventricles was disrupted and there was an incomplete denudation of ependymal cells lining the lateral (data not shown) and third ventricles (Figure 13J). In some instances the ependymal layer appears to be separated from the underlying tissue (Figure 13F). While it is possible that the separation is due to a fixation artifact, ependymal layer

separation was not observed in control mice.

Where intact, the ependymal cell layer appeared thinner (Figure 13*B,D,F*), the ependymal cells appeared flattened and had lost their cuboidal shape, and not all ependymal cells were ciliated. The rostral end of the aqueduct of Sylvius was completely obliterated (Figure 13*H*), suggesting that hydrocephalus results in part from the blocked flow of cerebral spinal fluid between the third and fourth ventricles. No ependymal cells were detected at the site of obstruction and the surrounding tissues appeared vacuolated. Abnormalities in the subcommissural organ (SCO) have been implicated in other models of hydrocephalus (Jones and Bucknall, 1988; Perez-Figares et al., 2001; Blackshear et al., 2003; Fernandez-Llebrez et al., 2004; Krebs et al., 2004); likewise in double tg brains the SCO appeared disorganized (Fig. 13*D*).

Adult onset of hydrocephalus. To determine if hydrocephalus results from Ro1 expression during development, mice were kept on 25 µg/mL dox until weaning (P21) and maintained




off dox until P60. Due to hardening of the skull, none of the double tg mice taken off dox developed overt enlargement of the cranium by P60; however double tg mice ( $n=4$ ) had enlarged lateral ventricles with an average ventricle to brain ratio of 0.30. By comparison, the ventricle to brain ratio for littermate controls ( $n=2$ ) was 0.11, similar to that of double tg mice ( $n=8$ , ratio=0.15) and controls ( $n=7$ , ratio=0.13)



that were maintained on dox continuously until P60 (Figure 14). Hydrocephalus appears to develop with Ro1 expression independent of age and thus is not dependent on processes linked to early stages of development.


Inhibition of Ro1 signaling. Since overexpression of a transgene may cause abnormalities due to disruption of normal cell transcription and translation machinery, signaling through Ro1 was suppressed with *nor*Binaltorphimine (*nor*BNI), a long-acting KOR inverse agonist (Jones and Holtzman, 1992; Ko et al., 1999; Holtzman, 2000). Double tg mice and single tg littermate controls were divided into three treatment groups (Figure 15). In the first group, double tg (n=2) and controls (n=2) were kept on 50 µg/mL dox. In the second group, double tg (n=4) and controls (n=4) were taken off 50 µg/mL dox at P30. In the third group, double

tg (n=6) and controls (n=5) were taken off 50 µg/mL dox at P30 and given injections of *nor*BNI (20 mg/kg, s.c.) every three days. All groups were sacrificed at P60 and ventricle to brain ratio



measured. As expected, no significant difference was found between double tg ( $0.16 \pm 0.08$ ) and littermate single tg ( $0.15 \pm 0.02$ ) controls on dox, while double tg mice taken off dox had a significantly larger ratio ( $0.29 \pm 0.07$ ) than control mice ( $0.08 \pm 0.05$ ). There was no difference in the ratios of double ( $0.17 \pm 0.06$ ) and single ( $0.13 \pm 0.09$ ) tg mice taken off dox and given *norBNI*. Since expression of the receptor alone did not result in enlarged ventricles, signaling through Ro1 appears to be required for HC to develop.

Blocking Ro1 signaling also prevented endymal denudation in double tg




mice. The same treatment groups listed above were sacrificed 10 days after dox removal and their brains processed for H and E staining. Only the double tg mice taken off dox showed loss of endymal cells

in the lateral ventricles, while *norBNI* treated double tg mice and double tg mice on dox had intact endymal cell layers similar to controls (Figure 16).

GFAP and phosphoErk. Increased GFAP expression, an indicator of reactive glia, is observed in most forms of damage to the brain, including hydrocephalus. Increased levels of GFAP staining were seen in hydrocephalic Ro1 mice. The cortex, a region typically low in

GFAP immunoreactivity, and the hippocampus showed the greatest increases in GFAP staining compared to control mice (Figure 17). Western blots confirmed that GFAP protein levels were elevated in the cortex and hippocampus of Ro1 mice; GFAP levels in cerebellar tissue extracts from the same mice appeared similar to controls (Figure 17A).

Because Gi-coupled receptors are known to activate the mitogen-activated protein (MAP) kinase pathway, coronal sections from Ro1 mice were stained for phosphorylated




extracellular signalregulated kinase (Erk) p44/42, the active form of Erk p44/42. Increased phosphoErk staining was observed in the superficial layers of entorhinal cortex and in the striatum surrounding the lateral ventricles of Ro1 animals off dox, but not in littermate single transgenic controls (Figure 18).

## Discussion

To investigate how Gi protein signaling in astrocytes affects brain neurophysiology, we expressed the RASSL Ro1, a Gi-coupled GPCR, in GFAP-positive using the hGFAP promoter and the tet-off system. Unexpectedly, all mice expressing Ro1 developed non-

communicating triventricular hydrocephalus. Hydrocephalus appears to result from Ro1 expression since single transgenic mice do not develop hydrocephalus as would be expected if it was caused by an insertional defect. Hydrocephalus is prevented by suppressing Ro1 expression with dox and by blocking Ro1 signaling with *norBIN*. Ro1 mice taken off dox at weaning (P21) develop enlarged ventricles, further supporting the idea that hydrocephalus is a consequence of Ro1 expression.

At birth double tg mice off dox appeared grossly normal and were indistinguishable



**Figure 18. Phospho-Erk p44/42 levels are elevated in hydrocephalic Ro1 mice.** Increased phospho-Erk p44/42 staining is observed around the lateral ventricles (**B**) and in the entorhinal cortex (**D**) of hydrocephalic Ro1 mice but not in littermate single-transgenic controls (P21) (**A,C**). No increases in phospho-Erk were observed in the hippocampus (**F**). Sections shown in **A** and **B** are at the level of the anterior commissure and striatum; **C-F** are at the level of the subcommissural organ. Scale bars, 150  $\mu$ m. LV=lateral ventricle, DG=dentate gyrus.

from their control littermates, but began to exhibit macrocephaly between P10 and P15. At P15 Ro1 mice have a lateral ventricle to brain ratio that exceeds 0.30 and by P30 have developed a severe hydrocephalus phenotype characterized by greatly dilated lateral and third ventricles that caudally displace surrounding brain structures. The rostral end of the Sylvius aqueduct is obliterated, with no lumen and no detectable ependymal cells visible in coronal sections of this

region, suggesting a denudation of ependymal cells has occurred. The collicular recess and caudal end of the Sylvius aqueduct, however, remain open. The subcommissural organ (SCO) also appears disorganized in hydrocephalic mice, although it is unclear whether this is due to Ro1 expression or if it is a result of hydrocephalus.

It is unclear how, in the absence of applied ligand, Ro1 expression leads to the development of hydrocephalus. Ro1 is likely to be highly over-expressed relative to other astrocytic receptors. Since G-protein-coupled receptors have a low level of constitutive activity (Lefkowitz et al., 1993; Costa and Cotecchia, 2005), Ro1 may be expressed at a high enough level that its constitutive activity is sufficient to trigger a series of events that ultimately result in hydrocephalus. Ro1 may be also expressed at a high enough level that dynorphin or another ligand that normally would have a low affinity for Ro1 can bind and activate the receptor. Significantly, mice with Ro1 expression targeted to the heart developed cardiomyopathy when taken off dox. When these mice were given an intermediate dose of dox to reduce Ro1 expression, their heart rates normalized. Treating these mice with pertussis toxin or the KOR inverse agonist *norbin* also restored normal heart rates, demonstrating that Ro1 signaling was responsible for the changes in heart (Redfern et al., 2000). A similar process where Ro1 overexpression and constitutive activity leads to pathology may be occurring in our mice.

Our finding is, to our knowledge, the first to implicate glial G<sub>i</sub> signaling pathways in the development of hydrocephalus. Ro1 expression in astrocytes may contribute to the development of hydrocephalus by altering the extracellular matrix so that more fluid from the extracellular space enters the ventricles. It could also alter the levels of various neurotransmitters, growth factors, or cytokines in the CSF resulting in a dysregulation of CSF

production by the choroid plexus. Astrocytic release of glutamine is known to be down regulated in the kaolin model of hydrocephalus (Kondziella et al., 2003); it is possible that increased G<sub>i</sub> signaling in astrocytes may affect glutamine production or release that in turn changes neuronal input into the SCO. It is interesting to note that gene expression analysis in the H-Tx rat, which has primary stenosis of the aqueduct similar to Ro1 mice, has implicated several genes linked to astrocytes and G-protein coupled receptors, including connexin 30, beta integrin 5, and somatostatin (Jones et al., 2001b; Miller et al., 2006).

Although SCO secretory function could not be determined from our histological examination of Ro1 brains, the observation that the SCO is disorganized in Ro1 mice is significant. The presence of a normal SCO appears to be necessary for the development and maintenance of the aqueduct, as impaired SCO formation and/or function are found in multiple animal hydrocephalus models and in human infantile hydrocephalus (Overholser et al., 1954; Newberne, 1962; Takeuchi and Takeuchi, 1986; Jones et al., 1987; Jones and Bucknall, 1988; Perez-Figares et al., 1998; Louvi and Wassef, 2000; Estivill-Torrus et al., 2001; Sakakibara et al., 2002; Fernandez-Llebrez et al., 2004). The SCO secretes negatively charged glycoproteins, such as SCO-spondin and RF-Gly I, that appear to be critical for maintaining an open aqueduct by their physical presence. The SCO may also regulate CSF formation. Receptors for SCO glycoproteins are found on the choroid plexus (Miranda et al., 2001), suggesting that CSF production could be influenced by SCO activity.

We have shown that phosphoErk levels are increased in hydrocephalic mice. G<sub>i</sub>-coupled receptors are known to signal via the MAP kinase pathway, thus Ro1 signaling may lead to changes in gene expression that contribute to the pathogenesis of hydrocephalus. We cannot rule out, however, the likely possibility that the observed increase in phosphoErk is a

consequence of hydrocephalus and not causal. Elevated phosphoErk has been associated with astrogliosis (Mandell and VandenBerg, 1999), a pathological response of astrocytes to many types of brain injury, including hydrocephalus (Fukumizu et al., 1996). Increased GFAP expression observed in severely hydrocephalic Ro1 mice is indicative of astrogliosis, particularly since GFAP levels are not elevated in Ro1 mice with mild hydrocephalus (data not shown).

The observed loss of ependymal cells in Ro1 mice is in agreement with other studies that demonstrate a denudation of ependymal cells in the ventricular system (Jimenez et al., 2001; Dominguez-Pinos et al., 2005). The underlying cause of ependymal detachment is unknown, although defects in certain adhesion molecules, such as L1, have been indicated in other HC models (Schmid et al., 2000; Itoh et al., 2004). As a subset of ependymal cells express GFAP (Takano et al., 1996; Rodriguez-Perez et al., 2003), it is possible that Ro1 may be expressed in these cells, causing a defect that leads to detachment. Ependymal cells may also be affected indirectly by changes in levels of secreted cytokines or altered neuronal inputs resulting from Ro1-induced changes in astrocyte signaling.


To develop more effective treatments, a better understanding of hydrocephalus pathogenesis is needed. The Ro1 model of hydrocephalus has several characteristics that will make it valuable for studying hydrocephalus. First, the Ro1 model does not require injection of foreign bodies into the ventricles to cause hydrocephalus, making the Ro1 model easier to work with and more relevant to human cases. Second, hydrocephalic animals are produced reliably; all Ro1 mice will develop hydrocephalus if maintained off dox. Inherited models of hydrocephalus, such as the H-Tx rat, reliably produce hydrocephalic offspring but at a much lower rate, and which animals will develop hydrocephalus cannot be predicted at early ages

(Jones et al., 2000; Jones et al., 2001b; Miller et al., 2006). Third, dox can be used to regulate when hydrocephalus will occur. It would be interesting to compare juvenile and adult onset to see if age changes how hydrocephalus develops. Fourth, the rate of development and the severity of hydrocephalus can be manipulated by giving Ro1 mice a lower concentration of dox. Slowing the development of hydrocephalus allows changes in the brain to be studied more thoroughly and could provide insights into how hydrocephalus progresses. The Ro1 model of hydrocephalus should prove useful in advancing our knowledge of hydrocephalus pathophysiology. Furthermore, the Ro1 model provides new evidence for the involvement of astrocytic G-protein coupled receptor signaling in the development of hydrocephalus.




## CHAPTER 5 EPENDYMAL CELL GENOMIC ARRAYS

Although the cause of HC is multi-factorial, the progression of pathology appears remarkably similar across various models of the disease. A common signaling pathway(s) may underlie the pathology. If so, identification of the pathway(s) involved would allow new



pharmacological approaches for treating HC to be developed. Because onset of the disease in our model is controlled by dox, the Ro1 model provides the opportunity to look for the earliest changes in disease progression without the complications caused by later pathology.

Ependymal denudation is a common feature of HC that appears to happen early in the progression of the disease. Ependymal denudation is reported in most, if not all, HC models and human cases. Whether denudation occurs prior to ventricular dilation or as a result of dilation is still unsettled, however there is strong evidence that denudation occurs early in the disease process, and loss of the ependyma cell layer or cilia function is also sufficient to cause HC (Davy and Robinson 2003, Ibanez-Tallon et al 2002, Grondona et al 1996). In the



Ro1 HC model, expression of the  $G_i$ -coupled Ro1 receptor in GFAP positive cells results in HC, although how enhanced  $G_i$  signaling leads to HC is unknown. Ependymal denudation is also seen in the Ro1 model, prompting us to question whether Ro1 signaling cascades could lead to ependymal denudation and subsequent onset of HC. To address this question, we used Affymetrix gene arrays to look for changes in gene expression in ependyma shortly after turning on Ro1 expression.


## Results

Expression of Ro1 in ependymal cells. Because of conflicting reports of GFAP expression in ependymal cells, we wanted to determine if ependymal cells express GFAP in our mouse line. Brains from adult mice were processed for immunostaining and cut into coronal sections. Sections containing lateral ventricles, third ventricle and aqueduct were probed with antibodies against GFAP and the ependymal cell marker S100β. A subpopulation of the ependymal cells was double-labeled with both markers, and the specialized ependyma of the SCO strongly expressed GFAP (Figure 19B). Additionally, in the lateral and third ventricles GFAP-positive processes could be seen wrapping around cell bodies in the subependymal cell layer and extending to the ventricular surface (Figure 19C). To further define the GFAP


labeled cells, sections were stained for GFAP and  $\alpha$ -tubulin, which is highly expressed in cilia. The cells lining the ventricles and the cells of the choroid plexus were edged on the luminal side with positive  $\alpha$ -tubulin staining. A subpopulation of the ependymal cells lining the lateral and third ventricles and the majority of ependyma lining the aqueduct were GFAP positive and labeled with anti-  $\alpha$ -tubulin antibodies on the apical surface, suggesting that the GFAP expressing cells are ciliated ependymal cells. Cells of the choroid plexus did not express GFAP in any of the sections (Figure 19).

Secondary antibodies can sometimes bind nonspecifically to the edges of tissue, resulting in falsely labeled cells. To reduce the likelihood that edge effect could be influencing staining in sections, the ependymal cell layer was isolated in sheets from the lateral ventricles and placed on slides for staining. Most cells in the isolated sheets expressed S100 $\beta$ , confirming that they were ependymal cells, and the majority of S100 $\beta$  positive cells were also labeled with GFAP (Figure 20).  $\alpha$ -tubulin staining of ependymal sheets revealed distinct tufts of cilia on the cells, and ciliated cells also expressed GFAP (Figure 21). Thus in the Ro1 mouse model at least a subpopulation of ependymal cells do express GFAP and would be able to express the Ro1 receptor.


To look for the Ro1 receptor expression in ependymal cells, double tg Ro1 mice and single tg littermate controls were taken off dox at P30 and



sacrificed two to four weeks after dox removal. The Ro1 receptor is FLAG-tagged, so fresh



frozen coronal sections and isolated ependymal cell sheets were probed with anti-FLAG epitope and GFAP antibodies. A small percentage of ependymal cells expressing GFAP stained positively for FLAG in both brain sections (Figure 22) and ependymal sheets from double tg mice but not in any tissue from control mice, confirming that Ro1 is indeed




expressed in ependymal cells from double tg mice.

Timing of Ro1 expression and ependymal denudation. In order to look for early events in HC development, the timing of Ro1 expression following dox removal needed to be determined. Double tg Ro1 mice and litter mate controls were taken off dox at P30 and

brains collected at day 0, day 3, and day 7 after dox removal. Fresh frozen coronal brain sections were probed for the FLAG epitope (Figure 23). On day 0, no FLAG staining was observed in either double or single tg mice. By day 3, positive FLAG staining was visible mainly in the thalamus of double tg mice, and by day 7 positive FLAG staining was found in the thalamus, hippocampus, and cortex. At no timepoint did FLAG antibodies positively label cells from single tg mice.

Onset of ependymal denudation after dox removal was determined by H&E staining of coronal brain sections. Double tg Ro1 mice and littermate controls were taken off dox at P30 and pairs were killed at day 3, day 6, day 10, and day 14. Ten days after dox removal small patches of ependyma lining the septal wall of the lateral ventricles were missing or detached from the underlying parenchyma in Ro1 mice. Most of the ependymal cells remained attached and retained a cuboidal shape, and there was no evidence for ependymal detachment in the aqueduct. Single tg mice did not have ependymal detachment at any timepoint. Thus ependymal detachment begins by 10 days after dox removal in double tg Ro1 mice.

Affymetrix gene arrays. Genomic arrays were utilized to identify target ependymal genes that may be responsible for subsequent cell detachment. To avoid gene changes that result as a consequence of HC, ependymal



**Figure 24. Time course of ependymal denudation after dox removal.** Coronal sections from control (A,B) and Ro1 mice (C,D) were stained with hematoxylin and eosin to look for ependymal denudation at 6 days (A,C) and 10 days (B,D) after dox removal. At six days off dox ependyma are normal in both control (A) and Ro1 mice (C). Patches of ependymal denudation are evident at 10 days off dox in Ro1 mice (D) but not in littermate controls (B).



cells needed to be collected very early in the disease process, but not so early that changes have not yet occurred. Based on the time course of Ro1 expression and ependymal denudation, day 5 and day 9 after dox removal were chosen as the timepoints to be used in gene arrays. Expression of Ro1 in double tg ependyma at these timepoints was confirmed using RT-PCR. Double tg (n=3) and single tg (n=3) mice were taken off dox at P30 and sacrificed at P35; four double tg and four single tg mice were taken off dox at P30 and sacrificed at P39. Approximately 0.5-1.0 µg of total RNA was extracted from the isolated ependymal cells of each animal and amplified for use in Affymetrix Mouse GeneChip Gene 1.0ST arrays. All chips passed internal quality controls, and the data was sent to Expression Analysis, Inc. for statistical analysis. Expression data for each genotype and timepoint was averaged together and two-group comparison with permutation analysis for differential expression (PADE) was used to identify differentially expressed genes. Using the criteria of  $p\text{-value} \leq 0.05$  and  $\text{fold change} \geq 0.20$ , 97 genes were differentially expressed at 5 days off dox and 343 genes were differentially expressed at 9 days off dox. Most gene changes were small. At 5 days off, the largest fold change was .67 and at 9 days off dox only 37 genes were changed by more than 20%. However, no genes were found to be differentially expressed with a false discovery rate (FDR) less than one, meaning that all changed genes are likely not truly different between double and single transgenics.

RT-PCR validation. Twelve genes from each timepoint were selected based on p-value and biological relevance for RT-PCR validation (Table 2). There was no significant difference between Ro1 and control mice for any of the selected genes, even when ependymal cells from multiple mice (same genotype) were combined to generate sufficient total RNA for

efficient reverse transcription. We were therefore unable to interpret the gene array data or draw any conclusions about the effect of Ro1 signaling in ependymal cells.

## **Discussion**

Similar to other HC models, ependymal denudation is a characteristic of HC in our Ro1 mouse model of HC. Because ependymal denudation is sufficient to cause HC (Grondona et al., 1996), we wanted to use gene arrays on ependymal cells to see if overexpression of Ro1 leads to changes in gene expression that ultimately causes denudation. Although some groups have not been able to detect GFAP in ependyma, other groups have shown that a subset of ependyma do express GFAP. The conflicting reports have been suggested to result from differences in fixation and staining methods, age, and type of animal (Bruni et al., 1985; Kasper et al., 1991; Sarnat, 1992; Xiuxin Liu, 2006). Immunostaining of adult ependyma in tissue sections and isolated ependymal sheets from our mice showed that a percentage of ciliated cells expressing the ependymal cell marker S100 $\beta$  also expressed GFAP. Thus in the Ro1 mice, we also detected GFAP in a subpopulation of ependymal cells similar to previous reports. More detailed studies would need to be carried out to determine the percentage, although GFAP expression seems to be higher in the SCO and aqueduct. Since ependymal cells do express GFAP, Ro1 should be expressed in the ependyma of double tg mice. Immunostaining for the FLAP epitope confirmed that at least a small subset of ependymal cells express the Ro1 receptor. This finding is not unexpected since not all ependyma detach during HC. If Ro1 expression does cause denudation in our model, then it is possible that GFAP could be a marker for the subsets of cells that detach in HC in other models.

Selecting timepoints for gene arrays was of some concern. The Ro1 model allows the disease process to be started by removing dox and turning on Ro1 expression, but the time required for dox to be cleared from the CNS and Ro1 expression to begin was unknown. If too early a timepoint was selected, Ro1 would not be expressed or would not have been expressed long enough for changes in gene expression to occur. If too late a timepoint was chosen, the early events of HC would be missed, and ependyma may have started to detach, making tissue collection extremely difficult. Timed studies of Ro1 expression and ependymal denudation after dox removal in double tg mice showed that Ro1 is expressed in the thalamus as early as three days after dox removal. Although isolated ependymal sheets were not included in the timed staining studies, subsequent RT-PCR of double tg ependyma

5 Days Off Dox	9 Days Off Dox
TNF $\alpha$	MID1
Caspase 8	MNS1
SIX2	CABLES2
SLC17A6	SCP2
TRHR	DSC3
LRP2	LRP2
MYL1	GFRA2
ITGAX	HYDIN
C1S	EPS8L1
ST8S1A6	ADAMTS3
PLEC1	TNFAIP6
	CSF3

**Table 2.** Gene targets for RT-PCR validation

at 5 and 9 days off dox showed that Ro1 can be detected at both timepoints and is more highly expressed at the 9 day timepoint, suggesting that dox is still being eliminated from the brain. H&E staining from similar timed studies showed normal ependyma in double tg mice at 3 and 6 days off dox. At ten days off dox, small areas of detaching ependyma were seen in the lateral ventricle. However, it should be noted that only

two mice were examined in this study, and disease progression is highly variable between animals. Based on the staining results, 5 and 9 days off dox were selected for the genomic array timepoints.



The gene array data cannot be interpreted since no differentially expressed genes between double tg mice and single tg controls at either timepoint could be validated by PADE analysis (for false discovery) or by RT-PCR. Although not many changes were expected to be found, the lack of differences is surprising and could be attributed to several different factors. First, Ro1 expression in ependyma may not cause changes at the gene level at these timepoints. Instead, Ro1 signaling may alter other signaling pathways that respond rapidly and result in denudation. Ro1 signaling in ependyma may not be the causal factor in detachment. Rather, Ro1 signaling in other cell types could lead to detachment. It is also possible that although Ro1 expression was detected in the brain at three days off dox, ependymal cells would be exposed to dox longer through contact with the CSF and thus not begin expressing Ro1 at high enough levels to cause pathology until later timepoints. Alternatively, the percentage of cells expressing Ro1 in the ependymal isolations may be small enough that changes in gene expression actually occurring could not be detected. The large variability in rate of disease progression between animals could also make actual changes in gene levels hard to detect statistically without using a much larger sample size. The lack of changes could also be due to technical issues. The amount of total RNA submitted in each sample was very small, less than 1  $\mu$ g. The necessary amplification steps could skew the relative levels of transcripts used in the array. Although data from the arrays cannot be interpreted, it is intriguing that a number of genes differentially expressed (by student's t test) were genes related to cilia function, extracellular matrix, and cell adhesion. However, these data cannot be interpreted, and further work is required to determine what changes, if any, Ro1 signaling causes in ependymal cells.

## **CHAPTER 6**

### **DISSCUSSION**

Despite decades of neuroscience research, many neurological disorders remain poorly understood. Interest in the role of glia in both normal and pathological states is increasing, particularly as evidence is presented for astrocyte modulation of neuronal activity. Recent research has provided evidence that astrocytes may have a greater function beyond providing structural and metabolic support for neurons. First, astrocytes account for approximately half of the brain's volume and are found in every part of the CNS. Many neuronal processes and synapses are surrounded by astrocytes; a single astrocyte can envelop as many as 80,000 synapses (Bushong et al., 2002). Second, astrocytes and neurons express a similar complement of receptors for neuroligands (Porter and McCarthy, 1996) and astrocytes are able to respond to numerous stimuli with an increase in intracellular calcium (Araque et al., 2001), which is generally accepted to be a form of glial excitability (Hansson and Ronnback, 2003). Third, astrocytes also appear to release gliotransmitters, including glutamate and ATP. For example, a calcium increase in astrocytes evokes an inward current in adjacent neurons that is dependent on glutamate release (Parpura et al., 1994; Carmignoto et al., 1997; Araque et al., 1998; Pasti et al., 2001) and uncaging IP<sub>3</sub> in a single astrocyte causes a glutamate-dependent increase in AMPA mEPSC frequency in a neighboring neuron (Fiacco and McCarthy, 2004). Yet little is actually known about what role astrocytes may play in listening to and modulating neuronal communication under physiological conditions.

Because an astrocyte-specific receptor and ligand have yet to be discovered, astrocytic responses cannot be isolated pharmacologically. The Ro1 transgenic mouse line was created to manipulate signaling events specifically in astrocytes in order to investigate the role of astrocyte GPCR signaling in modulating behavior and neuronal activity.

**Targeting a G<sub>i</sub>-coupled receptor to astrocytes.** The tetracycline transactivator (tTA) system was used to conditionally express Ro1 in astrocytes on a kappa opioid receptor knockout background. Ro1 is a kappa opioid receptor modified by replacing the second extracellular loop with the second extracellular loop of the delta opioid receptor. Ro1 has a greatly reduced affinity for endogenous ligands but is still activated by the highly selective small molecule drug spiradoline (Coward et al., 1998; Redfern et al., 1999). To restrict Ro1 expression to astrocytes, transgenic (tg) mice expressing the tetracycline transactivator (tTA) under the control of hGFAP, an astrocyte-specific tissue promoter, were crossed with a second line of mice carrying the Ro1 and LacZ genes under the control of the tet operon (tetO) promoter. Timing and level of Ro1 expression can be controlled with doxycycline, thus any potential developmental side effects of Ro1 expression could be eliminated by maintaining mice on doxycycline until maturity. Using spiradoline to activate Ro1 would allow the study of astrocytic G<sub>i</sub>-coupled signaling on animal behavior and neuronal activity.

**Ro1 expression and cellular localization.** Immunoprecipitation assays and  $\beta$ gal staining demonstrated that Ro1 is expressed only in double tg mice (positive for both tTA and Ro1) and only when the animals are maintained in the absence of dox, confirming that Ro1 expression is indeed regulated by doxycycline without any apparent “leakiness”. Cellular localization of Ro1 was determined using immunocytochemistry; the localization of Ro1

immunostaining to GFAP positive cells and its absence in NeuN positive cells confirmed that Ro1 expression is being driven by the GFAP promoter. Because spiradoline, the synthetic ligand for Ro1, activates the kappa opioid receptor, the Ro1 genotype was moved onto a KOR knockout background to ensure that the only response to spiradoline is mediated by Ro1.

**G<sub>i</sub> signaling and G<sub>i</sub> modulation of behavior.** G protein coupled receptors (GPCRs) are a class of seven transmembrane spanning receptors that are important mediators of intracellular signaling. The cytoplasmic tail of a GPCR interacts with the heterotrimeric G proteins, which are composed of  $\alpha$ ,  $\beta$  and  $\gamma$  subunits. In an inactive state, the  $\alpha$  subunit binds GDP. Activation of the GPCR causes a conformational change in G $\alpha$ , allowing GDP to be replaced with GTP and releasing the G $\beta\gamma$  dimer. There are multiple families of G $\alpha$  that couple to different effector molecules. G $\alpha_i$  is so named for its inhibition of adenylate cyclase and subsequent reduction of cAMP formation. A number of studies suggest that G<sub>i</sub>-coupled receptors in the CNS, including dopamine, serotonin, opioid and muscarinic acetylcholine receptors have roles in regulating both behavior and neuronal activity in normal and pathological states. The focus of many of these studies has been on neuronal receptors even though astrocytes also express many of the same receptors and there is indirect evidence that activation of astrocytic receptors may play a role in the observed effects. The Ro1 mice were designed to allow us to directly test if G<sub>i</sub> signaling in astrocytes alone can modulate behavior or neuronal activity.

Alterations in G<sub>i</sub> protein levels and activity have been implicated in multiple pathological states, including depression and panic disorders (Joseph et al., 1993; Saitoh et al., 1993; Gurguis et al., 1999c; Gurguis et al., 1999e; Gurguis et al., 1999d; Gurguis et al.,

1999b; Gurguis et al., 1999a). There is a significant reduction in stimulated cAMP formation in frontal cortex tissue from human suicide victims (Valdizan et al., 2003) and both increases (Garcia-Sevilla et al., 1999) and decreases (Pacheco et al., 1996) in G<sub>i</sub> protein levels have been found in the brains of depressed subjects. As these studies were done on brain tissue lysates, neuronal and astrocytic components cannot be determined. Dopamine D2 receptors signal primarily via G<sub>i</sub> pathways. Neuroleptic antipsychotic drugs, which are primarily dopamine D2 receptor antagonists, are used to treat depression and panic disorders. Since nearly 1/3 of all D2 receptors in the cerebral cortex are located on astrocytes and astrocytic processes rich in D2 receptors surround cortical interneurons lacking D2 receptors (Khan et al., 2001), astrocytes seem to be ideally placed to mediate some of the therapeutic effects. Furthermore, the D2 agonist quinpirole causes an increase in intracellular calcium in cultured cortical astrocytes that can be blocked with the D2 antagonist raclopride (Khan et al., 2001); increases in astrocytic calcium are correlated with changes in neuronal activity and release of gliotransmitters (Pasti et al., 1997; Newman and Zahs, 1998; Parri et al., 2001). It is possible that activity at astrocytic dopamine receptors accounts for at least part of the effect of antidepressant and antipsychotic drugs, and therefore defects in astrocytic G<sub>i</sub> signaling could have a critical role in these disorders.

**G<sub>i</sub> modulation of neuronal excitability.** G<sub>i</sub>-receptor signaling is also associated with changes in neuronal activity. For example, kappa opioid receptor activation *in situ* inhibits glutamate synaptic currents presynaptically (Svingos and Colago, 2002; Bie and Pan, 2003), reduces NMDA mediated glutamatergic transmission (Svingos and Colago, 2002), prevents glutamate release evoked by potassium (Nicol et al., 1996), and inhibits NMDA mediated neurotransmitter release (DA and Ach) (Schoffelfmeer et al., 1997). Dopamine receptor

agonists also suppress NMDA neurotransmission (Wang et al., 2006). While there is little question that neuronal receptors are involved in the observed results, astrocytes also express kappa opioid (Bunn et al., 1985; Eriksson et al., 1990; Belcheva et al., 2005) and dopamine receptors (Miyazaki et al., 2004), and activity at astrocytic  $G_i$  receptors could be involved in modulating the observed neuronal responses. As astrocytes express a number of different  $G_i$ - and  $G_s$ -coupled receptors it is likely that  $G_i$  signaling is important in regulating processes that effect astrocyte-neuron communication. For example, astrocytes respond to neuronal vasoactive intestinal peptide (VIP), a  $G_s$ -coupled receptor, with a release of neurotrophic factors, including activity-dependent neurotrophic factor (ADNF) (Gressens et al., 1997). ADNF causes the secretion of neurotrophin 3 in neurons and may function in long term potentiation by modulation of NMDA subunits 2A and 2B (Blondel et al., 2000). Since VIP receptors in astrocytes are  $G_s$ -linked GPCRs, activation of  $G_i$  could inhibit the release of ADNF and thus suppress long term potentiation induction.

**Mice expressing Ro1 develop hydrocephalus.** The original goal in creating the Ro1 mice was to allow selective activation of  $G_i$  signaling in a population of astrocytes while recording any subsequent changes in neuronal excitability or performing behavioral tests. However, severe hydrocephalus was observed in 100% of double tg mice off dox (expressing Ro1). All double tg mice maintained off dox developed visibly enlarged heads and had a 50% survival rate at 12 weeks of age, while all double tg mice on 25  $\mu$ g/mL dox were normal with a 100% survival rate at 12 weeks, demonstrating that suppression of Ro1 expression prevents hydrocephalus. Control mice heterozygous for the Ro1 gene but lacking tTA (GFAP-tTA<sup>-/-</sup> -

tetO:Ro1<sup>+/+</sup>) did not develop hydrocephalus, making it unlikely that the Ro1 gene is inserted in some critical location that results in hydrocephalus independent of receptor expression.

**Doxycycline dosing.** The appearance of hydrocephalus in double tg mice posed a serious problem for future studies of Ro1 signaling in astrocytes as differentiating between physiological or pathological responses would be impossible. The Conklin group had similar problems with G<sub>i</sub> associated pathology when they over-expressed Ro1 in heart tissue, but were able to eliminate pathology in the heart by administering an intermediate dose of doxycycline to their animals (Redfern et al., 2000). Extensive attempts were made to determine a minimum dose of doxycycline that would eliminate hydrocephalus while still allowing expression of Ro1 at a detectable level. Double tg mice maintained on 1, 2, 5, 10, and 12 µg/mL doxycycline developed noticeably enlarged heads and had enlarged lateral ventricles, although with increasing age of onset (unpublished observations). All double tg mice maintained on 17 µg/mL dox appeared outwardly normal but had enlarged lateral ventricles at p60, yet double tg mice on 25 µg/mL dox had normal ventricles at p60. The relatively small change in dox concentration between healthy and affected double tg mice, coupled with difficulty in detecting the Ro1 receptor in mice maintained on 17 µg/mL dox, prompted us to focus on the Ro1 mouse line as a HC model.

**The Ro1 model.** The unexpected development of HC in the Ro1 mouse line has resulted in a unique and valuable model of HC. The pathology of HC in the Ro1 model is very similar to other HC models with stenosis of the aqueduct. HC appears to develop postnatally in our model as newborn double tg pups (maintained off dox) are indistinguishable from control

littermates. At birth the ventricles are not enlarged, the ependymal cell layer appears intact and brain structures appear grossly normal. After birth HC develops progressively; by p15 double tg pups have domed skulls and significant enlargement of the lateral and third ventricles, optic chiasm and pinal recess. Similar to other HC models, discontinuous loss of ependymal cells is seen throughout the lateral ventricles, third ventricle and the aqueduct. Damage to periventricular white matter is prominent with thinning of the cortex and posterior displacement of the hippocampus. The cerebellum is smaller but appears otherwise normal, the SCO is small and disorganized, and the aqueduct is obliterated (Sweger et al., 2007). Many of these pathological changes are common in HC, both in experimental models and in human cases.

Perhaps it is not surprising that pathology resulting from accumulation of CSF would be similar across models. However, the initiating events in HC are still unclear, and the underlying mechanisms that cause neurological and motor deficits in patients are poorly understood. The similarities between models suggest that there may be a common pathway or process that causes HC when disrupted. Identification of such a process could provide the opportunity to develop sorely needed pharmacological treatments for HC. A number of other HC models already exist, but all have limitations that make isolating initiating events in HC difficult. In spontaneous models, onset of HC cannot be controlled and penetrance is not 100%. Thus the animals that will develop HC cannot be predicted, limiting studies to the pathology and not the causes of HC. In induced models, the onset of HC is controlled but does not reflect congenital HC and may be complicated by inflammatory responses. Other transgenic mice develop HC as part of their phenotype, but the presence of other pathologies or developmental abnormalities makes identifying the processes responsible for HC difficult.



In the Ro1 model HC is initiated by disruption of G<sub>i</sub> signaling, HC is the only pathology, onset of pathology is controlled, the rate of HC progression can be manipulated, and both congenital and adult onset HC can be studied. Thus the Ro1 model presents a new tool for understanding the molecular mechanisms of HC.

**Ro1 signaling.** Although the rate of HC development varies between animals, all double tg mice off dox will develop severe HC despite the absence of the Ro1 agonist spiradoline. GPCRs have basal levels of intrinsic signaling that are not blocked by an antagonist but can be blocked with an inverse agonist (Leurs et al., 2000; Kenakin, 2004; Wang et al., 2007). Because Ro1 is based on the kappa opioid receptor (KOR), a GPCR, basal signaling through Ro1 is the most likely explanation for development of HC in our model. *Nor*Binaltorphimine (*nor*BNI) is a KOR inverse agonist frequently used in opioid studies (Messer, 2004; Wang et al., 2007) and has been shown to be highly selective for KOR over other opioid receptors. A single dose can inhibit KOR signaling for weeks (Horan et al., 1992; Butelman et al., 1993; Powell and Holtzman, 1999; Ko et al., 2003; Messer, 2004), making *nor*BNI an ideal drug for long-term studies. Ro1 retains the binding site for *nor*BNI, and double tg mice treated with *nor*BNI after dox removal failed to develop ventriculomegaly or ependymal denudation after 30 days. Thus the signaling cascades activated by Ro1 are required for HC to occur; expression alone of Ro1 is insufficient to cause HC.

Elucidating the signaling pathways responsible for HC could provide valuable insights into HC pathology. While we know that Ro1 signaling triggers HC, it is unclear which cell types are primarily affected and should be the focus of further studies. Although GFAP positive cells are mostly astrocytes, astrocytes are not a homogenous population and

carry out a variety of diverse functions throughout the entire CNS. Other cells, including progenitors, ependymal cells and microglia, can express GFAP, particularly while undergoing development and maturation. A number of important postnatal developmental processes such as maturation of ependymal cells and cilia development occur during the first two postnatal weeks, coinciding with onset of HC in our model. However, it is unlikely that disruption of developmental processes by Ro1 signaling is the primary cause of HC because adult double tg mice kept on dox until p30 will also develop significantly enlarged ventricles and stenosis of the aqueduct when dox is removed. The rate of HC development is slower and variations in severity are more pronounced in adult onset, possibly reflecting differences in either genetic susceptibility or ability to compensate, yet the fact that even in adult onset all double tg mice develop HC indicates that Ro1 signaling is disrupting a critical process in normal CNS function.

**Possible mechanisms.** Ependymal cells are one of the cell types possibly affected by Ro1 expression. Ependymal denudation is seen across multiple experimental models of HC and in human cases. In at least one mouse model with stenosis of the aqueduct, denudation occurs in a distinct pattern early in the disease process, suggesting that denudation is not the result of stretching and could be the cause of HC (Raimondi et al., 1976; Jimenez et al., 2001). Other studies using neuraminidase to pharmacologically detach ependyma resulted in HC (Grondona et al., 1996). Because denudation was seen in Ro1 HC mice regardless of severity, we decided to focus on ependymal cells. While there is some contradiction in the literature regarding GFAP expression in ependyma (Bruni et al., 1985; Kasper et al., 1991; Sarnat, 1992; Xiuxin Liu, 2006), we were able to detect GFAP in ependymal cells by

immunocytochemistry. GFAP was not expressed by all ependyma, but ependyma are known to be a heterogeneous cell population, a fact that seems to be reflected in the discontinuous denudation seen in HC. Subsequently Ro1 expression in ependyma was confirmed by both immunocytochemistry and RT-PCR.

The consequences of Ro1 signaling in ependyma are not known. The classical action of  $G\alpha_i$  is to inhibit  $G\alpha_s$ -stimulated production of cAMP by adenylyl cyclase. While the functional consequences of altered cAMP levels are difficult to predict, both gain and loss of the  $G_s$ -coupled PAC1 receptor results in dysfunctional cilia, and abnormal cAMP and phosphoCREB levels in ependyma are found in some HC models (Otto et al., 2004; Banizs et al., 2005; Lang et al., 2006). Long term  $G\alpha_i$  activation also can sensitize some adenylyl cyclase isoforms, again with unknown functional effects (Thomas and Hoffman, 1996). More recently it has become clear that GPCRs have signaling functions outside of classical signaling pathways.  $G\alpha_i$  activates Rho via Rho guanine nucleotide exchange factors (Billington and Penn, 2003) and mitogen-activated protein kinase (MAPK) pathways can be activated via cross-talk of GPCRs with receptor tyrosine kinases and focal adhesion complexes, leading to transcriptional activation. Thus unregulated Ro1 signaling could result in abnormal gene transcription in ependyma, possibly resulting in denudation. GPCRs can act independently of G proteins, participating in protein signaling complexes with beta arrestins. These signaling complexes also activate MAPK pathways, but with a slower and longer-lasting activation that is concentrated mainly in the cytosol (Luttrell, 2005). The ERK1/2 (extracellular signal-regulated kinase) MAPK pathway regulates a number of processes, including proliferation and migration. Detachment could result from changes in adhesion molecules or gap junction proteins; other structural proteins could be affected,

including actin, myosin, tau proteins or microtubules. Changes in gene expression or signaling pathways that impact cilia function could also lead to HC in the Ro1 model.

**Gene arrays.** The decision was made to use gene arrays to determine what, if any, changes in ependymal gene expression are caused by Ro1 signaling. While other groups have done gene arrays on tissue collected from various brain regions from HC animals (Balasubramaniam and Del Bigio, 2002; Morgan et al., 2005; Miller et al., 2006; Ballarati et al., 2007; Nonaka et al., 2008; Zhang et al., 2008), our goal was to look for changes occurring in a specific cell type before the onset of HC. Thus the ependymal cells needed to be collected soon after onset of Ro1 expression to avoid changes that may occur in response to other pathology. The time course for dox to “wash” out of the CNS was unknown, so timed studies were done to look for Ro1 expression and ependymal detachment following removal of dox in adult mice. The appearance of Ro1 after just three days off dox was surprising as we had anticipated that the washout time would be longer. The mice had always been on dox, and dox is known to accumulate in bone. However, Ro1 expression was limited to the thalamus at 3 days off dox and no morphological changes of the ependyma could be detected. By seven days off dox Ro1 was seen in other areas of the brain, including the hippocampus and cortex, and detachment of ependyma was first seen in histological studies at 10 days off dox. Based on these results, 5 and 9 days after dox removal were the time points selected for gene arrays.

Unfortunately, the array data proved to be unusable with no genes being statistically differentially expressed between double tg mice and littermate controls at either time point. We do not interpret the data to mean that no changes occur in ependymal cells, but that the

design of the experiment was flawed and therefore no conclusions can be drawn from the data. Although no significantly changed genes were found to have a false discovery rate of less than one, pathway analysis (using the genes that were significantly changed between genotypes) found significant enrichment of pathways potentially relevant for denudation, including pathways for adhesion, cilia function, matrix proteins, and integrin signaling, suggesting that real changes may be occurring but cannot be adequately detected in our samples. This may be due to improper time point selection. The timed studies were done on a very limited number of animals and did not extend longer than 7 days for immunostaining or 9 days for RT-PCR. It is very likely that the range of animal to animal variability of Ro1 expression was not captured in these studies, and thus biological variation between our samples prevented the statistical detection of actual changes. Higher levels of Ro1 RNA were detected with RT-PCR at 9 days off dox than at 5 days off dox, suggesting longer time courses using RT-PCR would be very useful in determining the maximal level of Ro1 expression in ependymal cells so we would better understand if the levels seen at 5 and 9 days off dox should be sufficient to detect changes if present. Another source of error could come from the cellular composition of our samples. Isolation of the ependymal cell layer is technically challenging, and inclusion of some underlying subependymal cells is unavoidable. If actual changes in ependymal cells at these timepoints are small, the presence of another cell type could potentially dilute the sample.

Several attempts were made to verify by RT-PCR changes in selected genes from each gene array. The initial experiments proved to be unreliable due to inefficient reverse transcriptase activity resulting from very low amounts of total RNA used. Subsequent attempts using total RNA combined from three mice per sample produced more reliable

results, but no significant changes were found in the expression of the selected targets. Using a higher n would perhaps balance out biological variability, however, based on the questionable array data and preliminary RT-PCR results continuing the present study did not seem prudent.

**Alternate hypotheses.** Ro1 signaling may not cause changes at the transcriptional level in ependymal cells; denudation may instead result from altered signaling pathways in the cytosol. It is also possible that ependymal cell denudation is not the cause of HC in the Ro1 model. Ro1 signaling could impact the function of the subcommissural organ (SCO). Cells of the SCO strongly express GFAP, and Ro1 signaling in these cells could alter their secretory activity. The glycoproteins secreted by the SCO are theorized to play critical roles in keeping the narrow aqueduct patent and possibly in regulation CSF formation by the choroid plexus, thus loss of glycoprotein secretion could result in stenosis of the aqueduct and onset of HC. Ro1 signaling could also change the ability of astrocytes to carry out normal functions in the CNS. Astrocytes are found throughout the parenchyma and also directly contact the endothelial cells of blood vessels. Given the emerging importance of CSF absorption into venous blood, changes in astrocytic  $G_i$  signaling that impact the ability of CSF to move through the interstitial spaces or to be absorbed into blood vessels, perhaps by increasing production of extracellular matrix proteins (as seen in the S100 $\beta$  tg model) or by causing constriction of blood vessels, could result in the accumulation of CSF and onset of HC.

**Future directions.** The Ro1 HC model presents the field an opportunity to study the earliest events in HC pathology. Although treating HC by shunting has greatly improved the life expectancy and quality of life for patients with HC, new treatments capable of preventing neurological damage or motor deficits are greatly needed. Future studies of the Ro1 model should focus on understanding the Ro1 receptor and how enhanced  $G_i$  signaling in GFAP positive cells results in HC. While the affinity of Ro1 for endogenous KOR ligands is significantly reduced, overexpression of Ro1 may allow activation of the receptor by ligands that normally would not bind. Using a KOR antagonist, other than norBIN (an inverse agonist), to see if blocking signaling while maintaining intrinsic activity of Ro1 still results in HC would provide insight into whether Ro1 may be activated by endogenous ligands in our model.

Further work is needed to determine if ependymal cells are primarily involved in HC pathology. The timed studies previously mentioned should be expanded to gain a better understanding of the time range needed for Ro1 expression and ependymal denudation to begin after dox removal. Included in these studies should be immunostaining time courses designed to look for any set patterns of Ro1 expression, as well as RT-PCR experiments for detecting Ro1 levels in ependyma. Ependymal cells should be collected at the new time points established from these experiments and studied for changes at the kinase/phosphatase or protein levels.

To determine if SCO functions are impacted by Ro1 signaling, anti-RF antibodies could be used to look for changes in SCO secretions. If differences in glycoproteins are seen in Western blots of CSF from sick animals, a timed study should be done to see when these changes occur in relation to pathology. Changes in protein levels or phosphorylation of

proteins involved in vesicle docking and fusion could also be examined to see if secretory functions are impaired in Ro1 mice. PNA and RCA lectin binding to ependymal surfaces could also be used to determine if glycosylation is normal in ependyma.

Other potential areas of study include the progenitor cells of the subventricular zone (SVZ), cilia function, and extracellular matrix formation around blood vessels. Progenitor cells in the SVZ, specifically the B cells, express GFAP and thus should also express Ro1. Changes in the proliferation of progenitors may contribute to the pathology of HC; BrdU incorporation experiments in both immature and adult mice could provide insight on the impact of G<sub>i</sub> signaling in these populations. Immunostaining for SVZ cell markers could show alterations in relative numbers of A, B, and C cells and potential changes in migration. Previous  $\alpha$ -tubulin and H&E staining of double tg mice showed that cilia are present on ependymal cells, but did not provide information on whether cilia structure is normal. Using brain slices and fluorescent dyes in a fluid chamber would show if cilia are able to beat normally. Additional immunostaining using antibodies against various cilia proteins would also help identify any abnormalities in cilia structure and function. Changes in extracellular matrix proteins could also be detected by Western blots and immunostaining.

The development of hydrocephalus in the Ro1 transgenic mouse has resulted in a new inducible model for hydrocephalus with the potential to provide new insights into the causes of this disease. Finding pharmacological treatments for HC has been hindered by a lack of understanding of the molecular pathways that cause HC and are responsible for the neurological and motor deficits that frequently accompany the disease even after successful shunting. We have characterized the Ro1 model, confirming that the GFAP promoter drives expression of Ro1 and determining that Ro1 signaling is causing the onset of HC. While



some of the initial attempts at elucidating the signaling pathways perturbed by Ro1 were unsuccessful, the Ro1 model remains an exciting tool for HC research and future studies with the Ro1 mice have the potential to impact the course of HC treatment.

## LITERATURE CITED

- Abbott NJ (2004) Evidence for bulk flow of brain interstitial fluid: significance for physiology and pathology. *Neurochem Int* 45:545-552.
- Abbott NJ, Bundgaard M, Cserr HF (1985) Tightness of the blood-brain barrier and evidence for brain interstitial fluid flow in the cuttlefish, *Sepia officinalis*. *J Physiol* 368:213-226.
- Abdolvahabi RM, Mitchell JA, Diaz FG, McAllister JP, 2nd (2000) A brief review of the effects of chronic hydrocephalus on the gonadotropin releasing hormone system: implications for amenorrhea and precocious puberty. *Neurol Res* 22:123-126.
- Abramovitz M, Homma H, Ishigaki S, Tansey F, Cammer W, Listowsky I (1988) Characterization and localization of glutathione-S-transferases in rat brain and binding of hormones, neurotransmitters, and drugs. *J Neurochem* 50:50-57.
- Acikgoz B, Akpinar G, Bingol N, Usseli I (1999) Angiotensin II receptor content within the circumventricular organs increases after experimental hydrocephalus in rats. *Acta Neurochir (Wien)* 141:1095-1099.
- Ahlberg J, Norlen L, Blomstrand C, Wikkelso C (1988) Outcome of shunt operation on urinary incontinence in normal pressure hydrocephalus predicted by lumbar puncture. *J Neurol Neurosurg Psychiatry* 51:105-108.
- al-Shroof M, Karnik AM, Karnik AA, Longshore J, Sliman NA, Khan FA (2001) Ciliary dyskinesia associated with hydrocephalus and mental retardation in a Jordanian family. *Mayo Clin Proc* 76:1219-1224.
- Albeck MJ, Skak C, Nielsen PR, Olsen KS, Borgesen SE, Gjerris F (1998) Age dependency of resistance to cerebrospinal fluid outflow. *J Neurosurg* 89:275-278.
- Albrechtsen M, Sorensen PS, Gjerris F, Bock E (1985) High cerebrospinal fluid concentration of glial fibrillary acidic protein (GFAP) in patients with normal pressure hydrocephalus. *J Neurol Sci* 70:269-274.
- Ammirati M, Raimondi AJ (1987) Cerebrospinal fluid shunt infections in children. A study on the relationship between the etiology of hydrocephalus, age at the time of shunt placement, and infection rate. *Childs Nerv Syst* 3:106-109.
- Aoyama Y, Kinoshita Y, Yokota A, Hamada T (2006) Neuronal damage in hydrocephalus and its restoration by shunt insertion in experimental hydrocephalus: a study involving the neurofilament-immunostaining method. *J Neurosurg* 104:332-339.
- Araque A, Carmignoto G, Haydon PG (2001) Dynamic signaling between astrocytes and neurons. *Annu Rev Physiol* 63:795-813.

- Araque A, Sanzgiri RP, Parpura V, Haydon PG (1998) Calcium elevation in astrocytes causes an NMDA receptor-dependent increase in the frequency of miniature synaptic currents in cultured hippocampal neurons. *J Neurosci* 18:6822-6829.
- Araque A, Parpura V, Sanzgiri RP, Haydon PG (1999) Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* 22:208-215.
- Araque A, Li N, Doyle RT, Haydon PG (2000) SNARE protein-dependent glutamate release from astrocytes. *J Neurosci* 20:666-673.
- Araque A, Martin ED, Perea G, Arellano JI, Buno W (2002) Synaptically released acetylcholine evokes Ca<sup>2+</sup> elevations in astrocytes in hippocampal slices. *J Neurosci* 22:2443-2450.
- Arkhinova SE (1966) [Varieties in the structure and topography of arachnoidal villi in the human brain]. *Arkh Anat Gistol Embriol* 51:51-61.
- Arnold PM, Ma JY, Citron BA, Festoff BW (1999) Insulin-like growth factor binding proteins in cerebrospinal fluid during human development and aging. *Biochem Biophys Res Commun* 264:652-656.
- Asanuma H, Keller A (1991) Neuronal mechanisms of motor learning in mammals. *Neuroreport* 2:217-224.
- Ata AK, Funa K, Olsson Y (1997) Expression of various TGF-beta isoforms and type I receptor in necrotizing human brain lesions. *Acta Neuropathol* 93:326-333.
- Baas D, Meiniel A, Benadiba C, Bonnafé E, Meiniel O, Reith W, Durand B (2006) A deficiency in RFX3 causes hydrocephalus associated with abnormal differentiation of ependymal cells. *Eur J Neurosci* 24:1020-1030.
- Bachs A, Walker AE (1953) Experimental hydrocephalus. *J Neuropathol Exp Neurol* 12:283-292.
- Bahe S, Stierhof YD, Wilkinson CJ, Leiss F, Nigg EA (2005) Rootletin forms centriole-associated filaments and functions in centrosome cohesion. *J Cell Biol* 171:27-33.
- Bajpai M, Kataria R, Bhatnagar V, Agarwala S, Gupta DK, Bharadwaj M, Das K, Alladi A, Lama T, Srinivas M, Dave S, Arora M, Dutta H, Pandey RM, Mitra DK (1997) Management of hydrocephalus. *Indian J Pediatr* 64:48-56.
- Balasubramaniam J, Del Bigio MR (2002) Analysis of age-dependant alteration in the brain gene expression profile following induction of hydrocephalus in rats. *Exp Neurol* 173:105-113.
- Ball MJ (1976) Neurofibrillary tangles in the dementia of "normal pressure" hydrocephalus. *Can J Neurol Sci* 3:227-235.

- Ball MJ, Vis CL (1978) Relationship of granulovacuolar degeneration in hippocampal neurones to aging and to dementia in normal-pressure hydrocephalics. *J Gerontol* 33:815-824.
- Ballarati L, Rossi E, Bonati MT, Gimelli S, Maraschio P, Finelli P, Giglio S, Lapi E, Bedeschi MF, Gueneri S, Arrigo G, Patricelli MG, Mattina T, Guzzardi O, Pecile V, Police A, Scarano G, Larizza L, Zuffardi O, Giardino D (2007) 13q Deletion and central nervous system anomalies: further insights from karyotype-phenotype analyses of 14 patients. *J Med Genet* 44:e60.
- Banizs B, Komlosi P, Bevensee MO, Schwiebert EM, Bell PD, Yoder BK (2007) Altered pH(i) regulation and Na(+)/HCO3(-) transporter activity in choroid plexus of cilia-defective Tg737(orpk) mutant mouse. *Am J Physiol Cell Physiol* 292:C1409-1416.
- Banizs B, Pike MM, Millican CL, Ferguson WB, Komlosi P, Sheetz J, Bell PD, Schwiebert EM, Yoder BK (2005) Dysfunctional cilia lead to altered ependyma and choroid plexus function, and result in the formation of hydrocephalus. *Development* 132:5329-5339.
- Bannister CM, Mundy JE (1979) Some scanning electron microscopic observations of the ependymal surface of the ventricles of hydrocephalic Hy3 mice and a human infant. *Acta Neurochir (Wien)* 46:159-168.
- Barkovich AJ, Edwards MS (1992) Applications of neuroimaging in hydrocephalus. *Pediatr Neurosurg* 18:65-83.
- Barone FC, White RF, Spera PA, Ellison J, Currie RW, Wang X, Feuerstein GZ (1998) Ischemic preconditioning and brain tolerance: temporal histological and functional outcomes, protein synthesis requirement, and interleukin-1 receptor antagonist and early gene expression. *Stroke* 29:1937-1950; discussion 1950-1931.
- Baskin JJ, Manwaring KH, Rekate HL (1998a) Ventricular shunt removal: the ultimate treatment of the slit ventricle syndrome. *J Neurosurg* 88:478-484.
- Baskin JJ, Vishteh AG, Wesche DE, Rekate HL, Carrion CA (1998b) Ventriculoperitoneal shunt failure as a complication of laparoscopic surgery. *Jsls* 2:177-180.
- Bateman GA (2000) Vascular compliance in normal pressure hydrocephalus. *AJNR Am J Neuroradiol* 21:1574-1585.
- Batiz LF, Paez P, Jimenez AJ, Rodriguez S, Wagner C, Perez-Figares JM, Rodriguez EM (2006) Heterogeneous expression of hydrocephalic phenotype in the hyh mice carrying a point mutation in alpha-SNAP. *Neurobiol Dis* 23:152-168.
- Baykan N, Isbir O, Gercek A, Dagcinar A, Ozek MM (2005) Ten years of experience with pediatric neuroendoscopic third ventriculostomy: features and perioperative complications of 210 cases. *J Neurosurg Anesthesiol* 17:33-37.

- Bayle J, Letard S, Frank R, Dubreuil P, De Sepulveda P (2004) Suppressor of cytokine signaling 6 associates with KIT and regulates KIT receptor signaling. *J Biol Chem* 279:12249-12259.
- Bech RA, Juhler M, Waldemar G, Klinken L, Gjerris F (1997) Frontal brain and leptomeningeal biopsy specimens correlated with cerebrospinal fluid outflow resistance and B-wave activity in patients suspected of normal-pressure hydrocephalus. *Neurosurgery* 40:497-502.
- Becker DP, Nulsen FE (1968) Control of hydrocephalus by valve-regulated venous shunt: avoidance of complications in prolonged shunt maintenance. *J Neurosurg* 28:215-226.
- Beems T, Grotenhuis JA (2002) Is the success rate of endoscopic third ventriculostomy age-dependent? An analysis of the results of endoscopic third ventriculostomy in young children. *Childs Nerv Syst* 18:605-608.
- Beems T, Simons KS, Van Geel WJ, De Reus HP, Vos PE, Verbeek MM (2003) Serum- and CSF-concentrations of brain specific proteins in hydrocephalus. *Acta Neurochir (Wien)* 145:37-43.
- Belcheva MM, Clark AL, Haas PD, Serna JS, Hahn JW, Kiss A, Coscia CJ (2005) Mu and kappa opioid receptors activate ERK/MAPK via different protein kinase C isoforms and secondary messengers in astrocytes. *J Biol Chem* 280:27662-27669.
- Bellur SN, Chandra V, McDonald LW (1980) Arachnoidal cell hyperplasia. Its relationship to aging and chronic renal failure. *Arch Pathol Lab Med* 104:414-416.
- Benavides J, Quarteronet D, Imbault F, Malgouris C, Uzan A, Renault C, Dubroeuq MC, Gueremy C, Le Fur G (1983) Labelling of "peripheral-type" benzodiazepine binding sites in the rat brain by using [3H]PK 11195, an isoquinoline carboxamide derivative: kinetic studies and autoradiographic localization. *J Neurochem* 41:1744-1750.
- Beni-Adani L, Biani N, Ben-Sirah L, Constantini S (2006) The occurrence of obstructive vs absorptive hydrocephalus in newborns and infants: relevance to treatment choices. *Childs Nerv Syst* 22:1543-1563.
- Benveniste EN (1998) Cytokine actions in the central nervous system. *Cytokine Growth Factor Rev* 9:259-275.
- Bergsneider M (2001) Evolving concepts of cerebrospinal fluid physiology. *Neurosurg Clin N Am* 12:631-638, vii.
- Bergsneider M, Alwan AA, Falkson L, Rubinstein EH (1998) The relationship of pulsatile cerebrospinal fluid flow to cerebral blood flow and intracranial pressure: a new theoretical model. *Acta Neurochir Suppl* 71:266-268.

- Bickers DS, Adams RD (1949) Hereditary stenosis of the aqueduct of Sylvius as a cause of congenital hydrocephalus. *Brain* 72:246-262.
- Bie B, Pan ZZ (2003) Presynaptic mechanism for anti-analgesic and anti-hyperalgesic actions of kappa-opioid receptors. *J Neurosci* 23:7262-7268.
- Bifulco M, Laezza C, Stingo S, Wolff J (2002) 2',3'-Cyclic nucleotide 3'-phosphodiesterase: a membrane-bound, microtubule-associated protein and membrane anchor for tubulin. *Proc Natl Acad Sci U S A* 99:1807-1812.
- Billington CK, Penn RB (2003) Signaling and regulation of G protein-coupled receptors in airway smooth muscle. *Respir Res* 4:2.
- Birch NP, Rodriguez C, Dixon JE, Mezey E (1990) Distribution of carboxypeptidase H messenger RNA in rat brain using in situ hybridization histochemistry: implications for neuropeptide biosynthesis. *Brain Res Mol Brain Res* 7:53-59.
- Blaauwgeers HG, Sillevs Smitt PA, De Jong JM, Troost D (1993) Distribution of metallothionein in the human central nervous system. *Glia* 8:62-70.
- Black M, Graham DI (2002) Sudden unexplained death in adults caused by intracranial pathology. *J Clin Pathol* 55:44-50.
- Blackshear PJ, Graves JP, Stumpo DJ, Cobos I, Rubenstein JL, Zeldin DC (2003) Graded phenotypic response to partial and complete deficiency of a brain-specific transcript variant of the winged helix transcription factor RFX4. *Development* 130:4539-4552.
- Blomsterwall E, Bilting M, Stephensen H, Wikkelsø C (1995) Gait abnormality is not the only motor disturbance in normal pressure hydrocephalus. *Scand J Rehabil Med* 27:205-209.
- Blondel O, Collin C, McCarran WJ, Zhu S, Zamostiano R, Gozes I, Brenneman DE, McKay RD (2000) A glia-derived signal regulating neuronal differentiation. *J Neurosci* 20:8012-8020.
- Blount JP, Campbell JA, Haines SJ (1993) Complications in ventricular cerebrospinal fluid shunting. *Neurosurg Clin N Am* 4:633-656.
- Bonadio WA, Stanco L, Bruce R, Barry D, Smith D (1992) Reference values of normal cerebrospinal fluid composition in infants ages 0 to 8 weeks. *Pediatr Infect Dis J* 11:589-591.
- Bondurant CP, Jimenez DF (1995) Epidemiology of cerebrospinal fluid shunting. *Pediatr Neurosurg* 23:254-258; discussion 259.
- Boon AJ, Tans JT, Delwel EJ, Egeler-Peerdeman SM, Hanlo PW, Wurzer HA, Hermans J (1999) Dutch Normal-Pressure Hydrocephalus Study: the role of cerebrovascular disease. *J Neurosurg* 90:221-226.

- Booz KH (1975) Secretory phenomena at the ependyma of the IIIrd ventricle of the embryonic rat. *Anat Embryol (Berl)* 147:143-159.
- Booz KH, Wiesen B (1976) Fluorescence microscopic study of the behaviour of dans-marked histidine in the rat brain after intraventricular injection. *Anat Embryol (Berl)* 149:225-239.
- Botez MI, Ethier R, Leveille J, Botez-Marquard T (1977) A syndrome of early recognition of occult hydrocephalus and cerebral atrophy. *Q J Med* 46:365-380.
- Bottner M, Unsicker K, Suter-Crazzolara C (1996) Expression of TGF-beta type II receptor mRNA in the CNS. *Neuroreport* 7:2903-2907.
- Bourne A, Barnes K, Taylor BA, Turner AJ, Kenny AJ (1989) Membrane peptidases in the pig choroid plexus and on other cell surfaces in contact with the cerebrospinal fluid. *Biochem J* 259:69-80.
- Bouzinova EV, Praetorius J, Virkki LV, Nielsen S, Boron WF, Aalkjaer C (2005) Na<sup>+</sup>-dependent HCO<sub>3</sub><sup>-</sup> uptake into the rat choroid plexus epithelium is partially DIDS sensitive. *Am J Physiol Cell Physiol* 289:C1448-1456.
- Bradbury MW (1985) The blood-brain barrier. Transport across the cerebral endothelium. *Circ Res* 57:213-222.
- Bradley WG (2001a) Normal pressure hydrocephalus and deep white matter ischemia: which is the chicken, and which is the egg? *AJNR Am J Neuroradiol* 22:1638-1640.
- Bradley WG, Jr. (2001b) Diagnostic tools in hydrocephalus. *Neurosurg Clin N Am* 12:661-684, viii.
- Bradley WG, Jr., Whittemore AR, Watanabe AS, Davis SJ, Teresi LM, Homyak M (1991) Association of deep white matter infarction with chronic communicating hydrocephalus: implications regarding the possible origin of normal-pressure hydrocephalus. *AJNR Am J Neuroradiol* 12:31-39.
- Brailov I, Bancila M, Brisorgueil MJ, Miquel MC, Hamon M, Verge D (2000) Localization of 5-HT(6) receptors at the plasma membrane of neuronal cilia in the rat brain. *Brain Res* 872:271-275.
- Braun KP, Vandertop WP, Gooskens RH, Tulleken KA, Nicolay K (2000) NMR spectroscopic evaluation of cerebral metabolism in hydrocephalus: a review. *Neurol Res* 22:51-64.
- Braun KP, de Graaf RA, Vandertop WP, Gooskens RH, Tulleken KA, Nicolay K (1998) In vivo <sup>1</sup>H MR spectroscopic imaging and diffusion weighted MRI in experimental hydrocephalus. *Magn Reson Med* 40:832-839.

- Braun KP, Dijkhuizen RM, de Graaf RA, Nicolay K, Vandertop WP, Gooskens RH, Tulleken KA (1997) Cerebral ischemia and white matter edema in experimental hydrocephalus: a combined in vivo MRI and MRS study. *Brain Res* 757:295-298.
- Braun KP, van Eijsden P, Vandertop WP, de Graaf RA, Gooskens RH, Tulleken KA, Nicolay K (1999) Cerebral metabolism in experimental hydrocephalus: an in vivo <sup>1</sup>H and <sup>31</sup>P magnetic resonance spectroscopy study. *J Neurosurg* 91:660-668.
- Bret P, Chazal J, Janny P, Renaud B, Tommasi M, Lemaire JJ, Bret M, Jouvet A (1990) [Chronic hydrocephalus in adults]. *Neurochirurgie* 36 Suppl 1:1-159.
- Brightman MW (1965) The distribution within the brain of ferritin injected into cerebrospinal fluid compartments. I. Ependymal distribution. *J Cell Biol* 26:99-123.
- Broadwell RD, Sofroniew MV (1993) Serum proteins bypass the blood-brain fluid barriers for extracellular entry to the central nervous system. *Exp Neurol* 120:245-263.
- Brodbeck A, Stoodley M (2007) CSF pathways: a review. *Br J Neurosurg* 21:510-520.
- Brouns MR, Matheson SF, Hu KQ, Delalle I, Caviness VS, Silver J, Bronson RT, Settleman J (2000) The adhesion signaling molecule p190 RhoGAP is required for morphogenetic processes in neural development. *Development* 127:4891-4903.
- Brown PD, Davies SL, Speake T, Millar ID (2004) Molecular mechanisms of cerebrospinal fluid production. *Neuroscience* 129:957-970.
- Brummendorf T, Rathjen FG (1995) Cell adhesion molecules 1: immunoglobulin superfamily. *Protein Profile* 2:963-1108.
- Bruni JE, Del Bigio MR, Clattenburg RE (1985) Ependyma: normal and pathological. A review of the literature. *Brain Res* 356:1-19.
- Bruni JE, del Bigio MR, Cardoso ER, Persaud TV (1988a) Neuropathology of congenital hydrocephalus in the SUMS/NP mouse. *Acta Neurochir (Wien)* 92:118-122.
- Bruni JE, Del Bigio MR, Cardoso ER, Persaud TV (1988b) Hereditary hydrocephalus in laboratory animals and humans. *Exp Pathol* 35:239-246.
- Bugalho P, Guimaraes J (2007) Gait disturbance in normal pressure hydrocephalus: a clinical study. *Parkinsonism Relat Disord* 13:434-437.
- Buki A, Siman R, Trojanowski JQ, Povlishock JT (1999) The role of calpain-mediated spectrin proteolysis in traumatically induced axonal injury. *J Neuropathol Exp Neurol* 58:365-375.
- Bulat M, Lupret V, Orekhovic D, Klarica M (2008) Transventricular and transpial absorption of cerebrospinal fluid into cerebral microvessels. *Coll Antropol* 32 Suppl 1:43-50.



- Bunn SJ, Hanley MR, Wilkin GP (1985) Evidence for a kappa-opioid receptor on pituitary astrocytes: an autoradiographic study. *Neurosci Lett* 55:317-323.
- Bushong EA, Martone ME, Jones YZ, Ellisman MH (2002) Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains. *J Neurosci* 22:183-192.
- Butelman ER, Negus SS, Ai Y, de Costa BR, Woods JH (1993) Kappa opioid antagonist effects of systemically administered nor- binaltorphimine in a thermal antinociception assay in rhesus monkeys. *J Pharmacol Exp Ther* 267:1269-1276.
- Buxton N, Cartmill M, Vloeberghs M (1999) Endoscopic third ventriculostomy: outcome analysis of 100 consecutive procedures. *Neurosurgery* 45:957-959.
- Buxton N, Macarthur D, Mallucci C, Punt J, Vloeberghs M (1998) Neuroendoscopic third ventriculostomy in patients less than 1 year old. *Pediatr Neurosurg* 29:73-76.
- Carmignoto G, Pizzorusso T, Tia S, Vicini S (1997) Brain-derived neurotrophic factor and nerve growth factor potentiate excitatory synaptic transmission in the rat visual cortex. *J Physiol* 498 ( Pt 1):153-164.
- Casey AT, Kimmings EJ, Kleinlugtebeld AD, Taylor WA, Harkness WF, Hayward RD (1997) The long-term outlook for hydrocephalus in childhood. A ten-year cohort study of 155 patients. *Pediatr Neurosurg* 27:63-70.
- Casmiro M, Benassi G, Cacciatore FM, D'Alessandro R (1989a) Frequency of idiopathic normal pressure hydrocephalus. *Arch Neurol* 46:608.
- Casmiro M, D'Alessandro R, Cacciatore FM, Daidone R, Calbucci F, Lugaresi E (1989b) Risk factors for the syndrome of ventricular enlargement with gait apraxia (idiopathic normal pressure hydrocephalus): a case-control study. *J Neurol Neurosurg Psychiatry* 52:847-852.
- Castaneyra-Perdomo A, Meyer G, Carmona-Calero E, Banuelos-Pineda J, Mendez-Medina R, Ormazabal-Ramos C, Ferres-Torres R (1994) Alterations of the subcommissural organ in the hydrocephalic human fetal brain. *Brain Res Dev Brain Res* 79:316-320.
- Castaneyra-Perdomo A, Munoz R, Gonzalez-Marrero I, Fernandez-Rodriguez P, Perez-Gonzalez H, Castaneyra-Ruiz L, Castaneyra-Ruiz A, Ormazabal-Ramos C, Troyano J, Carmona-Calero E (2006) Protein changes in the cerebrospinal fluid of the different types of perinatal human hydrocephalus. *Cerebrospinal Fluid Research* 3:S56.
- Chae TH, Kim S, Marz KE, Hanson PI, Walsh CA (2004) The hyh mutation uncovers roles for alpha Snap in apical protein localization and control of neural cell fate. *Nat Genet* 36:264-270.
- Chakrabarti S, Fombonne E (2001) Pervasive developmental disorders in preschool children. *Jama* 285:3093-3099.

- Chemelli RM, Willie JT, Sinton CM, Elmquist JK, Scammell T, Lee C, Richardson JA, Williams SC, Xiong Y, Kisanuki Y, Fitch TE, Nakazato M, Hammer RE, Saper CB, Yanagisawa M (1999) Narcolepsy in orexin knockout mice: molecular genetics of sleep regulation. *Cell* 98:437-451.
- Chen C, Diaz Brinton RD, Shors TJ, Thompson RF (1993) Vasopressin induction of long-lasting potentiation of synaptic transmission in the dentate gyrus. *Hippocampus* 3:193-203.
- Chen J, Knowles HJ, Hebert JL, Hackett BP (1998) Mutation of the mouse hepatocyte nuclear factor/forkhead homologue 4 gene results in an absence of cilia and random left-right asymmetry. *J Clin Invest* 102:1077-1082.
- Chen J, Chang S, Duncan SA, Okano HJ, Fishell G, Aderem A (1996) Disruption of the MacMARCKS gene prevents cranial neural tube closure and results in anencephaly. *Proc Natl Acad Sci U S A* 93:6275-6279.
- Cherian S, Whitelaw A, Thoresen M, Love S (2004a) The pathogenesis of neonatal post-hemorrhagic hydrocephalus. *Brain Pathol* 14:305-311.
- Cherian S, Thoresen M, Silver IA, Whitelaw A, Love S (2004b) Transforming growth factor-beta<sub>1</sub> in a rat model of neonatal posthaemorrhagic hydrocephalus. *Neuropathol Appl Neurobiol* 30:585-600.
- Choi KH, Kim SI, Yoon SY, Kim JH, Kang SW, Ha SK, Lee HY, Han DS, Kim YS, Park K, Jeong HJ, Kim DK (2001) Long-term outcome of kidney transplantation in adult recipients with focal segmental glomerulosclerosis. *Yonsei Med J* 42:209-214.
- Chumas PD, Drake JM, Del Bigio MR, Da Silva M, Tuor UI (1994) Anaerobic glycolysis preceding white-matter destruction in experimental neonatal hydrocephalus. *J Neurosurg* 80:491-501.
- Chung JH, Whiteley M, Felsenfeld G (1993) A 5' element of the chicken beta-globin domain serves as an insulator in human erythroid cells and protects against position effect in *Drosophila*. *Cell* 74:505-514.
- Cifuentes M, Rodriguez S, Perez J, Grondona JM, Rodriguez EM, Fernandez-Llebrez P (1994) Decreased cerebrospinal fluid flow through the central canal of the spinal cord of rats immunologically deprived of Reissner's fibre. *Exp Brain Res* 98:431-440.
- Cinalli G, Salazar C, Mallucci C, Yada JZ, Zerah M, Sainte-Rose C (1998) The role of endoscopic third ventriculostomy in the management of shunt malfunction. *Neurosurgery* 43:1323-1327; discussion 1327-1329.
- Cinalli G, Sainte-Rose C, Chumas P, Zerah M, Brunelle F, Lot G, Pierre-Kahn A, Renier D (1999) Failure of third ventriculostomy in the treatment of aqueductal stenosis in children. *Neurosurg Focus* 6:e3.

- Cirovic S, Walsh C, Fraser WD (2003) Mathematical study of the role of non-linear venous compliance in the cranial volume-pressure test. *Med Biol Eng Comput* 41:579-588.
- Clark S, Sangra M, Hayhurst C, Kandasamy J, Jenkinson M, Lee M, Mallucci C (2008) The use of noninvasive electromagnetic neuronavigation for slit ventricle syndrome and complex hydrocephalus in a pediatric population. *J Neurosurg Pediatrics* 2:430-434.
- Cloez I, Tayarani I, Morel F, Bourre JM (1989) Alterations in protective enzymes against peroxidation in the central and peripheral nervous system of control and dysmyelinating mutant mice. *J Neurochem* 52:1353-1358.
- Cohen AR, Leifer DW, Zechel M, Flanigan DP, Lewin JS, Lust WD (1999) Characterization of a model of hydrocephalus in transgenic mice. *J Neurosurg* 91:978-988.
- Cohen NR, Taylor JS, Scott LB, Guillery RW, Soriano P, Furley AJ (1998) Errors in corticospinal axon guidance in mice lacking the neural cell adhesion molecule L1. *Curr Biol* 8:26-33.
- Collins P (1979) Experimental obstructive hydrocephalus in the rat: a scanning electron microscopic study. *Neuropathol Appl Neurobiol* 5:457-468.
- Collins P, Woollam DH (1979) The ventricular surface of the subcommissural organ: a scanning and transmission electron microscopic study. *J Anat* 129:623-631.
- Connell HM, McConnel TS (1981) Psychiatric sequelae in children treated operatively for hydrocephalus in infancy. *Dev Med Child Neurol* 23:505-517.
- Costa T, Cotecchia S (2005) Historical review: Negative efficacy and the constitutive activity of G-protein-coupled receptors. *Trends Pharmacol Sci* 26:618-624.
- Coward P, Wada HG, Falk MS, Chan SD, Meng F, Akil H, Conklin BR (1998) Controlling signaling with a specifically designed Gi-coupled receptor. *Proc Natl Acad Sci U S A* 95:352-357.
- Crews L, Wyss-Coray T, Masliah E (2004) Insights into the pathogenesis of hydrocephalus from transgenic and experimental animal models. *Brain Pathol* 14:312-316.
- Crone C, Christensen O (1981) Electrical resistance of a capillary endothelium. *J Gen Physiol* 77:349-371.
- Csillik AE, Okuno E, Csillik B, Knyihar E, Vecsei L (2002) Expression of kynurenine aminotransferase in the subplate of the rat and its possible role in the regulation of programmed cell death. *Cereb Cortex* 12:1193-1201.
- Czosnyka M, Czosnyka ZH, Whitfield PC, Donovan T, Pickard JD (2001) Age dependence of cerebrospinal pressure-volume compensation in patients with hydrocephalus. *J Neurosurg* 94:482-486.

- D'Ambrosio R, Gordon DS, Winn HR (2002) Differential role of KIR channel and Na(+)/K(+)-pump in the regulation of extracellular K(+) in rat hippocampus. *J Neurophysiol* 87:87-102.
- da Silva MC, Drake JM, Lemaire C, Cross A, Tuor UI (1994) High-energy phosphate metabolism in a neonatal model of hydrocephalus before and after shunting. *J Neurosurg* 81:544-553.
- da Silva MC, Michowicz S, Drake JM, Chumas PD, Tuor UI (1995) Reduced local cerebral blood flow in periventricular white matter in experimental neonatal hydrocephalus-restoration with CSF shunting. *J Cereb Blood Flow Metab* 15:1057-1065.
- Dahme M, Bartsch U, Martini R, Anliker B, Schachner M, Mantei N (1997) Disruption of the mouse L1 gene leads to malformations of the nervous system. *Nat Genet* 17:346-349.
- Dandy WE (1919) Experimental Hydrocephalus. *Ann Surg* 70:129-142.
- Danielian PS, McMahon AP (1996) Engrailed-1 as a target of the Wnt-1 signalling pathway in vertebrate midbrain development. *Nature* 383:332-334.
- Danik M, Chabot JG, Hassan-Gonzalez D, Suh M, Quirion R (1993) Localization of sulfated glycoprotein-2/clusterin mRNA in the rat brain by in situ hybridization. *J Comp Neurol* 334:209-227.
- Daszkiewicz P, Barszcz S (2007) Multiple shunt system revisions in patients with hydrocephalus - causes, effects, regularities and prognostic factors. *Neurol Neurochir Pol* 41:404-410.
- Dauvilliers Y, Baumann CR, Carlander B, Bischof M, Blatter T, Lecendreux M, Maly F, Besset A, Touchon J, Billiard M, Tafti M, Bassetti CL (2003) CSF hypocretin-1 levels in narcolepsy, Kleine-Levin syndrome, and other hypersomnias and neurological conditions. *J Neurol Neurosurg Psychiatry* 74:1667-1673.
- Davidoff LM, Chamlin M (1959) The "fontanometer"; adaptation of the Schiotz tonometer for the determination of intracranial pressure in the neonatal and early periods of infancy. *Pediatrics* 24:1065-1068.
- Davis SL, Tooley WH, Hunt JV (1987) Developmental outcome following posthemorrhagic hydrocephalus in preterm infants. Comparison of twins discordant for hydrocephalus. *Am J Dis Child* 141:1170-1174.
- Davy BE, Robinson ML (2003) Congenital hydrocephalus in hy3 mice is caused by a frameshift mutation in Hydin, a large novel gene. *Hum Mol Genet* 12:1163-1170.
- De Angelis E, Watkins A, Schafer M, Brummendorf T, Kenwrick S (2002) Disease-associated mutations in L1 CAM interfere with ligand interactions and cell-surface expression. *Hum Mol Genet* 11:1-12.

- De Santi MM, Magni A, Valletta EA, Gardi C, Lungarella G (1990) Hydrocephalus, bronchiectasis, and ciliary aplasia. *Arch Dis Child* 65:543-544.
- De SN (1950) A study of the changes in the brain in experimental internal hydrocephalus. *J Pathol Bacteriol* 62:197-208.
- Deckert-Schluter M, Schluter D, Hof H, Wiestler OD, Lassmann H (1994) Differential expression of ICAM-1, VCAM-1 and their ligands LFA-1, Mac-1, CD43, VLA-4, and MHC class II antigens in murine *Toxoplasma* encephalitis: a light microscopic and ultrastructural immunohistochemical study. *J Neuropathol Exp Neurol* 53:457-468.
- Deiber MP, Wise SP, Honda M, Catalan MJ, Grafman J, Hallett M (1997) Frontal and parietal networks for conditional motor learning: a positron emission tomography study. *J Neurophysiol* 78:977-991.
- Del Bigio MR (1989) Hydrocephalus-induced changes in the composition of cerebrospinal fluid. *Neurosurgery* 25:416-423.
- Del Bigio MR (1993) Neuropathological changes caused by hydrocephalus. *Acta Neuropathol* 85:573-585.
- Del Bigio MR (1995) The ependyma: a protective barrier between brain and cerebrospinal fluid. *Glia* 14:1-13.
- Del Bigio MR (2000) Calcium-mediated proteolytic damage in white matter of hydrocephalic rats? *J Neuropathol Exp Neurol* 59:946-954.
- Del Bigio MR (2001) Pathophysiologic consequences of hydrocephalus. *Neurosurg Clin N Am* 12:639-649, vii.
- Del Bigio MR (2004) Cellular damage and prevention in childhood hydrocephalus. *Brain Pathol* 14:317-324.
- Del Bigio MR, Bruni JE (1988) Changes in periventricular vasculature of rabbit brain following induction of hydrocephalus and after shunting. *J Neurosurg* 69:115-120.
- Del Bigio MR, Zhang YW (1998) Cell death, axonal damage, and cell birth in the immature rat brain following induction of hydrocephalus. *Exp Neurol* 154:157-169.
- Del Bigio MR, Massicotte EM (2001) Protective effect of nimodipine on behavior and white matter of rats with hydrocephalus. *J Neurosurg* 94:788-794.
- Del Bigio MR, Bruni JE, Fewer HD (1985) Human neonatal hydrocephalus. An electron microscopic study of the periventricular tissue. *J Neurosurg* 63:56-63.

- Del Bigio MR, Crook CR, Buist R (1997a) Magnetic resonance imaging and behavioral analysis of immature rats with kaolin-induced hydrocephalus: pre- and postshunting observations. *Exp Neurol* 148:256-264.
- Del Bigio MR, Cardoso ER, Halliday WC (1997b) Neuropathological changes in chronic adult hydrocephalus: cortical biopsies and autopsy findings. *Can J Neurol Sci* 24:121-126.
- Del Bigio MR, Kanfer JN, Zhang YW (1997c) Myelination delay in the cerebral white matter of immature rats with kaolin-induced hydrocephalus is reversible. *J Neuropathol Exp Neurol* 56:1053-1066.
- Del Bigio MR, Bruni JE, Vriend JP (1998) Monoamine neurotransmitters and their metabolites in the mature rabbit brain following induction of hydrocephalus. *Neurochem Res* 23:1379-1386.
- Del Bigio MR, Wang X, Wilson MJ (2002) Sodium channel-blocking agents are not of benefit to rats with kaolin-induced hydrocephalus. *Neurosurgery* 51:460-466; discussion 466-467.
- Del Bigio MR, Wilson MJ, Enno T (2003) Chronic hydrocephalus in rats and humans: white matter loss and behavior changes. *Ann Neurol* 53:337-346.
- del Pozo-Reyes D, Loyo-Varela M, Estanol-Vidal B, Mateos-Gomez JH (1978) Production and absorption of cerebrospinal fluid in the spinal subarachnoid space of the dog. *Arch Invest Med (Mex)* 9:511-518.
- Dennis M (2000) Developmental plasticity in children: the role of biological risk, development, time, and reserve. *J Commun Disord* 33:321-331; quiz 332.
- Di Chiro G (1966) Observations on the circulation of the cerebrospinal fluid. *Acta Radiol Diagn (Stockh)* 5:988-1002.
- Di Rocco C, Caldarelli M, Maira G, Rossi GF (1977) The study of cerebrospinal fluid dynamics in apparently 'arrested' hydrocephalus in children. *Childs Brain* 3:359-374.
- Di Terlizzi R, Platt S (2006) The function, composition and analysis of cerebrospinal fluid in companion animals: part I - function and composition. *Vet J* 172:422-431.
- Dickerman RD, McConathy WJ, Lustrin E, Schneider SJ (2003) Rapid neurological deterioration associated with minor head trauma in chronic hydrocephalus. *Childs Nerv Syst* 19:249-251; discussion 252-243.
- Ding Y, McAllister JP, 2nd, Yao B, Yan N, Canady AI (2001a) Neuron tolerance during hydrocephalus. *Neuroscience* 106:659-667.

- Ding Y, McAllister JP, 2nd, Yao B, Yan N, Canady AI (2001b) Axonal damage associated with enlargement of ventricles during hydrocephalus: a silver impregnation study. *Neurol Res* 23:581-587.
- Dobkin BH (1978) Syncope in the adult Chiari anomaly. *Neurology* 28:718-720.
- Dohrmann GJ (1971) The choroid plexus in experimental hydrocephalus. A light and electron microscopic study in normal, hydrocephalic, and shunted hydrocephalic dogs. *J Neurosurg* 34:56-69.
- Dominguez-Pinos MD, Paez P, Jimenez AJ, Weil B, Arraez MA, Perez-Figares JM, Rodriguez EM (2005) Ependymal denudation and alterations of the subventricular zone occur in human fetuses with a moderate communicating hydrocephalus. *J Neuropathol Exp Neurol* 64:595-604.
- Dominguez C, Broseta E, Alonso C, Munoz C, Server G, Jimenez-Cruz JF (1993) Epithelioid sarcoma of penis simulating Peyronie's disease. *Br J Urol* 72:975.
- Drayer BP, Rosenbaum AE (1977) Metrizamide brain penetrance. *Acta Radiol Suppl* 355:280-293.
- Du Boulay GH (1966) Pulsatile movements in the CSF pathways. *Br J Radiol* 39:255-262.
- Dusick JR, McArthur DL, Bergsneider M (2008) Success and complication rates of endoscopic third ventriculostomy for adult hydrocephalus: a series of 108 patients. *Surg Neurol* 69:5-15.
- Eckstein HB, Macnab GH (1966) Myelomeningocele and hydrocephalus. The impact of modern treatment. *Lancet* 1:842-845.
- Edwards RJ, Pople IK (2003) Death and third ventriculostomy. *J Neurosurg* 98:649-650; author reply 650-641.
- Edwards RJ, Dombrowski SM, Luciano MG, Pople IK (2004) Chronic hydrocephalus in adults. *Brain Pathol* 14:325-336.
- Egawa T, Mishima K, Egashira N, Fukuzawa M, Abe K, Yae T, Iwasaki K, Fujiwara M (2002) Impairment of spatial memory in kaolin-induced hydrocephalic rats is associated with changes in the hippocampal cholinergic and noradrenergic contents. *Behav Brain Res* 129:31-39.
- Elvidge AR, Branch CL, Thompson GB (1957) Observations in a case of hydrocephalus treated with diamox. *J Neurosurg* 14:628-638; discussion 638-629.
- Epstein BS (1952) Separation of the blood vessels from the walls of a lateral ventricle in rapidly progressing hydrocephalus. *Am J Roentgenol Radium Ther Nucl Med* 67:929-931.

- Eriksson PS, Hansson E, Ronnback L (1990) Delta and kappa opiate receptors in primary astroglial cultures from rat cerebral cortex. *Neurochem Res* 15:1123-1126.
- Erler T, Klaber HG (2001) [Treatment of posthemorrhagic hydrocephalus with intraventricular administration of recombinant plasminogen activator (rt-PA) and dexamethasone--possible prevention of permanent shunt implantation?]. *Z Geburtshilfe Neonatol* 205:27-32.
- Ersahin Y, Mutluer S, Guzelbag E (1994) Cerebrospinal fluid shunt infections. *J Neurosurg Sci* 38:161-165.
- Estivill-Torrus G, Vitalis T, Fernandez-Llebrez P, Price DJ (2001) The transcription factor Pax6 is required for development of the diencephalic dorsal midline secretory radial glia that form the subcommissural organ. *Mech Dev* 109:215-224.
- Factora R, Luciano M (2008) When to consider normal pressure hydrocephalus in the patient with gait disturbance. *Geriatrics* 63:32-37.
- Fan KJ, Pezeshkpour G (1987) Neurofibrillary tangles in association with congenital hydrocephalus. *J Natl Med Assoc* 79:1001-1003, 1006.
- Faraci FM, Mayhan WG, Heistad DD (1990) Vascular effects of acetazolamide on the choroid plexus. *J Pharmacol Exp Ther* 254:23-27.
- Feigin VL, Rinkel GJ, Algra A, Vermeulen M, van Gijn J (1998) Calcium antagonists in patients with aneurysmal subarachnoid hemorrhage: a systematic review. *Neurology* 50:876-883.
- Feng H, Huang G, Liao X, Fu K, Tan H, Pu H, Cheng Y, Liu W, Zhao D (2004) Endoscopic third ventriculostomy in the management of obstructive hydrocephalus: an outcome analysis. *J Neurosurg* 100:626-633.
- Fernandez-Llebrez P, Perez J, Cifuentes M, Alvial G, Rodriguez EM (1987) Immunocytochemical and ultrastructural evidence for a neurophysinergic innervation of the subcommissural organ of the snake *Natrix maura*. *Cell Tissue Res* 248:473-478.
- Fernandez-Llebrez P, Grondona JM, Perez J, Lopez-Aranda MF, Estivill-Torrus G, Llebrez-Zayas PF, Soriano E, Ramos C, Lallemand Y, Bach A, Robert B (2004) Msx1-deficient mice fail to form prosomere 1 derivatives, subcommissural organ, and posterior commissure and develop hydrocephalus. *J Neuropathol Exp Neurol* 63:574-586.
- Fernell E, Hagberg G, Hagberg B (1990) Infantile hydrocephalus--the impact of enhanced preterm survival. *Acta Paediatr Scand* 79:1080-1086.
- Fernell E, Gillberg C, von Wendt L (1991a) Behavioural problems in children with infantile hydrocephalus. *Dev Med Child Neurol* 33:388-395.



- Fernell E, Gillberg C, von Wendt L (1991b) Autistic symptoms in children with infantile hydrocephalus. *Acta Paediatr Scand* 80:451-457.
- Fiacco TA, McCarthy KD (2004) Intracellular astrocyte calcium waves in situ increase the frequency of spontaneous AMPA receptor currents in CA1 pyramidal neurons. *J Neurosci* 24:722-732.
- Fisher CM (1982) Hydrocephalus as a cause of disturbances of gait in the elderly. *Neurology* 32:1358-1363.
- Flanders KC, Ludecke G, Engels S, Cissel DS, Roberts AB, Kondaiah P, Lafyatis R, Sporn MB, Unsicker K (1991) Localization and actions of transforming growth factor-beta s in the embryonic nervous system. *Development* 113:183-191.
- Fletcher JM, Barnes M, Dennis M (2002) Language development in children with spina bifida. *Semin Pediatr Neurol* 9:201-208.
- Fletcher JM, Francis DJ, Thompson NM, Davidson KC, Miner ME (1992a) Verbal and nonverbal skill discrepancies in hydrocephalic children. *J Clin Exp Neuropsychol* 14:593-609.
- Fletcher JM, Brookshire BL, Landry SH, Bohan TP, Davidson KC, Francis DJ, Thompson NM, Miner ME (1995) Behavioral adjustment of children with hydrocephalus: relationships with etiology, neurological, and family status. *J Pediatr Psychol* 20:109-125.
- Fletcher JM, McCauley SR, Brandt ME, Bohan TP, Kramer LA, Francis DJ, Thorstad K, Brookshire BL (1996a) Regional brain tissue composition in children with hydrocephalus. Relationships with cognitive development. *Arch Neurol* 53:549-557.
- Fletcher JM, Bohan TP, Brandt ME, Brookshire BL, Beaver SR, Francis DJ, Davidson KC, Thompson NM, Miner ME (1992b) Cerebral white matter and cognition in hydrocephalic children. *Arch Neurol* 49:818-824.
- Fletcher JM, Bohan TP, Brandt ME, Kramer LA, Brookshire BL, Thorstad K, Davidson KC, Francis DJ, McCauley SR, Baumgartner JE (1996b) Morphometric evaluation of the hydrocephalic brain: relationships with cognitive development. *Childs Nerv Syst* 12:192-199.
- Flood C, Akinwunmi J, Lagord C, Daniel M, Berry M, Jackowski A, Logan A (2001) Transforming growth factor-beta1 in the cerebrospinal fluid of patients with subarachnoid hemorrhage: titers derived from exogenous and endogenous sources. *J Cereb Blood Flow Metab* 21:157-162.
- Foltz EL, Shurtleff DB (1963) Five-Year Comparative Study of Hydrocephalus in Children with and without Operation (113 Cases). *J Neurosurg* 20:1064-1079.

- Forni PE, Scuoppo C, Imayoshi I, Taulli R, Dastru W, Sala V, Betz UA, Muzzi P, Martinuzzi D, Vercelli AE, Kageyama R, Ponzetto C (2006) High levels of Cre expression in neuronal progenitors cause defects in brain development leading to microencephaly and hydrocephaly. *J Neurosci* 26:9593-9602.
- Fransen E, Van Camp G, Vits L, Willems PJ (1997) L1-associated diseases: clinical geneticists divide, molecular geneticists unite. *Hum Mol Genet* 6:1625-1632.
- Fransen E, Lemmon V, Van Camp G, Vits L, Coucke P, Willems PJ (1995) CRASH syndrome: clinical spectrum of corpus callosum hypoplasia, retardation, adducted thumbs, spastic paraparesis and hydrocephalus due to mutations in one single gene, L1. *Eur J Hum Genet* 3:273-284.
- Fransen E, D'Hooge R, Van Camp G, Verhoye M, Sijbers J, Reyniers E, Soriano P, Kamiguchi H, Willemsen R, Koekkoek SK, De Zeeuw CI, De Deyn PP, Van der Linden A, Lemmon V, Kooy RF, Willems PJ (1998) L1 knockout mice show dilated ventricles, vermis hypoplasia and impaired exploration patterns. *Hum Mol Genet* 7:999-1009.
- Fraser JJ, Fraser C (2007) Gait disorder is the cardinal sign of normal pressure hydrocephalus: a case study. *J Neurosci Nurs* 39:132-134, 192.
- Fukuhara T, Luciano MG, Kowalski RJ (2002) Clinical features of third ventriculostomy failures classified by fenestration patency. *Surg Neurol* 58:102-110.
- Fukumitsu H, Ohmiya M, Nitta A, Furukawa S, Mima T, Mori K (2000) Aberrant expression of neurotrophic factors in the ventricular progenitor cells of infant congenitally hydrocephalic rats. *Childs Nerv Syst* 16:516-521.
- Fukumizu M, Takashima S, Becker LE (1995) Neonatal posthemorrhagic hydrocephalus: neuropathologic and immunohistochemical studies. *Pediatr Neurol* 13:230-234.
- Fukumizu M, Takashima S, Becker LE (1996) Glial reaction in periventricular areas of the brainstem in fetal and neonatal posthemorrhagic hydrocephalus and congenital hydrocephalus. *Brain Dev* 18:40-45.
- Fukushima N, Yokouchi K, Kawagishi K, Ren G, Higashiyama F, Moriizumi T (2003) Proliferating cell populations in experimentally-induced hydrocephalus in developing rats. *J Clin Neurosci* 10:334-337.
- Furuta S, Ohta S, Hatakeyama T, Nakamura K, Sakaki S (1993) Recovery of protein synthesis in tolerance-induced hippocampal CA1 neurons after transient forebrain ischemia. *Acta Neuropathol* 86:329-336.
- Gabriel S, Kivi A, Kovacs R, Lehmann TN, Lanksch WR, Meencke HJ, Heinemann U (1998) Effects of barium on stimulus-induced changes in  $[K^+]_o$  and field potentials in dentate gyrus and area CA1 of human epileptic hippocampus. *Neurosci Lett* 249:91-94.

- Gadsdon DR, Variend S, Emery JL (1979) Myelination of the corpus callosum. II. The effect of relief of hydrocephalus upon the processes of myelination. *Z Kinderchir Grenzgeb* 28:314-321.
- Gadson DR, Emery JL (1976) Quantitative morphological studies of developing human cerebellar cortex in various disease states. *Arch Dis Child* 51:964-967.
- Galbreath E, Kim SJ, Park K, Brenner M, Messing A (1995) Overexpression of TGF-beta 1 in the central nervous system of transgenic mice results in hydrocephalus. *J Neuropathol Exp Neurol* 54:339-349.
- Gammal TE, Allen MB, Jr., Brooks BS, Mark EK (1987) MR evaluation of hydrocephalus. *AJR Am J Roentgenol* 149:807-813.
- Ganzler-Odenthal SI, Redies C (1998) Blocking N-cadherin function disrupts the epithelial structure of differentiating neural tissue in the embryonic chicken brain. *J Neurosci* 18:5415-5425.
- Garcia-Sevilla JA, Escriba PV, Ozaita A, La Harpe R, Walzer C, Eytan A, Guimon J (1999) Up-regulation of immunolabeled alpha2A-adrenoceptors, Gi coupling proteins, and regulatory receptor kinases in the prefrontal cortex of depressed suicides. *J Neurochem* 72:282-291.
- Gee P, Rhodes CH, Fricker LD, Angeletti RH (1993) Expression of neuropeptide processing enzymes and neurosecretory proteins in ependyma and choroid plexus epithelium. *Brain Res* 617:238-248.
- Gehlert DR, Gackenhimer SL, Schober DA (1991) Autoradiographic localization of subtypes of angiotensin II antagonist binding in the rat brain. *Neuroscience* 44:501-514.
- Gerlai R, Wojtowicz JM, Marks A, Roder J (1995) Overexpression of a calcium-binding protein, S100 beta, in astrocytes alters synaptic plasticity and impairs spatial learning in transgenic mice. *Learn Mem* 2:26-39.
- Giroud M, Dumas R (1995) Role of associated cortical lesions in motor partial seizures and lenticulostriate infarcts. *Epilepsia* 36:465-470.
- Go KG, Stokroos I, Blaauw EH, Zuiderveen F, Molenaar I (1976) Changes of ventricular ependyma and choroid plexus in experimental hydrocephalus, as observed by scanning electron microscopy. *Acta Neuropathol* 34:55-64.
- Gobron S, Monnerie H, Meiniel R, Creveaux I, Lehmann W, Lamalle D, Dastugue B, Meiniel A (1996) SCO-spondin: a new member of the thrombospondin family secreted by the subcommissural organ is a candidate in the modulation of neuronal aggregation. *J Cell Sci* 109 ( Pt 5):1053-1061.

- Gobron S, Creveaux I, Meiniel R, Didier R, Herbet A, Bamdad M, El Bitar F, Dastugue B, Meiniel A (2000) Subcommissural organ/Reissner's fiber complex: characterization of SCO-spondin, a glycoprotein with potent activity on neurite outgrowth. *Glia* 32:177-191.
- Goh D, Minns RA (1995) Intracranial pressure and cerebral arterial flow velocity indices in childhood hydrocephalus: current review. *Childs Nerv Syst* 11:392-396.
- Gonzalez-Marrero I, Carmona-Calero EM, Fernandez-Rodriguez P, Perez-Gonzalez H, Ormazabal-Ramos C, Castaneyra-Ruiz L, Perez-Garcia CG, Martinez-Pena-Valenzuela I, Castaneyra-Ruiz A, Castaneyra-Perdomo A, Ferres-Torres R (2007) Expression of certain proteins in the subfornical organ and cerebrospinal fluid of spontaneously hypertensive rats. *Histol Histopathol* 22:1371-1378.
- Gonzalez Santander R, Martinez Cuadrado G, Rubio Saez M, Bujan Varela J, Larana Sole A (1984) Ultrastructural basis of the coordination of ciliary movement. Contractile function of the interconnecting cross-striated roots in ependymal epithelium. *Acta Anat (Basel)* 118:82-90.
- Gopinath G, Bhatia R, Gopinath PG (1979) Ultrastructural observations in experimental hydrocephalus in the rabbit. *J Neurol Sci* 43:333-334.
- Graff-Radford NR, Godersky JC (1986) Normal-pressure hydrocephalus. Onset of gait abnormality before dementia predicts good surgical outcome. *Arch Neurol* 43:940-942.
- Grant JA, McLone DG (1997) Third ventriculostomy: a review. *Surg Neurol* 47:210-212.
- Green MC (1970) The developmental effects of congenital hydrocephalus (ch) in the mouse. *Dev Biol* 23:585-608.
- Greitz D (1993) Cerebrospinal fluid circulation and associated intracranial dynamics. A radiologic investigation using MR imaging and radionuclide cisternography. *Acta Radiol Suppl* 386:1-23.
- Greitz D, Hannerz J (1996) A proposed model of cerebrospinal fluid circulation: observations with radionuclide cisternography. *AJNR Am J Neuroradiol* 17:431-438.
- Greitz D, Greitz T, Hindmarsh T (1997a) We need a new understanding of the reabsorption of cerebrospinal fluid--II. *Acta Paediatr* 86:1148.
- Greitz D, Greitz T, Hindmarsh T (1997b) A new view on the CSF-circulation with the potential for pharmacological treatment of childhood hydrocephalus. *Acta Paediatr* 86:125-132.
- Gressens P, Marret S, Hill JM, Brenneman DE, Gozes I, Fridkin M, Evrard P (1997) Vasoactive intestinal peptide prevents excitotoxic cell death in the murine developing brain. *J Clin Invest* 100:390-397.

- Grondona JM, Perez-Martin M, Cifuentes M, Perez J, Jimenez AJ, Perez-Figares JM, Fernandez-Llebrez P (1996) Ependymal denudation, aqueductal obliteration and hydrocephalus after a single injection of neuraminidase into the lateral ventricle of adult rats. *J Neuropathol Exp Neurol* 55:999-1008.
- Grouzmann E, Borgeat A, Fathi M, Gaillard RC, Ravussin P (2000) Plasma and cerebrospinal fluid concentration of neuropeptide Y, serotonin, and catecholamines in patients under propofol or isoflurane anesthesia. *Can J Physiol Pharmacol* 78:100-107.
- Gruneberg H, Wickramaratne GA (1974) A re-examination of two skeletal mutants of the mouse, vestigial-tail (vt) and congenital hydrocephalus (ch). *J Embryol Exp Morphol* 31:207-222.
- Gueli U, Lato M, Cagini P (1976) [Prospects of medical treatment in communicating congenital hydrocephalus]. *Minerva Pediatr* 28:1530-1535.
- Gurguis GN, Vo SP, Griffith JM, Rush AJ (1999a) Neutrophil beta(2)-adrenoceptor function in major depression: G(s) coupling, effects of imipramine and relationship to treatment outcome. *Eur J Pharmacol* 386:135-144.
- Gurguis GN, Turkka J, Laruelle M, Kleinman J, Linnoila M (1999b) Coupling efficiency of brain beta-adrenergic receptors to Gs protein in suicide, alcoholism and control subjects. *Psychopharmacology (Berl)* 145:31-38.
- Gurguis GN, Andrews R, Antai-Otong D, Vo SP, Blakeley JE, Orsulak PJ, Rush AJ (1999c) Neutrophil beta2-adrenergic receptor coupling efficiency to Gs protein in subjects with post-traumatic stress disorder and normal controls. *Psychopharmacology (Berl)* 143:131-140.
- Gurguis GN, Blakeley JE, Antai-Otong D, Vo SP, Orsulak PJ, Petty F, Rush AJ (1999d) Adrenergic receptor function in panic disorder. II. Neutrophil beta 2 receptors: Gs protein coupling, effects of imipramine treatment and relationship to treatment outcome. *J Psychiatr Res* 33:309-322.
- Gurguis GN, Antai-Otong D, Vo SP, Blakeley JE, Orsulak PJ, Petty F, Rush AJ (1999e) Adrenergic receptor function in panic disorder. I. Platelet alpha 2 receptors: Gi protein coupling, effects of imipramine, and relationship to treatment outcome. *Neuropsychopharmacology* 20:162-176.
- Haan J, Jansen EN, Oostrom J, Roos RA (1987) Falling spells in normal pressure hydrocephalus: a favourable prognostic sign? *Eur Neurol* 27:216-220.
- Hader WJ, Drake J, Cochrane D, Sparrow O, Johnson ES, Kestle J (2002) Death after late failure of third ventriculostomy in children. Report of three cases. *J Neurosurg* 97:211-215.

- Hagberg B, Sjorgen I (1966) The chronic brain syndrome of infantile hydrocephalus. A follow-up study of 63 spontaneously arrested cases. *Am J Dis Child* 112:189-196.
- Haidri NH, Modi SM (1977) Normal pressure hydrocephalus and hypertensive cerebrovascular disease. *Dis Nerv Syst* 38:918-921.
- Hakim CA, Hakim R, Hakim S (2001) Normal-pressure hydrocephalus. *Neurosurg Clin N Am* 12:761-773, ix.
- Hakim S, Venegas JG, Burton JD (1976) The physics of the cranial cavity, hydrocephalus and normal pressure hydrocephalus: mechanical interpretation and mathematical model. *Surg Neurol* 5:187-210.
- Hale PM, McAllister JP, 2nd, Katz SD, Wright LC, Lovely TJ, Miller DW, Wolfson BJ, Salotto AG, Shroff DV (1992) Improvement of cortical morphology in infantile hydrocephalic animals after ventriculoperitoneal shunt placement. *Neurosurgery* 31:1085-1096; discussion 1096.
- Halliwell B (1992) Reactive oxygen species and the central nervous system. *J Neurochem* 59:1609-1623.
- Hamano T, Yoshimura M, Yamazaki T, Shinkai Y, Yanagisawa K, Kuriyama M, Ihara Y (1997) Amyloid beta-protein (A beta) accumulation in the leptomeninges during aging and in Alzheimer disease. *J Neuropathol Exp Neurol* 56:922-932.
- Hamer A, Sahar A (1978) Absence of formation of cerebrospinal fluid in the spinal subarachnoid space of the cat. *Isr J Med Sci* 14:741-744.
- Handel M, Schulz S, Stanarius A, Schreff M, Erdtmann-Vourliotis M, Schmidt H, Wolf G, Holtt V (1999) Selective targeting of somatostatin receptor 3 to neuronal cilia. *Neuroscience* 89:909-926.
- Hannibal J, Jamen F, Nielsen HS, Journot L, Brabet P, Fahrenkrug J (2001) Dissociation between light-induced phase shift of the circadian rhythm and clock gene expression in mice lacking the pituitary adenylate cyclase activating polypeptide type 1 receptor. *J Neurosci* 21:4883-4890.
- Hansen A, Whitelaw A, Lapp C, Brugnara C (1997a) Cerebrospinal fluid plasminogen activator inhibitor-1: a prognostic factor in posthaemorrhagic hydrocephalus. *Acta Paediatr* 86:995-998.
- Hansen AR, Volpe JJ, Goumnerova LC, Madsen JR (1997b) Intraventricular urokinase for the treatment of posthemorrhagic hydrocephalus. *Pediatr Neurol* 17:213-217.
- Hansson E, Ronnback L (2003) Glial neuronal signaling in the central nervous system. *Faseb J* 17:341-348.

- Harris NG, Plant HD, Briggs RW, Jones HC (1996a) Metabolite changes in the cerebral cortex of treated and untreated infant hydrocephalic rats studied using in vitro <sup>31</sup>P-NMR spectroscopy. *J Neurochem* 67:2030-2038.
- Harris NG, McAllister JP, 2nd, Conaughty JM, Jones HC (1996b) The effect of inherited hydrocephalus and shunt treatment on cortical pyramidal cell dendrites in the infant H-Tx rat. *Exp Neurol* 141:269-279.
- Harris NG, Plant HD, Inglis BA, Briggs RW, Jones HC (1997) Neurochemical changes in the cerebral cortex of treated and untreated hydrocephalic rat pups quantified with in vitro <sup>1</sup>H-NMR spectroscopy. *J Neurochem* 68:305-312.
- Harrison MJ, Robert CM, Uttley D (1974) Benign aqueduct stenosis in adults. *J Neurol Neurosurg Psychiatry* 37:1322-1328.
- Hassin GB (1948) Cerebrospinal fluid, its origin, nature and function. *J Neuropathol Exp Neurol* 7:172-181.
- Hatskelzon L, Dalal BI, Shalev A, Robertson C, Gerrard JM (1993) Wide distribution of granulophysin epitopes in granules of human tissues. *Lab Invest* 68:509-519.
- Hawkins D, Bowers TM, Bannister CM, Miyan JA (1997) The functional outcome of shunting H-Tx rat pups at different ages. *Eur J Pediatr Surg* 7 Suppl 1:31-34.
- Hebb AO, Cusimano MD (2001) Idiopathic normal pressure hydrocephalus: a systematic review of diagnosis and outcome. *Neurosurgery* 49:1166-1184; discussion 1184-1166.
- Herndon RM, Johnson RT, Davis LE, Descalzi LR (1974) Ependymitis in mumps virus meningitis. Electron microscopical studies of cerebrospinal fluid. *Arch Neurol* 30:475-479.
- Hidaka M, Matsumae M, Yamamura M, Tsugane R, Sato O (1997) Glucose metabolism and protective biochemical mechanisms in a rat brain affected by kaolin-induced hydrocephalus. *Childs Nerv Syst* 13:183-188.
- Higashi K, Asahisa H, Ueda N, Kobayashi K, Hara K, Noda Y (1986) Cerebral blood flow and metabolism in experimental hydrocephalus. *Neurol Res* 8:169-176.
- Hochwald GM, Sahar A, Sadik AR, Ransohoff J (1969) Cerebrospinal fluid production and histological observations in animals with experimental obstructive hydrocephalus. *Exp Neurol* 25:190-199.
- Hoffman HJ, Smith MS (1986) The use of shunting devices for cerebrospinal fluid in Canada. *Can J Neurol Sci* 13:81-87.

- Hoglund M, Tisell M, Wikkelso C (2001) [Incidence of surgery for hydrocephalus in adults surveyed: same number afflicted by hydrocephalus as by multiple sclerosis]. *Lakartidningen* 98:1681-1685.
- Holtzman SG (2000) Further characterization of the discriminative stimulus effects of spiradoline. *Pharmacol Biochem Behav* 66:517-522.
- Hong HK, Chakravarti A, Takahashi JS (2004) The gene for soluble N-ethylmaleimide sensitive factor attachment protein alpha is mutated in hydrocephaly with hop gait (hyh) mice. *Proc Natl Acad Sci U S A* 101:1748-1753.
- Hopf NJ, Grunert P, Fries G, Resch KD, Perneczky A (1999) Endoscopic third ventriculostomy: outcome analysis of 100 consecutive procedures. *Neurosurgery* 44:795-804; discussion 804-796.
- Hoppe-Hirsch E, Laroussinie F, Brunet L, Sainte-Rose C, Renier D, Cinalli G, Zerah M, Pierre-Kahn A (1998) Late outcome of the surgical treatment of hydrocephalus. *Childs Nerv Syst* 14:97-99.
- Horan P, Taylor J, Yamamura HI, Porreca F (1992) Extremely long-lasting antagonistic actions of nor-binaltorphimine (nor- BNI) in the mouse tail-flick test. *J Pharmacol Exp Ther* 260:1237-1243.
- Horinek D, Cihar M, Tichy M (2003) Current methods in the treatment of posthemorrhagic hydrocephalus in infants. *Bratisl Lek Listy* 104:347-351.
- Hortsch M (1996) The L1 family of neural cell adhesion molecules: old proteins performing new tricks. *Neuron* 17:587-593.
- Hough LB, Nalwalk JW, Chen Y, Schuller A, Zhu Y, Zhang J, Menge WM, Leurs R, Timmerman H, Pintar JE (2000) Imiprogan, a cimetidine analog, induces morphine-like antinociception in opioid receptor-knockout mice. *Brain Res* 880:102-108.
- Huang Y, Roelink H, McKnight GS (2002) Protein kinase A deficiency causes axially localized neural tube defects in mice. *J Biol Chem* 277:19889-19896.
- Ibanez-Tallon I, Gorokhova S, Heintz N (2002) Loss of function of axonemal dynein Mdnah5 causes primary ciliary dyskinesia and hydrocephalus. *Hum Mol Genet* 11:715-721.
- Ibanez-Tallon I, Pagenstecher A, Fliegauf M, Olbrich H, Kispert A, Ketelsen UP, North A, Heintz N, Omran H (2004) Dysfunction of axonemal dynein heavy chain Mdnah5 inhibits ependymal flow and reveals a novel mechanism for hydrocephalus formation. *Hum Mol Genet* 13:2133-2141.
- Iddon JL, Pickard JD, Cross JJ, Griffiths PD, Czosnyka M, Sahakian BJ (1999) Specific patterns of cognitive impairment in patients with idiopathic normal pressure



- hydrocephalus and Alzheimer's disease: a pilot study. *J Neurol Neurosurg Psychiatry* 67:723-732.
- Ikedo T, Xia XY, Xia YX, Ikenoue T, Choi BH (1999) Expression of glial cell line-derived neurotrophic factor in the brain and cerebrospinal fluid of the developing rat. *Int J Dev Neurosci* 17:681-691.
- Irigoin C, Rodriguez EM, Heinrichs M, Frese K, Herzog S, Oksche A, Rott R (1990) Immunocytochemical study of the subcommissural organ of rats with induced postnatal hydrocephalus. *Exp Brain Res* 82:384-392.
- Itoh K, Cheng L, Kamei Y, Fushiki S, Kamiguchi H, Gutwein P, Stoeck A, Arnold B, Altevogt P, Lemmon V (2004) Brain development in mice lacking L1-L1 homophilic adhesion. *J Cell Biol* 165:145-154.
- Jabourian Z, Lublin FD, Adler A, Gonzales C, Northrup B, Zwillenberg D (1986) Hydrocephalus in Kartagener's syndrome. *Ear Nose Throat J* 65:468-472.
- James AE, Jr., Flor WJ, Novak GR, Ribas JL, Parker JL, Sickel WL (1980) The ultrastructural basis of periventricular edema: preliminary studies. *Radiology* 135:747-750.
- Jan LY, Jan YN (1997) Receptor-regulated ion channels. *Curr Opin Cell Biol* 9:155-160.
- Jimenez AJ, Tome M, Paez P, Wagner C, Rodriguez S, Fernandez-Llebrez P, Rodriguez EM, Perez-Figares JM (2001) A programmed ependymal denudation precedes congenital hydrocephalus in the *hyh* mutant mouse. *J Neuropathol Exp Neurol* 60:1105-1119.
- Johanson C (2003) The choroid-plexus-CSF nexus: gateway to the brain. In: *Neuroscience in Medicine* (Conn PM, ed), pp 165-195. Totowa, NJ: Humana Press Inc.
- Johanson C, Del Bigio M, Kinsman S, Miyan J, Pattisapu J, Robinson M, Jones HC (2001) New models for analysing hydrocephalus and disorders of CSF volume transmission. *Br J Neurosurg* 15:281-283.
- Johanson CE, Jones HC (2001) Promising vistas in hydrocephalus and cerebrospinal fluid research. *Trends Neurosci* 24:631-632.
- Johanson CE, Preston JE, Chodobski A, Stopa EG, Szmydynger-Chodobska J, McMillan PN (1999a) AVP V1 receptor-mediated decrease in Cl<sup>-</sup> efflux and increase in dark cell number in choroid plexus epithelium. *Am J Physiol* 276:C82-90.
- Johanson CE, Szmydynger-Chodobska J, Chodobski A, Baird A, McMillan P, Stopa EG (1999b) Altered formation and bulk absorption of cerebrospinal fluid in FGF-2-induced hydrocephalus. *Am J Physiol* 277:R263-271.

- Johanson CE, Duncan JA, 3rd, Klinge PM, Brinker T, Stopa EG, Silverberg GD (2008) Multiplicity of cerebrospinal fluid functions: New challenges in health and disease. *Cerebrospinal Fluid Res* 5:10.
- Johnson G, Brane D, Block W, van Kammen DP, Gurklis J, Peters JL, Wyatt RJ, Kirch DG, Ghanbari HA, Merrill CR (1992a) Cerebrospinal fluid protein variations in common to Alzheimer's disease and schizophrenia. *Appl Theor Electrophor* 3:47-53.
- Johnson MD, Gold LI, Moses HL (1992b) Evidence for transforming growth factor-beta expression in human leptomeningeal cells and transforming growth factor-beta-like activity in human cerebrospinal fluid. *Lab Invest* 67:360-368.
- Johnson MJ, Ayzman I, Wood AS, Tkach JA, Klauschie J, Skarupa DJ, McAllister JP, Luciano MG (1999) Development and characterization of an adult model of obstructive hydrocephalus. *J Neurosci Methods* 91:55-65.
- Johnson RT, Johnson KP (1968) Hydrocephalus as a sequela of experimental mumps virus infection in the hamster. *J Neuropathol Exp Neurol* 27:136-137.
- Johnson RT, Johnson KP (1968) Hydrocephalus following viral infection: the pathology of aqueductal stenosis developing after experimental mumps virus infection. *J Neuropathol Exp Neurol* 27:591-606.
- Johnston I, Teo C (2000) Disorders of CSF hydrodynamics. *Childs Nerv Syst* 16:776-799.
- Johnston M, Zakharov A, Koh L, Armstrong D (2005) Subarachnoid injection of Microfil reveals connections between cerebrospinal fluid and nasal lymphatics in the non-human primate. *Neuropathol Appl Neurobiol* 31:632-640.
- Johnston M, Zakharov A, Papaiconomou C, Salmasi G, Armstrong D (2004) Evidence of connections between cerebrospinal fluid and nasal lymphatic vessels in humans, non-human primates and other mammalian species. *Cerebrospinal Fluid Res* 1:2.
- Jones DN, Holtzman SG (1992) Long term kappa-opioid receptor blockade following nor-binaltorphimine. *Eur J Pharmacol* 215:345-348.
- Jones HC (1985) Cerebrospinal fluid pressure and resistance to absorption during development in normal and hydrocephalic mutant mice. *Exp Neurol* 90:162-172.
- Jones HC, Sellars RA (1982) The movement of fluid out of the cerebral ventricles in fetal and neonatal rats. *Zeitschrift fur Kinderchirurgie* 37:130-133.
- Jones HC, Bucknall RM (1988) Inherited prenatal hydrocephalus in the H-Tx rat: a morphological study. *Neuropathol Appl Neurobiol* 14:263-274.
- Jones HC, Andersohn RW (1998) Progressive changes in cortical water and electrolyte content at three stages of rat infantile hydrocephalus and the effect of shunt treatment. *Exp Neurol* 154:126-136.

- Jones HC, Dack S, Ellis C (1987) Morphological aspects of the development of hydrocephalus in a mouse mutant (SUMS/NP). *Acta Neuropathol* 72:268-276.
- Jones HC, Bucknall RM, Harris NG (1991) The cerebral cortex in congenital hydrocephalus in the H-Tx rat: a quantitative light microscopy study. *Acta Neuropathol* 82:217-224.
- Jones HC, Rivera KM, Harris NG (1995a) Learning deficits in congenitally hydrocephalic rats and prevention by early shunt treatment. *Childs Nerv Syst* 11:655-660.
- Jones HC, Harris NG, Briggs RW, Williams SC (1995b) Shunt treatment at two postnatal ages in hydrocephalic H-Tx rats quantified using MR imaging. *Exp Neurol* 133:144-152.
- Jones HC, Harris NG, Rocca JR, Andersohn RW (1997) Progressive changes in cortical metabolites at three stages of infantile hydrocephalus studied by in vitro NMR spectroscopy. *J Neurotrauma* 14:587-602.
- Jones HC, Carter BJ, Depelteau JS, Roman M, Morel L (2001a) Chromosomal linkage associated with disease severity in the hydrocephalic H-Tx rat. *Behav Genet* 31:101-111.
- Jones HC, Depelteau JS, Carter BJ, Lopman BA, Morel L (2001b) Genome-wide linkage analysis of inherited hydrocephalus in the H-Tx rat. *Mamm Genome* 12:22-26.
- Jones HC, Lopman BA, Jones TW, Carter BJ, Depelteau JS, Morel L (2000) The expression of inherited hydrocephalus in H-Tx rats. *Childs Nerv Syst* 16:578-584.
- Joseph JA, Cutler R, Roth GS (1993) Changes in G protein-mediated signal transduction in aging and Alzheimer's disease. *Ann N Y Acad Sci* 695:42-45.
- Jung JS, Bhat RV, Preston GM, Guggino WB, Baraban JM, Agre P (1994) Molecular characterization of an aquaporin cDNA from brain: candidate osmoreceptor and regulator of water balance. *Proc Natl Acad Sci U S A* 91:13052-13056.
- Kang JK, Lee IW (1999) Long-term follow-up of shunting therapy. *Childs Nerv Syst* 15:711-717.
- Kasaian MT, Neet KE (1989) Nerve growth factor in human amniotic and cerebrospinal fluid. *Biofactors* 2:99-104.
- Kasper M, Perry G, Stosiek P (1991) Cytokeratin expression in human spinal meninges and ependymal cells. *J Hirnforsch* 32:19-25.
- Katayama K (1992) *Hydrocephalus: Pathogenesis and Treatment*: Springer.
- Kawamata T, Katayama Y, Tsuji N, Nishimoto H (2003) Metabolic derangements in interstitial brain edema with preserved blood flow: selective vulnerability of the hippocampal CA3 region in rat hydrocephalus. *Acta Neurochir Suppl* 86:545-547.

- Kazan S, Gura A, Ucar T, Korkmaz E, Ongun H, Akyuz M (2005) Hydrocephalus after intraventricular hemorrhage in preterm and low-birth weight infants: analysis of associated risk factors for ventriculoperitoneal shunting. *Surg Neurol* 64 Suppl 2:S77-81; discussion S81.
- Kenakin T (2004) Efficacy as a Vector: the Relative Prevalence and Paucity of Inverse Agonism. *Mol Pharmacol* 65:2-11.
- Kennedy CR, Ayers S, Campbell MJ, Elbourne D, Hope P, Johnson A (2001) Randomized, controlled trial of acetazolamide and furosemide in posthemorrhagic ventricular dilation in infancy: follow-up at 1 year. *Pediatrics* 108:597-607.
- Kenwrick S, Jouet M, Donnai D (1996) X linked hydrocephalus and MASA syndrome. *J Med Genet* 33:59-65.
- Keyser DO, Pellmar TC (1994) Synaptic transmission in the hippocampus: critical role for glial cells. *Glia* 10:237-243.
- Khajeh MR, Bannister CM, Miyan JA (2002) Study of the motor corticospinal system in the developing rat fetus: comparison of Wistar and normal and hydrocephalic HTx rats. *Eur J Pediatr Surg* 12 Suppl 1:S3-5.
- Khan OH, Enno T, Del Bigio MR (2003) Magnesium sulfate therapy is of mild benefit to young rats with kaolin-induced hydrocephalus. *Pediatr Res* 53:970-976.
- Khan OH, Enno TL, Del Bigio MR (2006) Brain damage in neonatal rats following kaolin induction of hydrocephalus. *Exp Neurol* 200:311-320.
- Khan ZU, Koulen P, Rubinstein M, Grandy DK, Goldman-Rakic PS (2001) An astroglia-linked dopamine D2-receptor action in prefrontal cortex. *Proc Natl Acad Sci U S A* 98:1964-1969.
- Kiefer M, Eymann R, von Tiling S, Muller A, Steudel WI, Booz KH (1998) The ependyma in chronic hydrocephalus. *Childs Nerv Syst* 14:263-270.
- Kielinen M, Rantala H, Timonen E, Linna SL, Moilanen I (2004) Associated medical disorders and disabilities in children with autistic disorder: a population-based study. *Autism* 8:49-60.
- Kim JC, Badano JL, Sibold S, Esmail MA, Hill J, Hoskins BE, Leitch CC, Venner K, Ansley SJ, Ross AJ, Leroux MR, Katsanis N, Beales PL (2004) The Bardet-Biedl protein BBS4 targets cargo to the pericentriolar region and is required for microtubule anchoring and cell cycle progression. *Nat Genet* 36:462-470.
- Kirsch J, Betz H (1998) Glycine-receptor activation is required for receptor clustering in spinal neurons. *Nature* 392:717-720.

- Kitazawa K, Tada T (1994) Elevation of transforming growth factor-beta 1 level in cerebrospinal fluid of patients with communicating hydrocephalus after subarachnoid hemorrhage. *Stroke* 25:1400-1404.
- Ko MC, Butelman ER, Woods JH (1999) Activation of peripheral kappa opioid receptors inhibits capsaicin-induced thermal nociception in rhesus monkeys. *J Pharmacol Exp Ther* 289:378-385.
- Ko MC, Willmont KJ, Lee H, Flory GS, Woods JH (2003) Ultra-long antagonism of kappa opioid agonist-induced diuresis by intracisternal nor-binaltorphimine in monkeys. *Brain Res* 982:38-44.
- Kobayashi Y, Watanabe M, Okada Y, Sawa H, Takai H, Nakanishi M, Kawase Y, Suzuki H, Nagashima K, Ikeda K, Motoyama N (2002) Hydrocephalus, situs inversus, chronic sinusitis, and male infertility in DNA polymerase lambda-deficient mice: possible implication for the pathogenesis of immotile cilia syndrome. *Mol Cell Biol* 22:2769-2776.
- Koh L, Zakharov A, Johnston M (2005) Integration of the subarachnoid space and lymphatics: is it time to embrace a new concept of cerebrospinal fluid absorption? *Cerebrospinal Fluid Res* 2:6.
- Kondziella D, Ludemann W, Brinker T, Sletvold O, Sonnewald U (2002) Alterations in brain metabolism, CNS morphology and CSF dynamics in adult rats with kaolin-induced hydrocephalus. *Brain Res* 927:35-41.
- Kondziella D, Qu H, Ludemann W, Brinker T, Sletvold O, Sonnewald U (2003) Astrocyte metabolism is disturbed in the early development of experimental hydrocephalus. *J Neurochem* 85:274-281.
- Korhonen L, Riikonen R, Nawa H, Lindholm D (1998) Brain derived neurotrophic factor is increased in cerebrospinal fluid of children suffering from asphyxia. *Neurosci Lett* 240:151-154.
- Kozminski KG, Johnson KA, Forscher P, Rosenbaum JL (1993) A motility in the eukaryotic flagellum unrelated to flagellar beating. *Proc Natl Acad Sci U S A* 90:5519-5523.
- Krauss JK, Regel JP, Vach W, Droste DW, Borremans JJ, Mergner T (1996) Vascular risk factors and arteriosclerotic disease in idiopathic normal-pressure hydrocephalus of the elderly. *Stroke* 27:24-29.
- Krebs DL, Hilton DJ (2000) SOCS: physiological suppressors of cytokine signaling. *J Cell Sci* 113 ( Pt 16):2813-2819.
- Krebs DL, Metcalf D, Merson TD, Voss AK, Thomas T, Zhang JG, Rakar S, O'Bryan M K, Willson TA, Viney EM, Mielke LA, Nicola NA, Hilton DJ, Alexander WS (2004) Development of hydrocephalus in mice lacking SOCS7. *Proc Natl Acad Sci U S A* 101:15446-15451.

- Kriebel RM, McAllister JP, 2nd (2000) Pathology of the hippocampus in experimental feline infantile hydrocephalus. *Neurol Res* 22:29-36.
- Kriebel RM, Shah AB, McAllister JP, 2nd (1993) The microstructure of cortical neuropil before and after decompression in experimental infantile hydrocephalus. *Exp Neurol* 119:89-98.
- Kuba H, Inamura T, Ikezaki K, Inoha S, Nakamizo A, Shono T, Fukui K, Fukui M (2002) Gait disturbance in patients with low pressure hydrocephalus. *J Clin Neurosci* 9:33-36.
- Kume T, Deng KY, Winfrey V, Gould DB, Walter MA, Hogan BL (1998) The forkhead/winged helix gene *Mf1* is disrupted in the pleiotropic mouse mutation congenital hydrocephalus. *Cell* 93:985-996.
- Laezza C, Wolff J, Bifulco M (1997) Identification of a 48-kDa prenylated protein that associates with microtubules as 2',3'-cyclic nucleotide 3'-phosphodiesterase in FRTL-5 cells. *FEBS Lett* 413:260-264.
- Lagace-Simard J, Descoteaux JP, Lussier G (1982) Experimental pneumovirus infections. 2. Hydrocephalus of hamsters and mice due to infection with human respiratory syncytial virus (RS). *Am J Pathol* 107:36-40.
- Lampson LA, Grabowska A, Whelan JP (1994) Class I and II MHC expression and its implications for regeneration in the nervous system. *Prog Brain Res* 103:307-317.
- Lang B, Song B, Davidson W, MacKenzie A, Smith N, McCaig CD, Harmar AJ, Shen S (2006) Expression of the human PAC1 receptor leads to dose-dependent hydrocephalus-related abnormalities in mice. *J Clin Invest* 116:1924-1934.
- Lappe-Siefke C, Goebbels S, Gravel M, Nicksch E, Lee J, Braun PE, Griffiths IR, Nave KA (2003) Disruption of *Cnp1* uncouples oligodendroglial functions in axonal support and myelination. *Nat Genet* 33:366-374.
- Laurence KM, Coates S (1962) The natural history of hydrocephalus. Detailed analysis of 182 unoperated cases. *Arch Dis Child* 37:345-362.
- Lawson RF, Raimondi AJ (1973) Hydrocephalus-3, a murine mutant: I. Alterations in fine structure of choroid plexus and ependyma. *Surg Neurol* 1:115-128.
- Lazareff JA, Peacock W, Holly L, Ver Halen J, Wong A, Olmstead C (1998) Multiple shunt failures: an analysis of relevant factors. *Childs Nerv Syst* 14:271-275.
- Lee HJ, Balasubramanian SV, Murer H, Biber J, Morris ME (1999) Modulation of sulfate renal transport by alterations in cell membrane fluidity. *J Pharm Sci* 88:976-980.
- Leech RW, Payne GG, Jr. (1991) Neural tube defects: epidemiology. *J Child Neurol* 6:286-287.

- Lefkowitz RJ, Cotecchia S, Samama P, Costa T (1993) Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins. *Trends Pharmacol Sci* 14:303-307.
- Leurs R, Pena MSR, Bakker RA, Alewijnse AE, Timmerman H, Ugo Gulini MGWQaGM (2000) Constitutive activity of G protein coupled receptors and drug action. In: *Pharmacochemistry Library*, pp 327-331: Elsevier.
- Lindquist B, Carlsson G, Persson EK, Uvebrant P (2005) Learning disabilities in a population-based group of children with hydrocephalus. *Acta Paediatr* 94:878-883.
- Lindquist B, Carlsson G, Persson EK, Uvebrant P (2006) Behavioural problems and autism in children with hydrocephalus : a population-based study. *Eur Child Adolesc Psychiatry* 15:214-219.
- Lindquist B, Persson EK, Uvebrant P, Carlsson G (2008) Learning, memory and executive functions in children with hydrocephalus. *Acta Paediatr* 97:596-601.
- Lindqvist G, Andersson H, Bilting M, Blomstrand C, Malmgren H, Wikkelso C (1993) Normal pressure hydrocephalus: psychiatric findings before and after shunt operation classified in a new diagnostic system for organic psychiatry. *Acta Psychiatr Scand Suppl* 373:18-32.
- Lippa CF, Flanders KC, Kim ES, Croul S (1998) TGF-beta receptors-I and -II immunoexpression in Alzheimer's disease: a comparison with aging and progressive supranuclear palsy. *Neurobiol Aging* 19:527-533.
- Liszcza TM, Black PM, Tzouras A, Foley L, Zervas NT (1984) Morphological changes of the basilar artery, ventricles, and choroid plexus after experimental SAH. *J Neurosurg* 61:486-493.
- Lopponen T, Saukkonen AL, Serlo W, Tapanainen P, Ruokonen A, Knip M (1996) Accelerated pubertal development in patients with shunted hydrocephalus. *Arch Dis Child* 74:490-496.
- Lorber J (1975) Isosorbide in treatment of infantile hydrocephalus. *Arch Dis Child* 50:431-436.
- Lorber J, Salfield S, Lonton T (1983) Isosorbide in the management of infantile hydrocephalus. *Dev Med Child Neurol* 25:502-511.
- Louvi A, Wassef M (2000) Ectopic engrailed 1 expression in the dorsal midline causes cell death, abnormal differentiation of circumventricular organs and errors in axonal pathfinding. *Development* 127:4061-4071.
- Luciano MG, Skarupa DJ, Booth AM, Wood AS, Brant CL, Gdowski MJ (2001) Cerebrovascular adaptation in chronic hydrocephalus. *J Cereb Blood Flow Metab* 21:285-294.

- Luttrell LM (2005) Composition and function of G protein-coupled receptor signalsomes controlling mitogen-activated protein kinase activity. *J Mol Neurosci* 26:253-264.
- Lux WE, Jr., Fenstermacher JD (1975) Cerebrospinal fluid formation in ventricles and spinal subarachnoid space of the rhesus monkey. *J Neurosurg* 42:674-678.
- Lux WE, Jr., Hochwald GM, Sahar A, Ransohoff J (1970) Periventricular water content. Effect of pressure in experimental chronic hydrocephalus. *Arch Neurol* 23:475-479.
- Mahjoub MR, Montpetit B, Zhao L, Finst RJ, Goh B, Kim AC, Quarmby LM (2002) The FA2 gene of *Chlamydomonas* encodes a NIMA family kinase with roles in cell cycle progression and microtubule severing during deflagellation. *J Cell Sci* 115:1759-1768.
- Mandell JW, VandenBerg SR (1999) ERK/MAP kinase is chronically activated in human reactive astrocytes. *Neuroreport* 10:3567-3572.
- Mangano FT, McAllister JP, 2nd, Jones HC, Johnson MJ, Kriebel RM (1998) The microglial response to progressive hydrocephalus in a model of inherited aqueductal stenosis. *Neurol Res* 20:697-704.
- Mann JD, Butler AB, Johnson RN, Bass NH (1979) Clearance of macromolecular and particulate substances from the cerebrospinal fluid system of the rat. *J Neurosurg* 50:343-348.
- Margolis G, Kilham L (1969) Hydrocephalus in hamsters, ferrets, rats, and mice following inoculations with reovirus type I. II. Pathologic studies. *Lab Invest* 21:189-198.
- Margolis G, Kilham L (1977) Induction of congenital hydrocephalus in hamsters with parainfluenza type 2 virus. *Exp Mol Pathol* 27:235-248.
- Marin-Padilla M (1998) Cajal-Retzius cells and the development of the neocortex. *Trends Neurosci* 21:64-71.
- Marinov MB, Harbaugh KS, Hoopes PJ, Pikus HJ, Harbaugh RE (1996) Neuroprotective effects of preischemia intraarterial magnesium sulfate in reversible focal cerebral ischemia. *J Neurosurg* 85:117-124.
- Marszalek JR, Ruiz-Lozano P, Roberts E, Chien KR, Goldstein LS (1999) Situs inversus and embryonic ciliary morphogenesis defects in mouse mutants lacking the KIF3A subunit of kinesin-II. *Proc Natl Acad Sci U S A* 96:5043-5048.
- Martens N, Wery M, Wang P, Braet F, Gertler A, Hooghe R, Vandenhoute J, Hooghe-Peters EL (2004) The suppressor of cytokine signaling (SOCS)-7 interacts with the actin cytoskeleton through vinexin. *Exp Cell Res* 298:239-248.



- Martinez-Lage JF, Perez-Espejo MA, Almagro MJ, Ros de San Pedro J, Lopez F, Piqueras C, Tortosa J (2005) [Syndromes of overdrainage of ventricular shunting in childhood hydrocephalus]. *Neurocirugia (Astur)* 16:124-133.
- Martinez-Pena y Valenzuela I, Carmona-Calero EM, Perez-Gonzalez H, Ormazabal-Ramos C, Fernandez-Rodriguez P, Gonzalez-Marrero I, Castaneyra-Perdomo A, Ferres-Torres R (2006) Alterations of the cerebrospinal fluid proteins and subcommissural organ secretion in the arterial hypertension and ventricular dilatation. A study in SHR rats. *Histol Histopathol* 21:179-185.
- Mashayekhi F, Draper CE, Bannister CM, Pourghasem M, Owen-Lynch PJ, Miyan JA (2002) Deficient cortical development in the hydrocephalic Texas (H-Tx) rat: a role for CSF. *Brain* 125:1859-1874.
- Massicotte EM, Buist R, Del Bigio MR (2000) Altered diffusion and perfusion in hydrocephalic rat brain: a magnetic resonance imaging analysis. *J Neurosurg* 92:442-447.
- Masters JC, O'Grady M (1992) Normal pressure hydrocephalus. A potentially reversible form of dementia. *J Psychosoc Nurs Ment Health Serv* 30:25-28.
- Matsas R, Kenny AJ, Turner AJ (1986) An immunohistochemical study of endopeptidase-24.11 ("enkephalinase") in the pig nervous system. *Neuroscience* 18:991-1012.
- Matuoka K, Miki H, Takahashi K, Takenawa T (1997) A novel ligand for an SH3 domain of the adaptor protein Nck bears an SH2 domain and nuclear signaling motifs. *Biochem Biophys Res Commun* 239:488-492.
- Maxwell WL, Irvine A, Graham, Adams JH, Gennarelli TA, Tipperman R, Sturatis M (1991) Focal axonal injury: the early axonal response to stretch. *J Neurocytol* 20:157-164.
- McAllister JP, 2nd, Chovan P (1998) Neonatal hydrocephalus. Mechanisms and consequences. *Neurosurg Clin N Am* 9:73-93.
- McAllister JP, 2nd, Maugans TA, Shah MV, Truex RC, Jr. (1985) Neuronal effects of experimentally induced hydrocephalus in newborn rats. *J Neurosurg* 63:776-783.
- McAllister JP, 2nd, Wood AS, Johnson MJ, Connelly RW, Skarupa DJ, Secic M, Luciano MG, Harris NG, Jones HC (1999) Decreased c-fos expression in experimental neonatal hydrocephalus: evidence for reduced neuronal activation. *Neurosurg Focus* 7:e13.
- McCallum JE, Turbeville D (1994) Cost and outcome in a series of shunted premature infants with intraventricular hemorrhage. *Pediatr Neurosurg* 20:63-67.
- McComb J (1997) Cerebrospinal fluid, hydrocephalus, and cerebral edema. In: *Textbook of Neuropathology*, 3rd Edition (Robertson RDaD, ed), pp 225-251. Baltimore: Williams and Wilkins.

- McComb JG (1983) Recent research into the nature of cerebrospinal fluid formation and absorption. *J Neurosurg* 59:369-383.
- McComb JG (1992) Cerebrospinal fluid physiology of the developing fetus. *AJNR Am J Neuroradiol* 13:595-599.
- McGirt MJ, Woodworth G, Coon AL, Thomas G, Williams MA, Rigamonti D (2008) Diagnosis, treatment, and analysis of long-term outcomes in idiopathic normal-pressure hydrocephalus. *Neurosurgery* 62 Suppl 2:670-677.
- McLone DG (2004) The anatomy of the ventricular system. *Neurosurg Clin N Am* 15:33-38.
- Meinzel A (2001) SCO-spondin, a glycoprotein of the subcommissural organ/Reissner's fiber complex: evidence of a potent activity on neuronal development in primary cell cultures. *Microsc Res Tech* 52:484-495.
- Meinkoth JH, Ewing SA, Cowell RL, Dawson JE, Warner CK, Mathew JS, Bowles M, Thiessen AE, Panciera RJ, Fox C (1998) Morphologic and molecular evidence of a dual species ehrlichial infection in a dog presenting with inflammatory central nervous system disease. *J Vet Intern Med* 12:389-393.
- Meller ST, Dennis BJ (1993) A scanning and transmission electron microscopic analysis of the cerebral aqueduct in the rabbit. *Anat Rec* 237:124-140.
- Messer WS, Jr. (2004) Bivalent ligands for G protein-coupled receptors. *Curr Pharm Des* 10:2015-2020.
- Mestres P, Rascher K (1994) The ventricular system of the pigeon brain: a scanning electron microscope study. *J Anat* 184 ( Pt 1):35-58.
- Mignone JL, Kukekov V, Chiang AS, Steindler D, Enikolopov G (2004) Neural stem and progenitor cells in nestin-GFP transgenic mice. *J Comp Neurol* 469:311-324.
- Milhorat TH (1969) Choroid plexus and cerebrospinal fluid production. *Science* 166:1514-1516.
- Milhorat TH (1974) Failure of choroid plexectomy as treatment for hydrocephalus. *Surg Gynecol Obstet* 139:505-508.
- Milhorat TH (1987) Acute hydrocephalus after aneurysmal subarachnoid hemorrhage. *Neurosurgery* 20:15-20.
- Milhorat TH, Hammock MK, Fenstermacher JD, Levin VA (1971) Cerebrospinal fluid production by the choroid plexus and brain. *Science* 173:330-332.
- Milhorat TH, Hammock MK, Chien T, Davis DA (1976) Normal rate of cerebrospinal fluid formation five years after bilateral choroid plexectomy. Case report. *J Neurosurg* 44:735-739.

- Miller AD, Leslie RA (1994) The area postrema and vomiting. *Front Neuroendocrinol* 15:301-320.
- Miller JM, Kumar R, McAllister JP, 2nd, Krause GS (2006) Gene expression analysis of the development of congenital hydrocephalus in the H-Tx rat. *Brain Res* 1075:36-47.
- Minamikawa J, Kikuchi H, Ishikawa M, Yamamura K, Kanashiro M (1994) The effects of atrial natriuretic peptide on brain edema, intracranial pressure and cerebral energy metabolism in rat congenital hydrocephalus. *Acta Neurochir Suppl (Wien)* 60:104-106.
- Miranda E, Almonacid JA, Rodriguez S, Perez J, Hein S, Cifuentes M, Fernandez-Llebrez P, Rodriguez EM (2001) Searching for specific binding sites of the secretory glycoproteins of the subcommissural organ. *Microsc Res Tech* 52:541-551.
- Mittendorf R, Dambrosia J, Pryde PG, Lee KS, Gianopoulos JG, Besinger RE, Tomich PG (2002) Association between the use of antenatal magnesium sulfate in preterm labor and adverse health outcomes in infants. *Am J Obstet Gynecol* 186:1111-1118.
- Miwa S, Inagaki C, Fujiwara M, Takaori S (1982) The activities of noradrenergic and dopaminergic neuron systems in experimental hydrocephalus. *J Neurosurg* 57:67-73.
- Miyagami M, Nakamura S, Mukakami T, Koga N, Moriyasu N (1976) Electron microscopic study of ventricular wall and choroid plexus in experimentally-induced hydrocephalic dogs. *Neurol Med Chir (Tokyo)* 16:15-21.
- Miyake H, Eghwudjakpor PO, Sakamoto T, Mori K (1992) Catecholamine alterations in experimental hydrocephalus. *Childs Nerv Syst* 8:243-246.
- Miyan JA, Mashayekhi F, Bannister CM (2001) Developmental abnormalities in early-onset hydrocephalus: clues to signalling. *Symp Soc Exp Biol*:91-106.
- Miyan JA, Khan MI, Kawarada Y, Sugiyama T, Bannister CM (1998) Cell death in the brain of the HTx rat. *Eur J Pediatr Surg* 8 Suppl 1:43-48.
- Miyazaki I, Asanuma M, Diaz-Corrales FJ, Miyoshi K, Ogawa N (2004) Direct evidence for expression of dopamine receptors in astrocytes from basal ganglia. *Brain Res* 1029:120-123.
- Monnerie H, Dastugue B, Meiniel A (1998) Effect of synthetic peptides derived from SCO-spondin conserved domains on chick cortical and spinal-cord neurons in cell cultures. *Cell Tissue Res* 293:407-418.
- Mooney RA, Senn J, Cameron S, Inamdar N, Boivin LM, Shang Y, Furlanetto RW (2001) Suppressors of cytokine signaling-1 and -6 associate with and inhibit the insulin receptor. A potential mechanism for cytokine-mediated insulin resistance. *J Biol Chem* 276:25889-25893.

- Moos T, Mollgard K (1993) A sensitive post-DAB enhancement technique for demonstration of iron in the central nervous system. *Histochemistry* 99:471-475.
- Morgan FW, Stewart JA, Smith AN, Tarnuzzer RW (2005) Differential expression of stress response genes in the H-Tx rat model of congenital hydrocephalus. *Brain Res Mol Brain Res* 138:273-290.
- Mori E (2008) [Gait disturbance in idiopathic normal pressure hydrocephalus]. *Brain Nerve* 60:219-224.
- Mori F, Tanji K, Yoshida Y, Wakabayashi K (2002) Thalamic retrograde degeneration in the congenitally hydrocephalic rat is attributable to apoptotic cell death. *Neuropathology* 22:186-193.
- Mori K, Shimada J, Kurisaka M, Sato K, Watanabe K (1995) Classification of hydrocephalus and outcome of treatment. *Brain Dev* 17:338-348.
- Murcia NS, Richards WG, Yoder BK, Mucenski ML, Dunlap JR, Woychik RP (2000) The Oak Ridge Polycystic Kidney (orpk) disease gene is required for left-right axis determination. *Development* 127:2347-2355.
- Nakada J, Oka N, Nagahori T, Endo S, Takaku A (1992) Changes in the cerebral vascular bed in experimental hydrocephalus: an angio-architectural and histological study. *Acta Neurochir (Wien)* 114:43-50.
- Nakamura Y, Sato K (1993) Role of disturbance of ependymal ciliary movement in development of hydrocephalus in rats. *Childs Nerv Syst* 9:65-71.
- Nakano H, Bando K, Miyaoka M, Sato K (1996) Evaluation of hydrocephalic periventricular radiolucency by dynamic computed tomography and xenon-computed tomography. *Neurosurgery* 39:758-762; discussion 762-753.
- Nayak PK, Agrawal D, Gowda NK, Bal CS, Mahapatra AK (2005) Does cerebral perfusion improve following CSF diversion in pediatric hydrocephalus? A prospective study using 99mTc ECD single photon emission computed tomography. *Pediatr Neurosurg* 41:117-121.
- Nelson KB, Grether JK (1995) Can magnesium sulfate reduce the risk of cerebral palsy in very low birthweight infants? *Pediatrics* 95:263-269.
- Nelson LH, Anderson SG, Penry MF (1984) The wavering midline: a diagnostic sign of fetal hydrocephalus. *Am J Obstet Gynecol* 149:662-665.
- Newberne PM (1962) The subcommissural organ of the vitamin B12-deficient rat. *J Nutr* 76:393-413.
- Newberne PM, O'Dell BL (1959) Pathology of vitamin B12 deficiency in infant rats. *J Nutr* 68:343-358.

- Newman EA, Zahs KR (1998) Modulation of neuronal activity by glial cells in the retina. *J Neurosci* 18:4022-4028.
- Newton H, Pickard JD, Weller RO (1989) Normal pressure hydrocephalus and cerebrovascular disease: findings of postmortem. *J Neurol Neurosurg Psychiatry* 52:804.
- Nicholson C (1999) Signals that go with the flow. *Trends Neurosci* 22:143-145.
- Nicholson C, Sykova E (1998) Extracellular space structure revealed by diffusion analysis. *Trends Neurosci* 21:207-215.
- Nicol B, Rowbotham DJ, Lambert DG (1996) mu- and kappa-opioids inhibit K<sup>+</sup> evoked glutamate release from rat cerebrocortical slices. *Neurosci Lett* 218:79-82.
- Nielsen SL, Baringer JR (1972) Reovirus-induced aqueductal stenosis in hamsters. Phase contrast and electron microscopic studies. *Lab Invest* 27:531-537.
- Nielsen SL, Gauger GE (1974) Experimental hydrocephalus: surface alterations of the lateral ventricle. Scanning electron microscopic studies. *Lab Invest* 30:618-625.
- Nilsson C, Lindvall-Axelsson M, Owman C (1991) Simultaneous and continuous measurement of choroid plexus blood flow and cerebrospinal fluid production: effects of vasoactive intestinal polypeptide. *J Cereb Blood Flow Metab* 11:861-867.
- Nilsson C, Stahlberg F, Thomsen C, Henriksen O, Herning M, Owman C (1992) Circadian variation in human cerebrospinal fluid production measured by magnetic resonance imaging. *Am J Physiol* 262:R20-24.
- Nogi H, Hashimoto H, Hagihara N, Shimada S, Yamamoto K, Matsuda T, Tohyama M, Baba A (1997) Distribution of mRNAs for pituitary adenylate cyclase-activating polypeptide (PACAP), PACAP receptor, vasoactive intestinal polypeptide (VIP), and VIP receptors in the rat superior cervical ganglion. *Neurosci Lett* 227:37-40.
- Nojima Y, Enzan H, Hayashi Y, Nakayama H, Kiyoku H, Hiroi M, Mori K (1998) Neuroepithelial and ependymal changes in HTX rats with congenital hydrocephalus: an ultrastructural and immunohistochemical study. *Pathol Int* 48:115-125.
- Nonaka Y, Miyajima M, Ogino I, Nakajima M, Arai H (2008) Analysis of neuronal cell death in the cerebral cortex of H-Tx rats with compensated hydrocephalus. *J Neurosurg Pediatrics* 1:68-74.
- Nowak DA, Topka HR (2006) Broadening a classic clinical triad: The hypokinetic motor disorder of normal pressure hydrocephalus also affects the hand. *Exp Neurol* 198:81-87.
- Nulsen FE, Spitz EB (1951) Treatment of hydrocephalus by direct shunt from ventricle to jugular vein. *Surg Forum*:399-403.

- Oi S, Matsumoto S (1985) [Postoperative management of patients with hydrocephalus and complications following shunts]. *Kango Gijutsu* 31:302-312.
- Oi S, Ijichi A, Matsumoto S (1989) Immunohistochemical evaluation of neuronal maturation in untreated fetal hydrocephalus. *Neurol Med Chir (Tokyo)* 29:989-994.
- Oi S, Yamada H, Sato O, Matsumoto S (1996) Experimental models of congenital hydrocephalus and comparable clinical problems in the fetal and neonatal periods. *Childs Nerv Syst* 12:292-302.
- Oi S, Shimoda M, Shibata M, Honda Y, Togo K, Shinoda M, Tsugane R, Sato O (2000) Pathophysiology of long-standing overt ventriculomegaly in adults. *J Neurosurg* 92:933-940.
- Oka N, Nakada J, Endo S, Takaku A (1985) Angioarchitecture in experimental hydrocephalus. *Pediatr Neurosci* 12:294-299.
- Okuyama T, Hashi K, Sasaki S, Sudo K, Kurokawa Y (1987) Changes in cerebral microvasculature in congenital hydrocephalus of the inbred rat LEW/Jms: light and electron microscopic examination. *Surg Neurol* 27:338-342.
- Oldendorf WH (1970) Measurement of brain uptake of radiolabeled substances using a tritiated water internal standard. *Brain Res* 24:372-376.
- Oliveira ES, Leite PE, Spillantini MG, Camargo AC, Hunt SP (1990) Localization of endo-oligopeptidase (EC 3.4.22.19) in the rat nervous tissue. *J Neurochem* 55:1114-1121.
- Otto C, Hein L, Brede M, Jahns R, Engelhardt S, Grone H-J, Schutz G (2004) Pulmonary Hypertension and Right Heart Failure in Pituitary Adenylate Cyclase-Activating Polypeptide Type I Receptor-Deficient Mice. *Circulation* 110:3245-3251.
- Overholser MD, Whitley JR, O'Dell BL, Hogan AG (1954) The ventricular system in hydrocephalic rat brains produced by a deficiency of vitamin B12 or of folic acid in the maternal diet. *Anat Rec* 120:917-933.
- Owen-Lynch PJ, Draper CE, Mashayekhi F, Bannister CM, Miyan JA (2003) Defective cell cycle control underlies abnormal cortical development in the hydrocephalic Texas rat. *Brain* 126:623-631.
- Pace AM, Wong YH, Bourne HR (1991) A mutant alpha subunit of Gi2 induces neoplastic transformation of Rat-1 cells. *Proc Natl Acad Sci U S A* 88:7031-7035.
- Pacheco MA, Stockmeier C, Meltzer HY, Overholser JC, Dilley GE, Jope RS (1996) Alterations in phosphoinositide signaling and G-protein levels in depressed suicide brain. *Brain Res* 723:37-45.
- Paez P, Batiz LF, Roales-Bujan R, Rodriguez-Perez LM, Rodriguez S, Jimenez AJ, Rodriguez EM, Perez-Figares JM (2007) Patterned neuropathologic events occurring

- in hyh congenital hydrocephalic mutant mice. *J Neuropathol Exp Neurol* 66:1082-1092.
- Page RB (1975) Scanning electron microscopy of the ventricular system in normal and hydrocephalic rabbits. Preliminary report and atlas. *J Neurosurg* 42:646-664.
- Page RB, Leure-duPree AE (1983) Ependymal alterations in hydrocephalus. In: *Neurobiology of Cerebrospinal Fluid* (Wood JH, ed), pp 789-820. New York: Plenum Publishing Corporation.
- Pan J, Misamore MJ, Wang Q, Snell WJ (2003) Protein transport and signal transduction during fertilization in chlamydomonas. *Traffic* 4:452-459.
- Parisi MA, Dobyns WB (2003) Human malformations of the midbrain and hindbrain: review and proposed classification scheme. *Mol Genet Metab* 80:36-53.
- Parpura V, Basarsky TA, Liu F, Jęftinija K, Jęftinija S, Haydon PG (1994) Glutamate-mediated astrocyte-neuron signalling. *Nature* 369:744-747.
- Parri HR, Gould TM, Crunelli V (2001) Spontaneous astrocytic Ca<sup>2+</sup> oscillations in situ drive NMDAR-mediated neuronal excitation. *Nat Neurosci* 4:803-812.
- Pasti L, Volterra A, Pozzan T, Carmignoto G (1997) Intracellular calcium oscillations in astrocytes: a highly plastic, bidirectional form of communication between neurons and astrocytes in situ. *J Neurosci* 17:7817-7830.
- Pasti L, Zonta M, Pozzan T, Vicini S, Carmignoto G (2001) Cytosolic calcium oscillations in astrocytes may regulate exocytotic release of glutamate. *J Neurosci* 21:477-484.
- Paulson JC (1989) Glycoproteins: what are the sugar chains for? *Trends Biochem Sci* 14:272-276.
- Peacock WJ, Currer TH (1984) Hydrocephalus in childhood. A study of 440 cases. *S Afr Med J* 66:323-324.
- Pearce RK, Collins P, Jenner P, Emmett C, Marsden CD (1996) Intraventricular infusion of basic fibroblast growth factor (bFGF) in the MPTP-treated common marmoset. *Synapse* 23:192-200.
- Pearson MM, Lu J, Mount DB, Delpire E (2001) Localization of the K(+)-Cl(-) cotransporter, KCC3, in the central and peripheral nervous systems: expression in the choroid plexus, large neurons and white matter tracts. *Neuroscience* 103:481-491.
- Perez-Figares JM, Jimenez AJ, Rodriguez EM (2001) Subcommissural organ, cerebrospinal fluid circulation, and hydrocephalus. *Microsc Res Tech* 52:591-607.
- Perez-Figares JM, Jimenez AJ, Perez-Martin M, Fernandez-Llebrez P, Cifuentes M, Riera P, Rodriguez S, Rodriguez EM (1998) Spontaneous congenital hydrocephalus in the

- mutant mouse hyh. Changes in the ventricular system and the subcommissural organ. *J Neuropathol Exp Neurol* 57:188-202.
- Persson EK, Hagberg G, Uvebrant P (2005) Hydrocephalus prevalence and outcome in a population-based cohort of children born in 1989-1998. *Acta Paediatr* 94:726-732.
- Persson EK, Hagberg G, Uvebrant P (2006) Disabilities in children with hydrocephalus--a population-based study of children aged between four and twelve years. *Neuropediatrics* 37:330-336.
- Petrella G, Czosnyka M, Keong N, Pickard JD, Czosnyka Z (2008) How does CSF dynamics change after shunting? *Acta Neurol Scand* 118:182-188.
- Picco P, Leveratto L, Cama A, Vigliarolo MA, Levato GL, Gattorno M, Zammarchi E, Donati MA (1993) Immobile cilia syndrome associated with hydrocephalus and precocious puberty: a case report. *Eur J Pediatr Surg* 3 Suppl 1:20-21.
- Pople IK (1992) Doppler flow velocities in children with controlled hydrocephalus: reference values for the diagnosis of blocked cerebrospinal fluid shunts. *Childs Nerv Syst* 8:124-125.
- Pople IK, Quinn MW, Bayston R (1990) Morbidity and outcome of shunted hydrocephalus. *Z Kinderchir* 45 Suppl 1:29-31.
- Porter JT, McCarthy KD (1996) Hippocampal astrocytes in situ respond to glutamate released from synaptic terminals. *J Neurosci* 16:5073-5081.
- Povlishock JT, Jenkins LW (1995) Are the pathobiological changes evoked by traumatic brain injury immediate and irreversible? *Brain Pathol* 5:415-426.
- Powell KR, Holtzman SG (1999) Differential antagonism of the rate-decreasing effects of kappa-opioid receptor agonists by naltrexone and norbinaltorphimine. *Eur J Pharmacol* 377:21-28.
- Praetorius HA, Spring KR (2001) Bending the MDCK cell primary cilium increases intracellular calcium. *J Membr Biol* 184:71-79.
- Praetorius J, Nejsum LN, Nielsen S (2004) A SLC4A10 gene product maps selectively to the basolateral plasma membrane of choroid plexus epithelial cells. *Am J Physiol Cell Physiol* 286:C601-610.
- Preston JE (2001) Ageing choroid plexus-cerebrospinal fluid system. *Microsc Res Tech* 52:31-37.
- Preston JE, McMillan PN, Stopa EG, Nashold JR, Duncan JA, Johanson CE (2003) Atrial natriuretic peptide induction of dark epithelial cells in choroid plexus: consistency with the model of CSF downregulation in hydrocephalus. *Eur J Pediatr Surg* 13 Suppl 1:S40-42.



- Price DL, James AE, Jr., Sperber E, Strecker EP (1976) Communicating hydrocephalus. Cisternographic and neuropathologic studies. *Arch Neurol* 33:15-20.
- Proescholdt MG, Hutto B, Brady LS, Herkenham M (2000) Studies of cerebrospinal fluid flow and penetration into brain following lateral ventricle and cisterna magna injections of the tracer [<sup>14</sup>C]inulin in rat. *Neuroscience* 95:577-592.
- Quencer RM (1992) Intracranial CSF flow in pediatric hydrocephalus: evaluation with cine-MR imaging. *AJNR Am J Neuroradiol* 13:601-608.
- Raimondi AJ, Soare P (1974) Intellectual development in shunted hydrocephalic children. *Am J Dis Child* 127:664-671.
- Raimondi AJ, Clark SJ, McLone DG (1976) Pathogenesis of aqueductal occlusion in congenital murine hydrocephalus. *J Neurosurg* 45:66-77.
- Rapoport SI (1976) Opening of the blood-brain barrier by acute hypertension. *Exp Neurol* 52:467-479.
- Rascher K, Booz KH, Donauer E, Nacimiento AC (1987) Structural alterations in the spinal cord during progressive communicating syringomyelia. An experimental study in the cat. *Acta Neuropathol* 72:248-255.
- Redfern CH, Coward P, Degtyarev MY, Lee EK, Kwa AT, Hennighausen L, Bujard H, Fishman GI, Conklin BR (1999) Conditional expression and signaling of a specifically designed Gi-coupled receptor in transgenic mice. *Nat Biotechnol* 17:165-169.
- Redfern CH, Degtyarev MY, Kwa AT, Salomonis N, Cotte N, Nanevich T, Fidelman N, Desai K, Vranizan K, Lee EK, Coward P, Shah N, Warrington JA, Fishman GI, Bernstein D, Baker AJ, Conklin BR (2000) Conditional expression of a Gi-coupled receptor causes ventricular conduction delay and a lethal cardiomyopathy. *Proc Natl Acad Sci U S A* 97:4826-4831.
- Reed DJ, Woodbury DM (1963) Kinetics of Movement of Iodide, Sucrose, Inulin and Radio-Iodinated Serum Albumin in the Central Nervous System and Cerebrospinal Fluid of the Rat. *J Physiol* 169:816-850.
- Rekate HL (2004) Selecting patients for endoscopic third ventriculostomy. *Neurosurg Clin N Am* 15:39-49.
- Rekate HL (2008) Shunt-related headaches: the slit ventricle syndromes. *Childs Nerv Syst* 24:423-430.
- Renier D, Lacombe J, Pierre-Kahn A, Sainte-Rose C, Hirsch JF (1984) Factors causing acute shunt infection. Computer analysis of 1174 operations. *J Neurosurg* 61:1072-1078.

- Richardson RR (1985) Congenital genetic murine (ch) hydrocephalus. A structural model of cellular dysplasia and disorganization with the molecular locus of deficient proteoglycan synthesis. *Childs Nerv Syst* 1:87-99.
- Rickert CH, Grabellus F, Varchmin-Schultheiss K, Stoss H, Paulus W (2001) Sudden unexpected death in young adults with chronic hydrocephalus. *Int J Legal Med* 114:331-337.
- Riikonen R, Somer M, Turpeinen U (1999) Low insulin-like growth factor (IGF-1) in the cerebrospinal fluid of children with progressive encephalopathy, hypsarrhythmia, and optic atrophy (PEHO) syndrome and cerebellar degeneration. *Epilepsia* 40:1642-1648.
- Ritter S, Dinh TT, Stone S, Ross N (1988) Cerebroventricular dilation in spontaneously hypertensive rats (SHRs) is not attenuated by reduction of blood pressure. *Brain Res* 450:354-359.
- Rivkin E, Cullinan EB, Tres LL, Kierszenbaum AL (1997) A protein associated with the manchette during rat spermiogenesis is encoded by a gene of the TBP-1-like subfamily with highly conserved ATPase and protease domains. *Mol Reprod Dev* 48:77-89.
- Roberts AB, Sporn MB (1996) Transforming Growth Factor-Beta. In: *The Molecular and Cellular Biology of Wound Repair*, Second Edition Edition (Clark RAF, ed), p 275. New York: Plenum Press.
- Robertson SP, Gattas M, Rogers M, Ades LC (2000) Macrocephaly--cutis marmorata telangiectatica congenita: report of five patients and a review of the literature. *Clin Dysmorphol* 9:1-9.
- Robinson ML, Allen CE, Davy BE, Durfee WJ, Elder FF, Elliott CS, Harrison WR (2002) Genetic mapping of an insertional hydrocephalus-inducing mutation allelic to hy3. *Mamm Genome* 13:625-632.
- Rodriguez-Perez LM, Perez-Martin M, Jimenez AJ, Fernandez-Llebrez P (2003) Immunocytochemical characterisation of the wall of the bovine lateral ventricle. *Cell Tissue Res* 314:325-335.
- Rodriguez EM (1976) The cerebrospinal fluid as a pathway in neuroendocrine integration. *J Endocrinol* 71:407-443.
- Rodriguez EM, Oksche A, Montecinos H (2001) Human subcommissural organ, with particular emphasis on its secretory activity during the fetal life. *Microsc Res Tech* 52:573-590.
- Rodriguez EM, Oksche A, Rodriguez S, Nualart F, Irigoien C, Fernandez-Llebrez P, Perez-Figueras JM (1990) Subcommissural organ and cerebrospinal fluid: a biochemical and experimental study. *Prog Clin Biol Res* 342:282-292.

- Rolf B, Kutsche M, Bartsch U (2001) Severe hydrocephalus in L1-deficient mice. *Brain Res* 891:247-252.
- Ronan A, Hogg GG, Klug GL (1995) Cerebrospinal fluid shunt infections in children. *Pediatr Infect Dis J* 14:782-786.
- Rosen H, Swigar ME (1976) Depression and normal pressure hydrocephalus. A dilemma in neuropsychiatric differential diagnosis. *J Nerv Ment Dis* 163:35-40.
- Rosenberg GA, Kyner WT, Fenstermacher JD, Patlak CS (1986) Effect of vasopressin on ependymal and capillary permeability to tritiated water in cat. *Am J Physiol* 251:F485-489.
- Roth Y, Kimhi Y, Edery H, Aharonson E, Priel Z (1985) Ciliary motility in brain ventricular system and trachea of hamsters. *Brain Res* 330:291-297.
- Rudolph D, Tafuri A, Gass P, Hammerling GJ, Arnold B, Schutz G (1998) Impaired fetal T cell development and perinatal lethality in mice lacking the cAMP response element binding protein. *Proc Natl Acad Sci U S A* 95:4481-4486.
- Rutter M, Tizard J, Yule W, Graham P, Whitmore K (1976) Research report: Isle of Wight Studies, 1964-1974. *Psychol Med* 6:313-332.
- Sainte-Rose C, Piatt JH, Renier D, Pierre-Kahn A, Hirsch JF, Hoffman HJ, Humphreys RP, Hendrick EB (1991) Mechanical complications in shunts. *Pediatr Neurosurg* 17:2-9.
- Saitoh T, Horsburgh K, Masliah E (1993) Hyperactivation of signal transduction systems in Alzheimer's disease. *Ann N Y Acad Sci* 695:34-41.
- Sakakibara S, Nakamura Y, Yoshida T, Shibata S, Koike M, Takano H, Ueda S, Uchiyama Y, Noda T, Okano H (2002) RNA-binding protein Musashi family: roles for CNS stem cells and a subpopulation of ependymal cells revealed by targeted disruption and antisense ablation. *Proc Natl Acad Sci U S A* 99:15194-15199.
- Sameshima H, Ikenoue T (2001) Long-term magnesium sulfate treatment as protection against hypoxic-ischemic brain injury in seven-day-old rats. *Am J Obstet Gynecol* 184:185-190.
- Sapiro R, Kostetskii I, Olds-Clarke P, Gerton GL, Radice GL, Strauss IJ (2002) Male infertility, impaired sperm motility, and hydrocephalus in mice deficient in sperm-associated antigen 6. *Mol Cell Biol* 22:6298-6305.
- Sarnat HB (1992) Regional differentiation of the human fetal ependyma: immunocytochemical markers. *J Neuropathol Exp Neurol* 51:58-75.
- Sarnat HB (1995) Ependymal reactions to injury. A review. *J Neuropathol Exp Neurol* 54:1-15.

- Satir P, Sleight MA (1990) The physiology of cilia and mucociliary interactions. *Annu Rev Physiol* 52:137-155.
- Sato H, Sato N, Tamaki N, Matsumoto S (1988) Threshold of cerebral perfusion pressure as a prognostic factor in hydrocephalus during infancy. *Childs Nerv Syst* 4:274-278.
- Sato O, Hara M, Asai T, Tsugane R, Kageyama N (1973) The effect of dexamethasone phosphate on the production rate of cerebrospinal fluid in the spinal subarachnoid space of dogs. *J Neurosurg* 39:480-484.
- Scarff JE (1963) Treatment of hydrocephalus: an historical and critical review of methods and results. *J Neurol Neurosurg Psychiatry* 26:1-26.
- Schick RW, Matson DD (1961) What is arrested hydrocephalus? *J Pediatr* 58:791-799.
- Schmid RS, Pruitt WM, Maness PF (2000) A MAP kinase-signaling pathway mediates neurite outgrowth on L1 and requires Src-dependent endocytosis. *J Neurosci* 20:4177-4188.
- Schmidt JF, Albeck M, Gjerris F (1990) The effect of nimodipine on ICP and CBF in patients with normal-pressure hydrocephalus. *Acta Neurochir (Wien)* 102:11-13.
- Schoeman J, Donald P, van Zyl L, Keet M, Wait J (1991) Tuberculous hydrocephalus: comparison of different treatments with regard to ICP, ventricular size and clinical outcome. *Dev Med Child Neurol* 33:396-405.
- Schoenbaum SC, Gardner P, Shillito J (1975) Infections of cerebrospinal fluid shunts: epidemiology, clinical manifestations, and therapy. *J Infect Dis* 131:543-552.
- Schoffelman AN, Hogenboom F, Mulder AH (1997) Kappa1- and kappa2-opioid receptors mediating presynaptic inhibition of dopamine and acetylcholine release in rat neostriatum. *Br J Pharmacol* 122:520-524.
- Scholey AB, Mileusnic R, Schachner M, Rose SP (1995) A role for a chicken homolog of the neural cell adhesion molecule L1 in consolidation of memory for a passive avoidance task in the chick. *Learn Mem* 2:17-25.
- Schossberger PF, Touya JJ (1976) Dynamic cisternography in normal dogs and in human beings. *Neurology* 26:254-260.
- Schwartz MW, Marks JL, Sipols AJ, Baskin DG, Woods SC, Kahn SE, Porte D, Jr. (1991) Central insulin administration reduces neuropeptide Y mRNA expression in the arcuate nucleus of food-deprived lean (Fa/Fa) but not obese (fa/fa) Zucker rats. *Endocrinology* 128:2645-2647.
- Schwartz TW (1994) Locating ligand-binding sites in 7TM receptors by protein engineering. *Curr Opin Biotechnol* 5:434-444.

- Scott MA, Fletcher JM, Brookshire BL, Davidson KC, Landry SH, Bohan TC, Kramer LA, Brandt ME, Francis DJ (1998) Memory functions in children with early hydrocephalus. *Neuropsychology* 12:578-589.
- Scriabine A, van den Kerckhoff W (1988) Pharmacology of nimodipine. A review. *Ann N Y Acad Sci* 522:698-706.
- Scriabine A, Schuurman T, Traber J (1989) Pharmacological basis for the use of nimodipine in central nervous system disorders. *Faseb J* 3:1799-1806.
- Segal HDaMB (1996) *Physiology of the CSF and Blood-Brain Barriers*. Boca Raton, FL: CRC Press.
- Senut MC, Jazat F, Choi NH, Lamour Y (1992) Protein SP40,40-like Immunoreactivity in the Rat Brain: Progressive Increase With Age. *Eur J Neurosci* 4:917-928.
- Serot JM, Bene MC, Faure GC (2003) Normal-pressure hydrocephalus and Alzheimer disease. *J Neurosurg* 99:797-798; author reply 798-799.
- Serot JM, Bene MC, Foliguet B, Faure GC (1997a) Altered choroid plexus basement membrane and epithelium in late-onset Alzheimer's disease: an ultrastructural study. *Ann N Y Acad Sci* 826:507-509.
- Serot JM, Christmann D, Dubost T, Couturier M (1997b) Cerebrospinal fluid transthyretin: aging and late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 63:506-508.
- Shadmehr R, Holcomb HH (1997) Neural correlates of motor memory consolidation. *Science* 277:821-825.
- Shank RP, Bennett GS, Freytag SO, Campbell GL (1985) Pyruvate carboxylase: an astrocyte-specific enzyme implicated in the replenishment of amino acid neurotransmitter pools. *Brain Res* 329:364-367.
- Shapira Y, Artru AA, Lam AM (1992) Changes in the rate of formation and resistance to reabsorption of cerebrospinal fluid during deliberate hypotension induced with adenosine or hemorrhage. *Anesthesiology* 76:432-439.
- Sheth RD (1998) Trends in incidence and severity of intraventricular hemorrhage. *J Child Neurol* 13:261-264.
- Shih WJ, Tasdemiroglu E (1995) Reversible hypoperfusion of the cerebral cortex in normal-pressure hydrocephalus on technetium-99m-HMPAO brain SPECT images after shunt operation. *J Nucl Med* 36:470-473.
- Shim I, Ha Y, Chung JY, Lee HJ, Yang KH, Chang JW (2003) Association of learning and memory impairments with changes in the septohippocampal cholinergic system in rats with kaolin-induced hydrocephalus. *Neurosurgery* 53:416-425; discussion 425.

- Shinnar S, Gammon K, Bergman EW, Jr., Epstein M, Freeman JM (1985) Management of hydrocephalus in infancy: use of acetazolamide and furosemide to avoid cerebrospinal fluid shunts. *J Pediatr* 107:31-37.
- Shoesmith CL, Buist R, Del Bigio MR (2000) Magnetic resonance imaging study of extracellular fluid tracer movement in brains of immature rats with hydrocephalus. *Neurol Res* 22:111-116.
- Shukla D, Singh BM, Strobos RJ (1980) Hypertensive cerebrovascular disease and normal pressure hydrocephalus. *Neurology* 30:998-1000.
- Silverberg GD (2004) Normal pressure hydrocephalus (NPH): ischaemia, CSF stagnation or both. *Brain* 127:947-948.
- Silverberg GD, Mayo M, Saul T, Rubenstein E, McGuire D (2003) Alzheimer's disease, normal-pressure hydrocephalus, and senescent changes in CSF circulatory physiology: a hypothesis. *Lancet Neurol* 2:506-511.
- Silverberg GD, Huhn S, Jaffe RA, Chang SD, Saul T, Heit G, Von Essen A, Rubenstein E (2002) Downregulation of cerebrospinal fluid production in patients with chronic hydrocephalus. *J Neurosurg* 97:1271-1275.
- Simpson JI, Eide TR, Schiff GA, Clagnaz JF, Hossain I, Tverskoy A, Koski G (1994) Intrathecal magnesium sulfate protects the spinal cord from ischemic injury during thoracic aortic cross-clamping. *Anesthesiology* 81:1493-1499; discussion 1426A-1427A.
- Siomin V, Weiner H, Wisoff J, Cinalli G, Pierre-Kahn A, Saint-Rose C, Abbott R, Elran H, Beni-Adani L, Ouaknine G, Constantini S (2001) Repeat endoscopic third ventriculostomy: is it worth trying? *Childs Nerv Syst* 17:551-555.
- Slotkin TA, Wang XF, Symonds HS, Seidler FJ (1997) Expression of mRNAs coding for the transforming growth factor-beta receptors in brain regions of euthyroid and hypothyroid neonatal rats and in adult brain. *Brain Res Dev Brain Res* 99:61-65.
- Socci DJ, Bjugstad KB, Jones HC, Pattisapu JV, Arendash GW (1999) Evidence that oxidative stress is associated with the pathophysiology of inherited hydrocephalus in the H-Tx rat model. *Exp Neurol* 155:109-117.
- Soderborg B, Lying-Tunell U (1972) How much radioactivity reaches the basal cisterns at cisternography? *Strahlentherapie [Sonderb]* 72:203-207.
- Soelberg Sorensen P, Jansen EC, Gjerris F (1986) Motor disturbances in normal-pressure hydrocephalus. Special reference to stance and gait. *Arch Neurol* 43:34-38.
- Somera KC, Jones HC (2004) Reduced subcommissural organ glycoprotein immunoreactivity precedes aqueduct closure and ventricular dilatation in H-Tx rat hydrocephalus. *Cell Tissue Res* 315:361-373.

- Sorensen PS (1986) Studies of vasopressin in the human cerebrospinal fluid. *Acta Neurol Scand* 74:81-102.
- Spiegel AM (1992a) Heterotrimeric GTP-binding proteins: an expanding family of signal transducers. *Med Res Rev* 12:55-71.
- Spiegel AM (1992b) G proteins in cellular control. *Curr Opin Cell Biol* 4:203-211.
- Srebro Z (1969) The effect of intracranial puromycin and ouabain administration on the morphology of the circumventricular organs and the ependyma in the frog, *Rana esculenta*. *Folia Biol (Krakow)* 17:249-258.
- Starr R, Willson TA, Viney EM, Murray LJ, Rayner JR, Jenkins BJ, Gonda TJ, Alexander WS, Metcalf D, Nicola NA, Hilton DJ (1997) A family of cytokine-inducible inhibitors of signalling. *Nature* 387:917-921.
- Strazielle N, Gherzi-Egea JF (2000) Choroid plexus in the central nervous system: biology and physiopathology. *J Neuropathol Exp Neurol* 59:561-574.
- Stys PK (1998) Anoxic and ischemic injury of myelinated axons in CNS white matter: from mechanistic concepts to therapeutics. *J Cereb Blood Flow Metab* 18:2-25.
- Suzaki I, Hara T, Tanaka C, Dejima S, Takeshita K (1997) Elevated nerve growth factor levels in cerebrospinal fluid associated with progressive cortical atrophy. *Neuropediatrics* 28:268-271.
- Suzer T, Coskun E, Islekel H, Tahta K (1999) Neuroprotective effect of magnesium on lipid peroxidation and axonal function after experimental spinal cord injury. *Spinal Cord* 37:480-484.
- Suzuki K, Nakajima K, Otaki N, Kimura M, Kawaharada U, Uehara K, Hara F, Nakazato Y, Takatama M (1994) Localization of metallothionein in aged human brain. *Pathol Int* 44:20-26.
- Svingos AL, Colago EE (2002) Kappa-Opioid and NMDA glutamate receptors are differentially targeted within rat medial prefrontal cortex. *Brain Res* 946:262-271.
- Sweger EJ, Casper KB, Searce-Levie K, Conklin BR, McCarthy KD (2007) Development of hydrocephalus in mice expressing the G(i)-coupled GPCR Ro1 RASSL receptor in astrocytes. *J Neurosci* 27:2309-2317.
- Sykova E, Fiala J, Antonova T, Vorisek I (2001) Extracellular space volume changes and diffusion barriers in rats with kaolin-induced and inherited hydrocephalus. *Eur J Pediatr Surg* 11 Suppl 1:S34-37.
- Tada T, Kanaji M, Kobayashi S (1994) Induction of communicating hydrocephalus in mice by intrathecal injection of human recombinant transforming growth factor-beta 1. *J Neuroimmunol* 50:153-158.

- Takahashi Y (2006) Long-term outcome and neurologic development after endoscopic third ventriculostomy versus shunting during infancy. *Childs Nerv Syst* 22:1591-1602.
- Takano T, Rutka JT, Becker LE (1996) Overexpression of nestin and vimentin in ependymal cells in hydrocephalus. *Acta Neuropathol* 92:90-97.
- Takei F, Sato O (1995) Morphological analysis of progressive hydrocephalus and shunt-dependent arrested hydrocephalus. An experimental study. *Pediatr Neurosurg* 23:246-253.
- Takeuchi IK, Takeuchi YK (1986) Congenital hydrocephalus following X-irradiation of pregnant rats on an early gestational day. *Neurobehav Toxicol Teratol* 8:143-150.
- Takeuchi IK, Kimura R, Matsuda M, Shoji R (1987) Absence of subcommissural organ in the cerebral aqueduct of congenital hydrocephalus spontaneously occurring in MT/HokIdr mice. *Acta Neuropathol* 73:320-322.
- Tardieu M, Goffinet A, Harmant-van Rijckevorsel G, Lyon G (1982) Ependymitis, leukoencephalitis, hydrocephalus, and thrombotic vasculitis following chronic infection by mouse hepatitis virus 3 (MHV 3). *Acta Neuropathol* 58:168-176.
- Tarlow MJ, Jenkins R, Comis SD, Osborne MP, Stephens S, Stanley P, Crocker J (1993) Ependymal cells of the choroid plexus express tumour necrosis factor- $\alpha$ . *Neuropathol Appl Neurobiol* 19:324-328.
- Tashiro Y, Drake JM, Chakraborty S, Hattori T (1997) Functional injury of cholinergic, GABAergic and dopaminergic systems in the basal ganglia of adult rat with kaolin-induced hydrocephalus. *Brain Res* 770:45-52.
- Taulman PD, Haycraft CJ, Balkovetz DF, Yoder BK (2001) Polaris, a protein involved in left-right axis patterning, localizes to basal bodies and cilia. *Mol Biol Cell* 12:589-599.
- Tayarani I, Cloez I, Clement M, Bourre JM (1989) Antioxidant enzymes and related trace elements in aging brain capillaries and choroid plexus. *J Neurochem* 53:817-824.
- Thomas JM, Hoffman BB (1996) Isoform-specific sensitization of adenylyl cyclase activity by prior activation of inhibitory receptors: role of beta gamma subunits in transducing enhanced activity of the type VI isoform. *Mol Pharmacol* 49:907-914.
- Tomita S, Ueno M, Sakamoto M, Kitahama Y, Ueki M, Maekawa N, Sakamoto H, Gassmann M, Kageyama R, Ueda N, Gonzalez FJ, Takahama Y (2003) Defective brain development in mice lacking the Hif-1 $\alpha$  gene in neural cells. *Mol Cell Biol* 23:6739-6749.
- Torikata C, Kijimoto C, Koto M (1991) Ultrastructure of respiratory cilia of WIC-Hyd male rats. An animal model for human immotile cilia syndrome. *Am J Pathol* 138:341-347.



- Toumane A, Durkin T, Marighetto A, Galey D, Jaffard R (1988) Differential hippocampal and cortical cholinergic activation during the acquisition, retention, reversal and extinction of a spatial discrimination in an 8-arm radial maze by mice. *Behav Brain Res* 30:225-234.
- Tullberg M, Jensen C, Ekholm S, Wikkelsö C (2001) Normal pressure hydrocephalus: vascular white matter changes on MR images must not exclude patients from shunt surgery. *AJNR Am J Neuroradiol* 22:1665-1673.
- Tullberg M, Rosengren L, Blomsterwall E, Karlsson JE, Wikkelsö C (1998) CSF neurofilament and glial fibrillary acidic protein in normal pressure hydrocephalus. *Neurology* 50:1122-1127.
- Tung H, Raffel C, McComb JG (1991) Ventricular cerebrospinal fluid eosinophilia in children with ventriculoperitoneal shunts. *J Neurosurg* 75:541-544.
- Turgut M, Alabaz D, Erbey F, Kocabas E, Erman T, Alhan E, Aksaray N (2005) Cerebrospinal fluid shunt infections in children. *Pediatr Neurosurg* 41:131-136.
- Ulfing N (2002) Neuronal vacuolation in the basal nucleus of meynert caused by fetal hydrocephalus. *Pediatr Neurosurg* 36:320-323.
- Ulfing N, Szabo A, Bohl J (2001) Effect of fetal hydrocephalus on the distribution patterns of calcium-binding proteins in the human occipital cortex. *Pediatr Neurosurg* 34:20-32.
- Uren D, Hwang HK, Hara Y, Takeda K, Kawamoto S, Tullio AN, Yu ZX, Ferrans VJ, Tresser N, Grinberg A, Preston YA, Adelstein RS (2000) Gene dosage affects the cardiac and brain phenotype in nonmuscle myosin II-B-depleted mice. *J Clin Invest* 105:663-671.
- Valdizan EM, Gutierrez O, Pazos A (2003) Adenylate cyclase activity in postmortem brain of suicide subjects: reduced response to beta-adrenergic stimulation. *Biol Psychiatry* 54:1457-1464.
- Vale FA, Miranda SJ (2002) Clinical and demographic features of patients with dementia attended in a tertiary outpatient clinic. *Arq Neuropsiquiatr* 60:548-552.
- Van Camp G, Vits L, Coucke P, Lyonnet S, Schrander-Stumpel C, Darby J, Holden J, Munnich A, Willems PJ (1993) A duplication in the L1CAM gene associated with X-linked hydrocephalus. *Nat Genet* 4:421-425.
- van der Knaap MS, Valk J, de Neeling N, Nauta JJ (1991) Pattern recognition in magnetic resonance imaging of white matter disorders in children and young adults. *Neuroradiology* 33:478-493.
- van der Knaap MS, Smit LM, Barth PG, Catsman-Berrevoets CE, Brouwer OF, Begeer JH, de Coö IF, Valk J (1997) Magnetic resonance imaging in classification of congenital muscular dystrophies with brain abnormalities. *Ann Neurol* 42:50-59.

- Vanneste JA (2000) Diagnosis and management of normal-pressure hydrocephalus. *J Neurol* 247:5-14.
- Vernadakis A (1996) Glia-neuron intercommunications and synaptic plasticity. *Prog Neurobiol* 49:185-214.
- Vertinsky AT, Barnes PD (2007) Macrocephaly, increased intracranial pressure, and hydrocephalus in the infant and young child. *Top Magn Reson Imaging* 18:31-51.
- Vetsika EK, Bannister CM, Buckle AM, Miyan JA (1999) The effects of CSF blockage in early-onset hydrocephalus on the activity of the germinal epithelium. *Eur J Pediatr Surg* 9 Suppl 1:43-44.
- Vigers AJ, Pfenninger KH (1991) N-type and L-type calcium channels are present in nerve growth cones. Numbers increase on synaptogenesis. *Brain Res Dev Brain Res* 60:197-203.
- Vink R, Nimmo AJ, Cernak I (2001) An overview of new and novel pharmacotherapies for use in traumatic brain injury. *Clin Exp Pharmacol Physiol* 28:919-921.
- Vio K, Rodriguez S, Navarrete EH, Perez-Figares JM, Jimenez AJ, Rodriguez EM (2000) Hydrocephalus induced by immunological blockage of the subcommissural organ-Reissner's fiber (RF) complex by maternal transfer of anti-RF antibodies. *Exp Brain Res* 135:41-52.
- Virella AA, Galarza M, Masterman-Smith M, Lemus R, Lazareff JA (2002) Distal slit valve and clinically relevant CSF overdrainage in children with hydrocephalus. *Childs Nerv Syst* 18:15-18.
- Vladic A, Klarica M, Bulat M (2009) Dynamics of distribution of 3H-inulin between the cerebrospinal fluid compartments. *Brain Res* 1248:127-135.
- Wagner C, Batiz LF, Rodriguez S, Jimenez AJ, Paez P, Tome M, Perez-Figares JM, Rodriguez EM (2003) Cellular mechanisms involved in the stenosis and obliteration of the cerebral aqueduct of hyh mutant mice developing congenital hydrocephalus. *J Neuropathol Exp Neurol* 62:1019-1040.
- Walters BC, Hoffman HJ, Hendrick EB, Humphreys RP (1984) Cerebrospinal fluid shunt infection. Influences on initial management and subsequent outcome. *J Neurosurg* 60:1014-1021.
- Walz W (1989) Role of glial cells in the regulation of the brain ion microenvironment. *Prog Neurobiol* 33:309-333.
- Walz W (2000) Role of astrocytes in the clearance of excess extracellular potassium. *Neurochem Int* 36:291-300.

- Wang D, Sun X, Sadee W (2007) Different Effects of Opioid Antagonists on {micro}-, {delta}-, and {kappa}-Opioid Receptors with and without Agonist Pretreatment. *J Pharmacol Exp Ther* 321:544-552.
- Wang Q, Snell WJ (2003) Flagellar adhesion between mating type plus and mating type minus gametes activates a flagellar protein-tyrosine kinase during fertilization in *Chlamydomonas*. *J Biol Chem* 278:32936-32942.
- Wang X, Gu Z, Zhong P, Chen G, Feng J, Yan Z (2006) Aberrant regulation of NMDA receptors by dopamine D4 signaling in rats after phencyclidine exposure. *Mol Cell Neurosci* 31:15-25.
- Wang YQ, Sizeland A, Wang XF, Sassoon D (1995) Restricted expression of type-II TGF beta receptor in murine embryonic development suggests a central role in tissue modeling and CNS patterning. *Mech Dev* 52:275-289.
- Welch K, Friedman V (1960) The cerebrospinal fluid valves. *Brain* 83:454-469.
- Weller RO, Shulman K (1972) Infantile hydrocephalus: clinical, histological, and ultrastructural study of brain damage. *J Neurosurg* 36:255-265.
- Weller RO, Kida S, Zhang ET (1992) Pathways of fluid drainage from the brain--morphological aspects and immunological significance in rat and man. *Brain Pathol* 2:277-284.
- Weller RO, Mitchell J, Griffin RL, Gardner MJ (1978) The effects of hydrocephalus upon the developing brain. Histological and quantitative studies of the ependyma and subependyma in hydrocephalic rats. *J Neurol Sci* 36:383-402.
- Weller S, Gartner J (2001) Genetic and clinical aspects of X-linked hydrocephalus (L1 disease): Mutations in the L1CAM gene. *Hum Mutat* 18:1-12.
- Welty TE, Horner TG (1990) Pathophysiology and treatment of subarachnoid hemorrhage. *Clin Pharm* 9:35-39.
- Wenk GL (1984) Pharmacological manipulations of the substantia innominata-cortical cholinergic pathway. *Neurosci Lett* 51:99-103.
- Wessels MW, den Hollander NS, Willems PJ (2003) Mild fetal cerebral ventriculomegaly as a prenatal sonographic marker for Kartagener syndrome. *Prenat Diagn* 23:239-242.
- Whalen MJ, Carlos TM, Kochanek PM, Wisniewski SR, Bell MJ, Clark RS, DeKosky ST, Marion DW, Adelson PD (2000) Interleukin-8 is increased in cerebrospinal fluid of children with severe head injury. *Crit Care Med* 28:929-934.
- Whitelaw A (2001) Intraventricular streptokinase after intraventricular hemorrhage in newborn infants. *Cochrane Database Syst Rev*:CD000498.

- Whitelaw A, Christie S, Pople I (1999) Transforming growth factor-beta1: a possible signal molecule for posthemorrhagic hydrocephalus? *Pediatr Res* 46:576-580.
- Whitelaw A, Kennedy CR, Brion LP (2001) Diuretic therapy for newborn infants with posthemorrhagic ventricular dilatation. *Cochrane Database Syst Rev*:CD002270.
- Whittle IR, Johnston IH, Besser M (1985) Intracranial pressure changes in arrested hydrocephalus. *J Neurosurg* 62:77-82.
- Wickman KD, Clapham DE (1995) G-protein regulation of ion channels. *Curr Opin Neurobiol* 5:278-285.
- Wilkinson HA, Wilson RB, Patel PP, Esmaili M (1974) Corticosteroid therapy of experimental hydrocephalus after intraventricular-subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry* 37:224-229.
- Williams J, Lyttle S (1998) Mother and teacher reports of behaviour and perceived self-competence of children with hydrocephalus. *Eur J Pediatr Surg* 8 Suppl 1:5-9.
- Williams MA, Thomas G, de Lateur B, Imteyaz H, Rose JG, Shore WS, Kharkar S, Rigamonti D (2008) Objective assessment of gait in normal-pressure hydrocephalus. *Am J Phys Med Rehabil* 87:39-45.
- Wisniewski H, Weller RO, Terry RD (1969) Experimental hydrocephalus produced by the subarachnoid infusion of silicone oil. *J Neurosurg* 31:10-14.
- Witte MK, Stork JE, Blumer JL (1986) Diuretic therapeutics in the pediatric patient. *Am J Cardiol* 57:44A-53A.
- Wong EV, Kenwrick S, Willems P, Lemmon V (1995) Mutations in the cell adhesion molecule L1 cause mental retardation. *Trends Neurosci* 18:168-172.
- Woody CD (1986) Understanding the cellular basis of memory and learning. *Annu Rev Psychol* 37:433-493.
- Wozniak M, McLone DG, Raimondi AJ (1975) Micro- and macrovascular changes as the direct cause of parenchymal destruction in congenital murine hydrocephalus. *J Neurosurg* 43:535-545.
- Wright LC, McAllister JP, 2nd, Katz SD, Miller DW, Lovely TJ, Salotto AG, Wolfson BJ (1990) Cytological and cytoarchitectural changes in the feline cerebral cortex during experimental infantile hydrocephalus. *Pediatr Neurosurg* 16:139-155.
- Wyss-Coray T, Feng L, Masliah E, Ruppe MD, Lee HS, Toggas SM, Rockenstein EM, Mucke L (1995) Increased central nervous system production of extracellular matrix components and development of hydrocephalus in transgenic mice overexpressing transforming growth factor-beta 1. *Am J Pathol* 147:53-67.

- Xiuxin Liu AJBDMBOHAB (2006) GFAP-expressing cells in the postnatal subventricular zone display a unique glial phenotype intermediate between radial glia and astrocytes. *Glia* 54:394-410.
- Yamada H, Oi SZ, Tamaki N, Matsumoto S, Sudo K (1991) Prenatal aqueductal stenosis as a cause of congenital hydrocephalus in the inbred rat LEW/Jms. *Childs Nerv Syst* 7:218-222.
- Yamada H, Oi S, Tamaki N, Matsumoto S, Sudo K (1992) Histological changes in the midbrain around the aqueduct in congenital hydrocephalic rat LEW/Jms. *Childs Nerv Syst* 8:394-398.
- Yang J, Adamian M, Li T (2006) Rootletin interacts with C-Nap1 and may function as a physical linker between the pair of centrioles/basal bodies in cells. *Mol Biol Cell* 17:1033-1040.
- Yapicioglu H, Narli N, Satar M, Soyupak S, Altunbasak S (2003) Intraventricular streptokinase for the treatment of posthaemorrhagic hydrocephalus of preterm. *J Clin Neurosci* 10:297-299.
- Yasargil MG, Symon L, Teddy PJ (1984) Arteriovenous malformations of the spinal cord. *Adv Tech Stand Neurosurg* 11:61-102.
- Yashon D (1963) Prognosis in Infantile Hydrocephalus: Past and Present. *J Neurosurg* 20:105-111.
- Yau YH, Piper IR, Contant CF, Dunn LT, Whittle IR (2002) Clinical experience in the use of the Spiegelberg automated compliance device in the assessment of patients with hydrocephalus. *Acta Neurochir Suppl* 81:171-172.
- Yogev R, Davis AT (1980) Neurosurgical shunt infections. A review. *Childs Brain* 6:74-81.
- Yoshino M, Yoshimi Y, Taniguchi M, Nakamura S, Ikeda T (1999) Syndrome of inappropriate secretion of antidiuretic hormone associated with idiopathic normal pressure hydrocephalus. *Intern Med* 38:290-292.
- Young HF, Nulsen FE, Weiss MH, Thomas P (1973) The relationship of intelligence and cerebral mantle in treated infantile hydrocephalus. (IQ potential in hydrocephalic children). *Pediatrics* 52:38-44.
- Zakharov A, Papaiconomou C, Koh L, Djenic J, Bozanovic-Sosic R, Johnston M (2004) Integrating the roles of extracranial lymphatics and intracranial veins in cerebrospinal fluid absorption in sheep. *Microvasc Res* 67:96-104.
- Zammarchi E, Calzolari C, Pignotti MS, Pezzati P, Lignana E, Cama A (1993) Unusual presentation of the immotile cilia syndrome in two children. *Acta Paediatr* 82:312-313.

- Zechel J, Gohil H, Lust WD, Cohen A (2002) Alterations in matrix metalloproteinase-9 levels and tissue inhibitor of matrix metalloproteinases-1 expression in a transforming growth factor-beta transgenic model of hydrocephalus. *J Neurosci Res* 69:662-668.
- Zeiner HK, Prigatano GP, Pollay M, Biscoe CB, Smith RV (1985) Ocular motility, visual acuity and dysfunction of neuropsychological impairment in children with shunted uncomplicated hydrocephalus. *Childs Nerv Syst* 1:115-122.
- Zemack G, Romner B (2002) Adjustable valves in normal-pressure hydrocephalus: a retrospective study of 218 patients. *Neurosurgery* 51:1392-1400; discussion 1400-1392.
- Zhang D, Zeldin DC, Blackshear PJ (2007) Regulatory factor X4 variant 3: a transcription factor involved in brain development and disease. *J Neurosci Res* 85:3515-3522.
- Zhang J, Williams MA, Rigamonti D (2008) Heritable essential tremor-idiopathic normal pressure hydrocephalus (ETINPH). *Am J Med Genet A* 146A:433-439.
- Zhang Q, Davenport JR, Croyle MJ, Haycraft CJ, Yoder BK (2005) Disruption of IFT results in both exocrine and endocrine abnormalities in the pancreas of Tg737(orpk) mutant mice. *Lab Invest* 85:45-64.